

A.I. ESHNIYOZOV

OLIV MATEMATIKADAN MUSTAQIL ISHLAR

$$\begin{aligned} \frac{\partial}{\partial a} \ln f_{a, \sigma^2}(\xi_1) &= \frac{(\xi_1 - a)}{\sigma^2} f_{a, \sigma^2}(\xi_1) = \frac{1}{\sqrt{2\pi\sigma}} \exp\left\{-\frac{(\xi_1 - a)^2}{2\sigma^2}\right\} \\ \int_{R_n} T(x) \cdot \frac{\partial}{\partial \theta} f(x, \theta) dx &= M\left(T(\xi) \cdot \frac{\partial}{\partial \theta} \ln L(\xi, \theta)\right) \\ \int_{R_n} T(x) \cdot \left(\frac{\partial}{\partial \theta} \ln L(x, \theta)\right) \cdot f(x, \theta) dx &= \int_{R_n} T(x) \cdot \left(\frac{\frac{\partial}{\partial \theta} f(x, \theta)}{f(x, \theta)}\right) \cdot f(x, \theta) dx \\ \frac{\partial}{\partial \theta} \int_{R_n} T(x) f(x, \theta) dx &= \int_{R_n} \frac{\partial}{\partial \theta} T(x) f(x, \theta) dx \end{aligned}$$

**O‘ZBEKISTON RESPUBLIKASI OLIY VA O‘RTA
MAXSUS TA‘LIM VAZIRLIGI**

ESHNIYOZOV A.I.

**OLIV MATEMATIKADAN
MUSTAQIL ISHLAR**

O‘quv qo‘llanma

Guliston-2022

UO'K 514.124, 517.2/3.

A.I.Eshniyozov. Oliy matematikadan mustaqil ishlar. Amaliy mashg'ulotlar uchun o'quv qo'llanma – Guliston., 2022, 252 bet

O'quv qo'llanmada analitik geometriya, limitlar nazariyasi, bir o'zgaruvchi funksiyalarning differensial va integral hisobi va ularning tatbiqlari bo'limlari qaralgan. Har bir mavzu uchun qisqacha nazariy ma'lumotlar, namunaviy misol va masalalar to'liq echimi bilan keltirilgan, mustaqil o'zlashtirish uchun berilgan misol va masalalarning javoblari berilgan.

Oliy o'quv yurtlari talabalari uchun mo'ljallangan.

В пособие рассматриваются основные разделы аналитическая геометрия, теория пределов, дифференциальное и интегральное исчисления функций с одной переменных и их применение. По каждой теме кратко излагается основные теоретические сведения, детально разобраны типовые задачи, даются задачи и упражнения для самостоятельной работы с ответами.

Для студентов высших учебных заведений.

In a manual examined basic divisions analytical geometry, theory of limits, differential and integral calculations of functions with one the variables and their application. On every topic briefly expounded basic theoretical information, model tasks are taken apart in detail, tasks and exercises are given for independent work with answers.

For the students of higher educational establishments.

Taqrizchilar :

O'zbekiston Milliy Universitet professori,
fizika-matematika fanlari doktori **R.N. Ganixo'jayev**
Guliston Davlat Universiteti dotsenti, fizika-
matematika fanlari nomzodi **H. Norjigitov**

SO'Z BOSHI

Hozirgi zamon ilmiy – texnika taraqqiyoti sharoitida Oliy o'quv yurtlarida yuqori malakali mutaxassislar tayyorlash borasida fizika – matematika fanlariga katta e'tibor berilayapti.

Oliy matematika kursi bo'yicha chuqur va har tomonlama bilim egallash uchun faqat asosiy nazariy materialning o'zi yetarli bo'lmasdan, maxsus tanlangan misol va masalalarni yetarlicha yechish ham zarur bo'ladi.

Bunda masala va misollarning aniq matematik ko'yilishi, yechimlarning asoslanganligi va to'laligi, javoblarning to'g'riligi katta ahamiyatga egadir.

E'tiboringizga havola qilinayotgan mazkur qo'llanma Oliy o'quv yurtlarining bakalavr yo'nalishidagi Oliy matematika dasturiga moslab tuzilgan.

Qo'llanmaning har qaysi bobida qisqacha nazariy ma'lumot bayon qilinib, tipik masalalar to'liq yechimlari berilgan. Bulardan tashqari, har qaysi paragraf oxirida mustaqil yechish uchun masalalar va ulaning javoblari keltirilgan. Ana shu misol va masalalarni yechish bilangina – cheklanib qolish yaramaydi, zero ushbu qo'llanma tegishli to'plamlarning o'rnini bosishga da'vo – qilmaydi.

Qo'llanma oliy o'quv yurtlarining barcha bakalar yo'nalishidagi talabalari uchun mo'ljallangan.

Muallif qimmatli metodik maslahatlari uchun UzMU, mexanika – matematika fakulteti professori, fizika – matematika fanlari doktori R.N.G'anixo'jayevga, GulDU fizika – matematika fakulteti "Umumiy matematika" kafedrasi dotsenti, fizika – matematika fanlari nomzodi K.Jomuratovga, hamda "Umumiy matematika" kafedrasi mudiri dotsent, fizika – matematika fanlari nomzodi X. Norjigitovga o'zining chuqur minnatdorchiligini bildiradi.

Muallif.

I BOB. Analitik geometriya

Analitik geometriya bobida siz analitik geometriyaning har xil masalalari: to'g'ri chiziq, tekislik tenglamalari va ularga oid masalalar, vektorlar ustida amallar, vektorlarning skalyar, vektor va aralash ko'paytmalari, geometrik masalalarga ularning tatbiqlari o'rganib olasiz.

Vektorni bazis bo'yicha yoyilmasi

Uchta chiziqli bog'liqmas vektorlar sistemasi p, q, r berilgan bo'lib, agar ixtiyoriy x vektorni ularning chiziqli kombinatsiya, ya'ni

$$\vec{x} = a \vec{\times p} + b \vec{\times q} + g \vec{\times r}$$

shaklida ifodalash mumkin bo'lsa, u holda berilgan sistema bazis deyiladi.

Bu tenglik x vektorning p, q, r bazis bo'yicha yoyilmasi deyiladi.

\vec{x} vektorning yoyilmasi quyidagi ko'rinishda izlanadi:

$$\vec{x} = a \vec{\times p} + b \vec{\times q} + g \vec{\times r}.$$

Bu tenglama a, b, g larga nisbatan vektor tenglama bo'lib, uch o'zgaruvchili uchta chiziqli tenglamalar sistemasi yordamida quyidagicha yoziladi:

$$\begin{cases} a \times p_1 + b \times q_1 + g \times r_1 = x_1 \\ a \times p_2 + b \times q_2 + g \times r_2 = x_2 \\ a \times p_3 + b \times q_3 + g \times r_3 = x_3 \end{cases}$$

Tenglamalar sistemasini yechib a, b, g larni topib,

$$\vec{x} = a \vec{\times p} + b \vec{\times q} + g \vec{\times r}$$

vektorni ko'rinishini topamiz.

1-masala. x vektorni p, q, r vektorlar orqali yoyilmasini yozing.

$$x = \{-13, 2, 18\},$$

$$p = \{1, 1, 4\},$$

$$q = \{-3, 0, 2\},$$

$$r = \{1, 2, -1\}.$$

Echim:

x vektorni p, q, r vektorlar orqali yoyilmasi: $x = a \succ p + b \succ q + g \succ r$.

yoki sistema ko'rinishida

$$\begin{cases} a \succ p_1 + b \succ q_1 + g \succ r_1 = x_1 \\ a \times p_2 + b \times q_2 + g \times r_2 = x_2 \\ a \times p_3 + b \times q_3 + g \times r_3 = x_3 \end{cases}$$

Natijada

$$\begin{cases} a - 3b + g = -13, \\ a + 2g = 2, & | \times 2 \\ 4a + 2b - g = 18 \end{cases}$$

$$\begin{cases} a - 3b + g = -13, \\ -a + 6g = 28, \\ 4a + 2b - g = 18 \end{cases}$$

Uchinchi satrga birinchi satrni qo'shib:

$$\begin{cases} a - 3b + g = -13, \\ -a + 6g = 28, \\ 5a - b = 5 \end{cases}$$

$$\begin{cases} a - 3b + g = -13, \\ -a + 6g = 28, \\ 5a - b = 5 & | \times 6 \end{cases}$$

$$\begin{cases} a - 3b + g = -13, \\ 29a = 58, \\ 5a - b = 5 \end{cases}$$

$$\begin{cases} a - 3b + g = -13, \\ a = 2, \\ 5a - b = 5 \end{cases}$$

$$\begin{cases} a - 3b + g = -13, \\ a = 2, \\ 5 \times 2 - b = 5 \end{cases}$$

$$\begin{cases} 2 - 3b + g = -13, \\ a = 2, \\ b = 5 \end{cases}$$

$$\begin{cases} -3b + g = -15, \\ a = 2, \\ b = 5 \end{cases}$$

$$\begin{cases} -3 \times 5 + g = -15, \\ a = 2, \\ b = 5 \end{cases}$$

$$\begin{cases} a = 2, \\ b = 5, \\ g = 0. \end{cases}$$

Izlanayotgan yoyilma: $x = 2p + 5q$ ko'rinishida bo'lar ekan.

1–masala. x vektorni p , q , r vektorlar orqali yoyilmasini yozing.

- | | | |
|---|---|---|
| $x = \{-2, 4, 7\},$
$p = \{0, 1, 2\},$
^{1.} $q = \{1, 0, 1\},$
$r = \{-1, 2, 4\}.$ | $x = \{6, 12, -1\},$
$p = \{1, 3, 0\},$
^{2.} $q = \{2, -1, 1\},$
$r = \{0, -1, 2\}.$ | $x = \{1, -4, 4\},$
$p = \{2, 1, -1\},$
^{3.} $q = \{0, 3, 2\},$
$r = \{1, -1, 1\}.$ |
| $x = \{-9, 5, 5\},$
$p = \{4, 1, 1\},$
^{4.} $q = \{2, 0, -3\},$
$r = \{-1, 2, 1\}.$ | $x = \{-5, -5, 5\},$
$p = \{-2, 0, 1\},$
^{5.} $q = \{1, 3, -1\},$
$r = \{0, 4, 1\}.$ | $x = \{13, 2, 7\},$
$p = \{5, 1, 0\},$
^{6.} $q = \{2, -1, 3\},$
$r = \{1, 0, -1\}.$ |
| $x = \{-19, -1, 7\},$
$p = \{0, 1, 1\},$
^{7.} $q = \{-2, 0, 1\},$
$r = \{3, 1, 0\}.$ | $x = \{3, -3, 4\},$
$p = \{1, 0, 2\},$
^{8.} $q = \{0, 1, 1\},$
$r = \{2, -1, 4\}.$ | $x = \{3, 3, -1\},$
$p = \{3, 1, 0\},$
^{9.} $q = \{-1, 2, 1\},$
$r = \{-1, 0, 2\}.$ |
| $x = \{-1, 7, -4\},$
$p = \{-1, 2, 1\},$
^{10.} $q = \{2, 0, 3\},$
$r = \{1, 1, -1\}.$ | $x = \{6, 5, -14\},$
$p = \{1, 1, 4\},$
^{11.} $q = \{0, -3, 2\},$
$r = \{2, 1, -1\}.$ | $x = \{6, -1, 7\},$
$p = \{1, -2, 0\},$
^{12.} $q = \{-1, 1, 3\},$
$r = \{1, 0, 4\}.$ |
| $x = \{5, 15, 0\},$
$p = \{1, 0, 5\},$
^{13.} $q = \{-1, 3, 2\},$
$r = \{0, -1, 1\}.$ | $x = \{2, -1, 11\},$
$p = \{1, 1, 0\},$
^{14.} $q = \{0, 1, -2\},$
$r = \{1, 0, 3\}.$ | $x = \{11, 5, -3\},$
$p = \{1, 0, 2\},$
^{15.} $q = \{-1, 0, 1\},$
$r = \{2, 5, -3\}.$ |
| $x = \{8, 0, 5\},$
$p = \{2, 0, 1\},$
^{16.} $q = \{1, 1, 0\},$
$r = \{4, 1, 2\}.$ | $x = \{3, 1, 8\},$
$p = \{0, 1, 3\},$
^{17.} $q = \{1, 2, -1\},$
$r = \{2, 0, -1\}.$ | $x = \{8, 1, 12\},$
$p = \{1, 2, -1\},$
^{18.} $q = \{3, 0, 2\},$
$r = \{-1, 1, 1\}.$ |

$$\begin{array}{lll}
 x = \{-9, -8, -3\}, & x = \{-5, 9, -13\}, & x = \{-15, 5, 6\}, \\
 p = \{1, 4, 1\}, & p = \{0, 1, -2\}, & p = \{0, 5, 1\}, \\
 19. \quad q = \{-3, 2, 0\}, & 20. \quad q = \{3, -1, 1\}, & 21. \quad q = \{3, 2, -1\}, \\
 r = \{1, -1, 2\}. & r = \{4, 1, 0\}. & r = \{-1, 1, 0\}.
 \end{array}$$

$$\begin{array}{lll}
 x = \{8, 9, 4\}, & x = \{23, -14, -30\}, & x = \{3, 1, 3\}, \\
 p = \{1, 0, 1\}, & p = \{2, 1, 0\}, & p = \{2, 1, 0\}, \\
 22. \quad q = \{0, -2, 1\}, & 23. \quad q = \{1, -1, 0\}, & 24. \quad q = \{1, 0, 1\}, \\
 r = \{1, 3, 0\}. & r = \{-3, 2, 5\}. & r = \{4, 2, 1\}.
 \end{array}$$

$$\begin{array}{lll}
 x = \{1, 7, 0\}, & x = \{11, -1, 4\}, & x = \{0, -8, 9\}, \\
 p = \{0, 3, 1\}, & p = \{1, -1, 2\}, & p = \{0, -2, 1\}, \\
 25. \quad q = \{1, -1, 2\}, & 26. \quad q = \{3, 2, 0\}, & 27. \quad q = \{3, 1, -1\}, \\
 r = \{2, -1, 0\}. & r = \{-1, 1, 1\}. & r = \{4, 0, 1\}.
 \end{array}$$

$$\begin{array}{lll}
 x = \{8, -7, -13\}, & x = \{2, 7, 5\}, & x = \{-15, -20, -1\}, \\
 p = \{0, 1, 5\}, & p = \{1, 0, 1\}, & p = \{0, 2, 1\}, \\
 28. \quad q = \{3, -1, 2\}, & 29. \quad q = \{1, -2, 0\}, & 30. \quad q = \{0, 1, -1\}, \\
 r = \{-1, 0, 1\}. & r = \{0, 3, 1\}. & r = \{5, -3, 2\}.
 \end{array}$$

Javoblar. 1.1 $x = 2p - q + r$; 1.2 $x = 4p + q - r$; 1.3 $x = -p + 3r$; 1.4 $x = -p - q + 3r$;
 1.5 $x = p - 3q + r$; 1.6 $x = 3p + q - 4r$; 1.7 $x = 2p + 5q - 3r$; 1.8 $x = p - 2q + r$;
 1.9 $x = -p + q - r$; 1.10 $x = 2p - q + 3r$; 1.11 $x = -2p - q + 4r$;
 1.12 $x = -p - 3q + 4r$; 1.13 $x = 4p - q - 18r$; 1.14 $x = -3p + 2q + 5r$;
 1.15 $x = 3p - 6q + r$; 1.16 $x = p - 2q + 2r$; 1.17 $x = 3p - q + 2r$;
 1.18 $x = -p + 4q + 2,2r$; 1.19 $x = -3p + 2q$; 1.20 $x = 5p - 3q + r$ 1.21 $x = 2p - 4q + 3r$;
 1.22 $x = 7p - 3q + r$; 1.23 $x = 13p + 15q - 6r$;
 1.24 $x = -3p + q + 2r$; 1.25 $x = 2p - q$; 1.26 $x = 3p + 2q - 2r$; 1.27 $x = 2p - 4q + 3r$; 1.28
 $x = -4p + 3q + r$; 1.29 $x = 4p - 2q + r$; 1.30 $x = -6p + q + 3r$.

Vektorlarning kollinearligi

Bitta to'g'ri chiziqda yoki parallel chiziqlarda yotgan c_1 va c_2 vektorlar kollinear vektorlar deyiladi. Boshqacha aytganda shunday a topilsaki, $\vec{p} = a\vec{q}$ bo'lsa, ya'ni ularning koordinatalari o'zaro proporsional bo'lishi zarur va yetarli bo'ladi.

Demak,

$$\frac{p_1}{q_1} = \frac{p_2}{q_2} = \frac{p_3}{q_3}$$

bo'lsa, kollinear, tenglik bajarilmasa \vec{p} va \vec{q} vektorlar kollinear emas bo'lar ekan.

2-masala. a va b yordamida qurilgan c_1 va c_2 vektorlar kollinear mi?

$$a = \{-1, 2, -1\}, \quad b = \{2, -7, 1\}, \quad c_1 = 6a - 2b, \quad c_2 = b - 3a.$$

$$c_1 = 6a - 2b = \{6 \cdot (-1) - 2 \cdot 2; 6 \cdot 2 - 2 \cdot (-7); 1 - 6 \cdot (-1) - 2 \cdot 1\} = \{-10, 26, -8\}.$$

$$c_2 = b - 3a = \{2 - 3 \cdot (-1); -7 - 3 \cdot 2; 1 - 3 \cdot (-1)\} = \{5, -13, 4\}.$$

$$\frac{-10}{-5} = \frac{26}{-13} = \frac{-8}{4}, \text{ ya'ni } c_1 = -2c_2.$$

Demak, c_1 va c_2 kollinear ekan.

$$1. a = \{1, -2, 3\}, b = \{3, 0, -1\}, c_1 = 2a + 4b, c_2 = 3b - a.$$

$$2. a = \{1, 0, 1\}, b = \{-2, 3, 5\}, c_1 = a + 2b, c_2 = 3a - b.$$

$$3. a = \{-2, 4, 1\}, b = \{1, -2, 7\}, c_1 = 5a + 3b, c_2 = 2a - b.$$

$$4. a = \{1, 2, -3\}, b = \{2, -1, -1\}, c_1 = 4a + 3b, c_2 = 8a - b.$$

$$5. a = \{3, 5, 4\}, b = \{5, 9, 7\}, c_1 = -2a + b, c_2 = 3a - 2b.$$

$$6. a = \{1, 4, -2\}, b = \{1, 1, -1\}, c_1 = a + b, c_2 = 4a + 2b.$$

$$7. a = \{1, -2, 5\}, b = \{3, -1, 0\}, c_1 = 4a - 2b, c_2 = b - 2a.$$

$$8. a = \{3, 4, -1\}, b = \{2, -1, 1\}, c_1 = 6a - 3b, c_2 = b - 2a.$$

$$9. a = \{-2, -3, -2\}, b = \{1, 0, -5\}, c_1 = 3a + 9b, c_2 = -a - 3b.$$

$$10. a = \{-1, 4, 2\}, b = \{3, -2, 6\}, c_1 = 2a - b, c_2 = 3b - 6a.$$

$$11. a = \{5, 0, -1\}, b = \{7, 2, 3\}, c_1 = 2a - b, c_2 = 3b - 6a.$$

$$12. a = \{0, 3, -2\}, b = \{1, -2, 1\}, c_1 = 5a - 2b, c_2 = 3a + 5b.$$

$$13. a = \{-2, 7, -1\}, b = \{-3, 5, 2\}, c_1 = 2a + 3b, c_2 = 3a + 2b.$$

$$14. a = \{3, 7, 0\}, b = \{1, -3, 4\}, c_1 = 4a - 2b, c_2 = b - 2a.$$

$$15. a = \{7, 9, -2\}, b = \{5, 4, 3\}, c_1 = 4a - b, c_2 = 4b - a.$$

$$16. a = \{5, 0, -2\}, b = \{6, 4, 3\}, c_1 = 5a - 3b, c_2 = 6b - 10a.$$

$$17. a = \{8, 3, -1\}, b = \{4, 1, 3\}, c_1 = 2a - b, c_2 = 2b - 4a.$$

$$18. a = \{3, -1, 6\}, b = \{5, 7, 10\}, c_1 = 4a - 2b, c_2 = b - 2a.$$

$$19. a = \{1, -2, 4\}, b = \{7, 3, 5\}, c_1 = 6a - 3b, c_2 = b - 2a.$$

$$20. a = \{3, 7, 0\}, b = \{4, 6, -1\}, c_1 = 3a + 2b, c_2 = 5a - 7b.$$

$$21. a = \{2, -1, 4\}, b = \{3, -7, -6\}, c_1 = 2a - 3b, c_2 = 3a - 2b.$$

$$22. a = \{5, -1, -2\}, b = \{6, 0, 7\}, c_1 = 3a - 2b, c_2 = 4b - 6a.$$

$$23. a = \{-9, 5, 3\}, b = \{7, 1, -2\}, c_1 = 2a - b, c_2 = 3a + 5b.$$

$$24. a = \{4, 2, 9\}, b = \{0, -1, 3\}, c_1 = 4b - 3a, c_2 = 4a - 3b.$$

$$25. a = \{2, -1, 6\}, b = \{-1, 3, 8\}, c_1 = 5a - 2b, c_2 = 2a - 5b.$$

$$26. a = \{5, 0, 8\}, b = \{-3, 1, 7\}, c_1 = 3a - 4b, c_2 = 12b - 9a.$$

$$27. a = \{-1, 3, 4\}, b = \{2, -1, 0\}, c_1 = 6a - 2b, c_2 = b - 3a.$$

$$28. a = \{4, 2, -7\}, b = \{5, 0, -3\}, c_1 = a - 3b, c_2 = 6b - 2a.$$

$$29. a = \{2, 0, -5\}, b = \{1, -3, 4\}, c_1 = 2a - 5b, c_2 = 5a - 2b.$$

$$30. a = \{-1, 2, 8\}, b = \{3, 7, -1\}, c_1 = 4a - 3b, c_2 = 9b - 12a.$$

Javoblar. 2.1 yo'q; 2.2 yo'q; 2.3 yo'q; 2.4 yo'q; 2.5 yo'q; 2.6 yo'q; 2.7 ha; 2.8 ha; 2.9 ha; 2.10 ha; 2.11 ha; 2.12 yo'q; 2.13 yo'q; 2.14 ha; 2.15 yo'q; 2.16 ha; 2.17 ha; 2.18 ha; 2.19 ha; 2.20 yo'q; 2.21 yo'q; 2.22 ha; 2.23 yo'q; 2.24 yo'q; 2.25 yo'q; 2.26 ha; 2.27 ha; 2.28 yo'q; 2.29 yo'q; 2.30 ha.

Vektorlar orasidagi burchak

Ikkita \overrightarrow{AB} va \overrightarrow{AC} vektorning skalyar ko'paytmasi deb, $(\overrightarrow{AB}, \overrightarrow{AC})$ ko'rinishda belgilanuvchi va shu vektorlar uzunliklari ko'paytmasining ular orasidagi burchak kosinusi bilan ko'paytmasiga teng bo'lgan songa aytiladi:

$$(\overrightarrow{AB}, \overrightarrow{AC}) = |\overrightarrow{AB}| \times |\overrightarrow{AC}| \times \cos j.$$

$A(x_1, y_1, z_1)$, $B(x_2, y_2, z_2)$ va $C(x_3, y_3, z_3)$ berilganda vektorlarning skalyar ko'paytmasi

$$\overrightarrow{AB} = (x_2 - x_1, y_2 - y_1, z_2 - z_1)$$

$$\overrightarrow{AC} = (x_3 - x_1, y_3 - y_1, z_3 - z_1)$$

$$(\overrightarrow{AB}, \overrightarrow{AC}) = (x_2 - x_1)(x_3 - x_1) + (y_2 - y_1)(y_3 - y_1) + (z_2 - z_1)(z_3 - z_1)$$

va vektorlarning uzunliklari mos ravishda:

$$|\overrightarrow{AB}| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2},$$

$$|\overrightarrow{AC}| = \sqrt{(x_3 - x_1)^2 + (y_3 - y_1)^2 + (z_3 - z_1)^2},$$

formulalar bilan topiladi.

\overrightarrow{AB} va \overrightarrow{AC} vektor orasidagi burchak ushbu formula bilan hisoblanadi:

$$\cos j = \frac{(\overrightarrow{AB}, \overrightarrow{AC})}{|\overrightarrow{AB}| \times |\overrightarrow{AC}|}.$$

3–masala. \overrightarrow{AB} va \overrightarrow{AC} vektorlar orasidagi burchak kosinusini toping.

$$A(1, -2, 3),$$

$$B(3, 4, -6),$$

$$C(1, 1, -1).$$

$$\overline{AB} = \{4, 2, -3\}, \quad |\overline{AB}| = \sqrt{4^2 + 2^2 + (-3)^2} = \sqrt{29}.$$

$$\overline{AC} = \{2, -1, 2\}, \quad |\overline{AC}| = \sqrt{2^2 + (-1)^2 + (2)^2} = 3.$$

$$\cos(\overline{AB} \wedge \overline{AC}) = \frac{4 \times 2 - 2 \times 1 - 3 \times 2}{3 \times \sqrt{29}} = 0.$$

$$(\overline{AB} \wedge \overline{AC}) = \frac{\rho}{2}.$$

$$A(1, -2, 3),$$

$$1. B(0, -1, 2),$$

$$C(3, -4, 5).$$

$$A(0, -3, 6),$$

$$2. B(-12, -3, -3),$$

$$C(-9, -3, -6).$$

$$A(3, 3, -1),$$

$$3. B(5, 5, -2),$$

$$C(4, 1, 1).$$

$$A(-4, -2, 0),$$

$$4. B(-1, -2, 4),$$

$$C(3, -2, 1).$$

$$A(5, 3, -1),$$

$$5. B(5, 2, 0),$$

$$C(6, 4, -1).$$

$$A(-3, -7, -5),$$

$$6. B(0, -1, -2),$$

$$C(2, 3, 0).$$

$$A(2, -4, 6),$$

$$7. B(0, -2, 4),$$

$$C(6, -8, 10).$$

$$A(0, 1, -2),$$

$$8. B(3, 1, 2),$$

$$C(4, 1, 1).$$

$$A(3, 3, -1),$$

$$9. B(1, 5, -2),$$

$$C(4, 1, 1).$$

$$A(2, 1, -1),$$

$$10. B(6, -1, -4),$$

$$C(4, 2, 1).$$

$$A(-1, -2, 1),$$

$$11. B(-4, -2, 5),$$

$$C(-8, -2, 2).$$

$$A(6, 2, -3),$$

$$12. B(6, 3, -2),$$

$$C(7, 3, -3).$$

$$A(0, 0, 4),$$

$$13. B(-3, -6, 1),$$

$$C(-5, -10, -1).$$

$$A(2, -8, -1),$$

$$14. B(4, -6, 0),$$

$$C(-2, -5, -1).$$

$$A(3, -6, 9),$$

$$15. B(0, -3, 6),$$

$$C(9, -12, 15).$$

- $A(0, 2, -4),$ $A(3, 3, -1),$ $A(-4, 3, 0),$
 16. $B(8, 2, 2),$ 17. $B(5, 1, -2),$ 18. $B(0, 1, 3),$
 $C(6, 2, 4).$ $C(4, 1, 1).$ $C(-2, 4, -2).$
- $A(1, -1, 0),$ $A(7, 0, 2),$ $A(2, 3, 2),$
 19. $B(-2, -1, 4),$ 20. $B(7, 1, 3),$ 21. $B(-1, -3, -1),$
 $C(8, -1, -1).$ $C(8, -1, 2).$ $C(-3, -7, -3).$
- $A(2, 2, 7),$ $A(-1, 2, -3),$ $A(0, 3, -6),$
 22. $B(0, 0, 6),$ 23. $B(0, 1, -2),$ 24. $B(9, 3, 6),$
 $C(-2, 5, 7).$ $C(-3, 4, -5).$ $C(12, 3, 3).$
- $A(3, 3, -1),$ $A(-2, 1, 1),$ $A(1, 4, -1),$
 25. $B(5, 1, -2),$ 26. $B(2, 3, -2),$ 27. $B(-2, 4, -5),$
 $C(4, 1, -3).$ $C(0, 0, 3).$ $C(8, 4, 0).$
- $A(0, 1, 0),$ $A(-4, 0, 4),$ $A(-2, 4, -6),$
 28. $B(0, 2, 1),$ 29. $B(-1, 6, 7),$ 30. $B(0, 2, -4),$
 $C(1, 2, 0).$ $C(1, 10, 9).$ $C(-6, 8, -10).$

Javoblar. 3.1 p ; 3.2 $16^0 15' 37''$; 3.3 $116^0 23' 16''$; 3.4 $\frac{p}{4}$; 3.5 $\frac{2p}{3}$; 3.6 p ; 3.7 p ;
 3.8 $\arccos 0,96$; 3.9 $152^0 44' 2''$; 3.10 $\frac{p}{2}$; 3.11 $\frac{p}{4}$; 3.12 $\frac{p}{3}$; 3.13 0^0 ; 3.14 $97^0 39' 44''$;
 3.15 p ; 3.16 $16^0 15' 37''$; 3.17 $63^0 36' 44''$; 3.18 $\frac{p}{2}$; 3.19 135^0 ; 3.20 120^0 ; 3.21 0^0 ;
 3.22 $82^0 20' 15''$; 3.23 p ; 3.24 $16^0 15' 37''$; 3.25 $27^0 15' 58''$; 3.26 $\frac{p}{2}$; 3.27 135^0 ; 3.28 $\frac{p}{3}$;
 3.29 0^0 ; 3.30 p .

Parallelogrammning yuzi

a va b vektorlarga qurilgan parallelogramm yuzi

$$S = |\vec{a}, \vec{b}|$$

shu vektorlarning vektor ko'paytmasidan olingan modulga teng.

$$[\vec{a}, \vec{b}] = [a_1 \vec{p} + a_2 \vec{q}, b_1 \vec{p} + b_2 \vec{q}] = a_1 b_1 [\vec{p}, \vec{p}] + a_1 b_2 [\vec{p}, \vec{q}] + a_2 b_1 [\vec{q}, \vec{p}] + a_2 b_2 [\vec{q}, \vec{q}] = (a_1 b_2 - a_2 b_1) [\vec{p}, \vec{q}].$$

Bundan,

$$S = |\vec{a}, \vec{b}| = (a_1 b_2 - a_2 b_1) [\vec{p}, \vec{q}] \sin j.$$

4-masala. a va b vektorlarga qurilgan parallelogramm yuzini hisoblang.

$$a = 6p - q,$$

$$b = 5q + p.$$

$$|p| = \frac{1}{2}, \quad |q| = 4, \quad (p \wedge q) = \frac{5p}{6}.$$

$$\begin{aligned} S &= |(6p - q) \wedge (5q + p)| = |6p \wedge 5q + 6p \wedge p - 5q \wedge q - q \wedge p| = \\ &= |6p \wedge 5q + p \wedge q| = 31|p| \times |q| \sin(p \wedge q) = \\ &= 31 \times \frac{1}{2} \times 4 \times \sin \frac{5p}{6} = 31 \times 2 \times \frac{1}{2} = 31. \end{aligned}$$

1. $a = p + 2q, \quad b = 3p - q; \quad |p| = 1, \quad |q| = 2, \quad (p \wedge q) = p/6.$
2. $a = 3p + q, \quad b = p - 2q; \quad |p| = 4, \quad |q| = 1, \quad (p \wedge q) = p/4.$
3. $a = p - 3q, \quad b = p + 2q; \quad |p| = 1/5, \quad |q| = 1, \quad (p \wedge q) = p/2.$
4. $a = 3p - 2q, \quad b = p + 5q; \quad |p| = 4, \quad |q| = 1/2, \quad (p \wedge q) = 5p/6.$
5. $a = p - 2q, \quad b = 2p + q; \quad |p| = 2, \quad |q| = 3, \quad (p \wedge q) = 3p/4.$
6. $a = p + 3q, \quad b = p - 2q; \quad |p| = 2, \quad |q| = 3, \quad (p \wedge q) = p/3.$
7. $a = 2p - q, \quad b = p + 3q; \quad |p| = 3, \quad |q| = 2, \quad (p \wedge q) = p/2.$
8. $a = 4p + q, \quad b = p - q; \quad |p| = 7, \quad |q| = 2, \quad (p \wedge q) = p/4.$
9. $a = p - 4q, \quad b = 3p + q; \quad |p| = 1, \quad |q| = 2, \quad (p \wedge q) = p/6.$
10. $a = p + 4q, \quad b = 2p - q; \quad |p| = 7, \quad |q| = 2, \quad (p \wedge q) = p/3.$

11. $a = 3p + 2q, \quad b = p - q; \quad |p| = 10, \quad |q| = 1, \quad (p \wedge q) = p/2.$
12. $a = 4p - q, \quad b = p + 2q; \quad |p| = 5, \quad |q| = 4, \quad (p \wedge q) = p/4.$
13. $a = 2p + 3q, \quad b = p - 2q; \quad |p| = 6, \quad |q| = 7, \quad (p \wedge q) = p/3.$
14. $a = 3p - q, \quad b = p + 2q; \quad |p| = 3, \quad |q| = 4, \quad (p \wedge q) = p/3.$
15. $a = 2p + 3q, \quad b = p - 2q; \quad |p| = 2, \quad |q| = 3, \quad (p \wedge q) = p/4.$
16. $a = 2p - 3q, \quad b = 3p + q; \quad |p| = 4, \quad |q| = 1, \quad (p \wedge q) = p/6.$
17. $a = 5p + q, \quad b = p - 3q; \quad |p| = 1, \quad |q| = 2, \quad (p \wedge q) = p/3.$
18. $a = 7p - 2q, \quad b = p + 3q; \quad |p| = 1/2, \quad |q| = 2, \quad (p \wedge q) = p/2.$
19. $a = 6p - q, \quad b = p + q; \quad |p| = 3, \quad |q| = 4, \quad (p \wedge q) = p/4.$
20. $a = 10p + q, \quad b = 3p - 2q; \quad |p| = 4, \quad |q| = 1, \quad (p \wedge q) = p/6.$
21. $a = 6p - q, \quad b = p + 2q; \quad |p| = 8, \quad |q| = 1/2, \quad (p \wedge q) = p/3.$
22. $a = 3p + 4q, \quad b = q - p; \quad |p| = 2,5, \quad |q| = 2, \quad (p \wedge q) = p/2.$
23. $a = 7p + q, \quad b = p - 3q; \quad |p| = 3, \quad |q| = 1, \quad (p \wedge q) = 3p/4.$
24. $a = p + 3q, \quad b = 3p - q; \quad |p| = 3, \quad |q| = 5, \quad (p \wedge q) = 2p/3.$
25. $a = 3p + q, \quad b = p - 3q; \quad |p| = 7, \quad |q| = 2, \quad (p \wedge q) = p/4.$
26. $a = 5p - q, \quad b = p + q; \quad |p| = 5, \quad |q| = 3, \quad (p \wedge q) = 5p/6.$
27. $a = 3p - 4q, \quad b = p + 3q; \quad |p| = 2, \quad |q| = 3, \quad (p \wedge q) = p/4.$
28. $a = 2p + 3q, \quad b = p - 2q; \quad |p| = 2, \quad |q| = 1, \quad (p \wedge q) = p/3.$
29. $a = 2p - 3q, \quad b = 5p + q; \quad |p| = 2, \quad |q| = 3, \quad (p \wedge q) = p/2.$
30. $a = 3p + 2q, \quad b = 2p - q; \quad |p| = 4, \quad |q| = 4, \quad (p \wedge q) = 3p/4.$

Javoblar. 4.1 7; 4.2 $14\sqrt{2}$; 4.3 1; 4.4 17; 4.5 $15\sqrt{2}$; 4.6 $15\sqrt{3}$; 4.7 42; 4.8 $35\sqrt{2}$; 4.9 13; 4.10 $63\sqrt{3}$; 4.11 50; 4.12 $90\sqrt{2}$; 4.13 $147\sqrt{3}$; 4.14 $42\sqrt{3}$; 4.15 $21\sqrt{2}$; 4.16 22; 4.17 $16\sqrt{3}$; 4.18 23; 4.19 $42\sqrt{2}$; 4.20 46; 4.21 $26\sqrt{3}$; 4.22 35; 4.23 $33\sqrt{2}$; 4.24 $75\sqrt{3}$; 4.25 $70\sqrt{2}$; 4.26 45; 4.27 $39\sqrt{2}$; 4.28 $7\sqrt{3}$; 4.29 102; 4.30 $42\sqrt{2}$.

Vektorlar komplanarligi

Bir tekislikda yoki parallel tekisliklarda yotuchi vektorlarni

komplanar vektorlar deyiladi.

Agar $a = (a_1, a_2, a_3)$, $b = (b_1, b_2, b_3)$ va $c = (c_1, c_2, c_3)$ vektorlar komplanar bo'lishi uchun ularning aralash ko'paytmasi nolga teng bo'lishi zarur va yetarli, ya'ni

$$(\vec{a}, \vec{b}, \vec{c}) = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = 0.$$

5–masala. a , b va c vektorlar komplanarmi?

$$a = \{7, 3, 4\}, \quad b = \{-1, 2, -1\}, \quad c = \{4, 2, 4\}.$$

$$(\vec{a}, \vec{b}, \vec{c}) = \begin{vmatrix} 7 & 3 & 4 \\ -1 & 2 & -1 \\ 4 & 2 & 4 \end{vmatrix} = -56 - 12 - 8 + 32 + 14 + 12 = -18 \neq 0$$

a, b va c vektorlar komplanar emas.

- | | | |
|-------------------------|------------------------|-------------------------|
| $a = \{2, 3, 1\},$ | $a = \{3, 2, 1\},$ | $a = \{1, 5, 2\},$ |
| 1. $b = \{-1, 0, -1\},$ | 2. $b = \{2, 3, 4\},$ | 3. $b = \{-1, 1, -1\},$ |
| $c = \{2, 2, 2\}.$ | $c = \{3, 1, -1\}$ | $c = \{1, 1, 1\}.$ |
| $a = \{1, -1, -3\},$ | $a = \{3, 3, 1\},$ | $a = \{3, 1, -1\},$ |
| 4. $b = \{3, 2, 1\},$ | 5. $b = \{1, -2, 1\},$ | 6. $b = \{-2, -1, 0\},$ |
| $c = \{2, 3, 4\}.$ | $c = \{1, 1, 1\}.$ | $c = \{5, 2, -1\}.$ |
| $a = \{4, 3, 1\},$ | $a = \{4, 3, 1\},$ | $a = \{3, 2, 1\},$ |
| 7. $b = \{1, -2, 1\},$ | 8. $b = \{6, 7, 4\},$ | 9. $b = \{1, -3, -7\},$ |
| $c = \{2, 2, 2\}.$ | $c = \{2, 0, -1\}.$ | $c = \{1, 2, 3\}.$ |

$$\begin{array}{lll}
 a = \{3, 7, 2\}, & a = \{1, -2, 6\}, & a = \{6, 3, 4\}, \\
 10. \ b = \{-2, 0, -1\}, & 11. \ b = \{1, 0, 1\}, & 12. \ b = \{-1, 2, -1\}, \\
 c = \{2, 2, 1\}. & c = \{2, -6, 17\}. & c = \{2, 1, 2\}.
 \end{array}$$

$$\begin{array}{lll}
 a = \{2, 3, 2\}, & a = \{5, 3, 4\}, & a = \{3, 10, 5\}, \\
 13. \ b = \{4, 7, 5\}, & 14. \ b = \{-1, 0, -1\}, & 15. \ b = \{-2, -2, -3\}, \\
 c = \{2, 0, -1\}. & c = \{4, 2, 4\}. & c = \{2, 4, 3\}.
 \end{array}$$

$$\begin{array}{lll}
 a = \{-2, -4, -3\}, & a = \{3, 1, -1\}, & a = \{4, 2, 2\}, \\
 16. \ b = \{4, 3, 1\}, & 17. \ b = \{1, 0, -1\}, & 18. \ b = \{-3, -3, -3\}, \\
 c = \{6, 7, 4\}. & c = \{8, 3, -2\}. & c = \{2, 1, 2\}.
 \end{array}$$

$$\begin{array}{lll}
 a = \{4, 1, 2\}, & a = \{5, 3, 4\}, & a = \{3, 4, 2\}, \\
 19. \ b = \{9, 2, 5\}, & 20. \ b = \{4, 3, 3\}, & 21. \ b = \{1, 1, 0\}, \\
 c = \{1, 1, -1\}. & c = \{9, 5, 8\}. & c = \{8, 11, 6\}.
 \end{array}$$

$$\begin{array}{lll}
 a = \{4, -1, -6\}, & a = \{3, 1, 0\}, & a = \{3, 0, 3\}, \\
 22. \ b = \{1, -3, -7\}, & 23. \ b = \{-5, -4, -5\}, & 24. \ b = \{8, 1, 6\}, \\
 c = \{2, -1, -4\}. & c = \{4, 2, 4\}. & c = \{1, 1, -1\}.
 \end{array}$$

$$\begin{array}{lll}
 a = \{1, -1, 4\}, & a = \{6, 3, 4\}, & a = \{4, 1, 1\}, \\
 25. \ b = \{1, 0, 3\}, & 26. \ b = \{-1, -2, -1\}, & 27. \ b = \{-9, -4, -9\}, \\
 c = \{1, -3, 8\}. & c = \{2, 1, 2\}. & c = \{6, 2, 6\}.
 \end{array}$$

$$\begin{array}{lll}
 a = \{-3, 3, 3\}, & a = \{-7, 10, -5\}, & a = \{7, 4, 6\}, \\
 28. \ b = \{-4, 7, 6\}, & 29. \ b = \{0, -2, -1\}, & 30. \ b = \{2, 1, 1\}, \\
 c = \{3, 0, -1\}. & c = \{-2, 4, -1\}. & c = \{19, 11, 17\}.
 \end{array}$$

Javoblar. 5.1 yo'q; 5.2 ha; 5.3 yo'q; 5.4 ha; 5.5 yo'q; 5.6 ha; 5.7 yo'q; 5.8 ha; 5.9 ha;
 5.10 yo'q; 5.11 ha; 5.12 yo'q; 5.13 ha; 5.14 yo'q; 5.15 yo'q; 5.16 ha; 5.17 ha; 5.18 yo'q;
 5.19 ha; 5.20 yo'q; 5.21 ha; 5.22 ha; 5.23 yo'q; 5.24 ha; 5.25 yo'q; 5.26 yo'q; 5.27 yo'q;
 5.28 ha; 5.29 yo'q; 5.30 ha .

Tetraedrning balandligi va hajmi

Uchlari $A_1(x_1, y_1, z_1)$, $A_2(x_2, y_2, z_2)$, $A_3(x_3, y_3, z_3)$, $A_4(x_4, y_4, z_4)$ bo'lgan, hamda A_4 uchidan $A_1A_2A_3$ yog'iga balandlik tushirilgan tetraedrning hajmini topish masalasini qaraylik.

A_1 uchdan quyidagi vektorlarni o'tkazamiz:

$$\overrightarrow{A_1A_2} = \{x_2 - x_1, y_2 - y_1, z_2 - z_1\},$$

$$\overrightarrow{A_1A_3} = \{x_3 - x_1, y_3 - y_1, z_3 - z_1\},$$

$$\overrightarrow{A_1A_4} = \{x_4 - x_1, y_4 - y_1, z_4 - z_1\}.$$

Vektorlarning aralash ko'patmasining geometrik ma'nosidan quyidagiga egamiz:

$$V_t = \frac{1}{6} \times V_{pp} = \frac{1}{6} \left| \left(\overrightarrow{A_1A_2}, \overrightarrow{A_1A_3}, \overrightarrow{A_1A_4} \right) \right|,$$

bu yerda V_t va V_{pp} —lar mos ravishda $\overrightarrow{A_1A_2}$, $\overrightarrow{A_1A_3}$, $\overrightarrow{A_1A_4}$ vektorlar yordamida qurilgan tetraedr va parallelepipedning hajmlari.

Ikkinchi tomondan

$$V_t = \frac{1}{3} S_{DA_1A_2A_3} \times h,$$

vektor ko'paytmaning geometrik ma'nosidan esa,

$$S_{DA_1A_2A_3} = \frac{1}{2} \left| \left[\overrightarrow{A_1A_2}, \overrightarrow{A_1A_3} \right] \right|.$$

Demak, tetraedrning balandligi

$$h = \frac{3V_t}{S_{DA_1A_2A_3}} = \frac{\left| \left(\overrightarrow{A_1A_2}, \overrightarrow{A_1A_3}, \overrightarrow{A_1A_4} \right) \right|}{\left| \left[\overrightarrow{A_1A_2}, \overrightarrow{A_1A_3} \right] \right|}$$

ga teng bo'ladi.

6–masala. Uchlari A_1, A_2, A_3, A_4 nuqtalarda yordamida berilgan tetraedrning, A_4 uchidan $A_1A_2A_3$ yog'iga balandlik tushirilgan tetraedrning hajmini toping.

$$A_1(0, -1, -1),$$

$$A_2(-2, 3, 5),$$

$$A_3(1, -5, -9),$$

$$A_4(-1, -6, 3).$$

$$\overline{A_1A_2} = \{-2, 4, 6\},$$

$$\overline{A_1A_3} = \{1, -4, -8\},$$

$$\overline{A_1A_4} = \{-1, -5, 4\}.$$

$$V = \frac{1}{6} |(\overline{A_1A_2}, \overline{A_1A_3}, \overline{A_1A_4})| = \frac{1}{6} \times \begin{vmatrix} -2 & 4 & 6 \\ 1 & -4 & -8 \\ -1 & -5 & 4 \end{vmatrix} =$$

$$= \frac{1}{6} \times |32 - 30 + 32 - 24 + 80 - 16| = \frac{74}{6}.$$

$$V_{A_1A_2A_3A_4} = \frac{1}{3} S_{A_1A_2A_3} \times h \Rightarrow h = \frac{3V}{S}.$$

$$S_{A_1A_2A_3} = \frac{1}{2} |\overline{A_1A_2} \times \overline{A_1A_3}| = \frac{1}{2} \begin{vmatrix} i & j & k \\ -2 & 4 & 6 \\ 1 & -4 & -8 \end{vmatrix} =$$

$$= \frac{1}{2} |-8i - 10j + 4k| = \frac{1}{2} \sqrt{64 + 100 + 16} =$$

$$= \frac{1}{2} \sqrt{180} = \sqrt{45}.$$

$$h = \frac{3 \times 74}{6 \times \sqrt{45}} = \frac{37}{\sqrt{45}}.$$

$$\begin{aligned}
&A_1(1, 3, 6), \\
&A_2(2, 2, 1), \\
1. \quad &A_3(-1, 0, 1), \\
&A_4(-4, 6, -3).
\end{aligned}$$

$$\begin{aligned}
&A_1(-4, 2, 6), \\
&A_2(2, -3, 0), \\
2. \quad &A_3(-10, 5, 8), \\
&A_4(-5, 2, -4).
\end{aligned}$$

$$\begin{aligned}
&A_1(7, 2, 4), \\
&A_2(7, -1, -2), \\
3. \quad &A_3(3, 3, 1), \\
&A_4(-4, 2, 1).
\end{aligned}$$

$$\begin{aligned}
&A_1(2, 1, 4), \\
&A_2(-1, 5, -2), \\
4. \quad &A_3(-7, -3, 2), \\
&A_4(-6, -3, 6).
\end{aligned}$$

$$\begin{aligned}
&A_1(-1, -5, 2), \\
&A_2(-6, 0, -3), \\
5. \quad &A_3(3, 6, -3), \\
&A_4(-10, 6, 7).
\end{aligned}$$

$$\begin{aligned}
&A_1(5, 2, 0), \\
&A_2(2, 5, 0), \\
6. \quad &A_3(1, 2, 4), \\
&A_4(-1, 1, 1).
\end{aligned}$$

$$\begin{aligned}
&A_1(2, -1, -2), \\
&A_2(1, 2, 1), \\
7. \quad &A_3(5, 0, -6), \\
&A_4(-10, 9, -7).
\end{aligned}$$

$$\begin{aligned}
&A_1(-2, 0, -4), \\
&A_2(-1, 7, 1), \\
8. \quad &A_3(4, -8, -4), \\
&A_4(1, -4, 6).
\end{aligned}$$

$$\begin{aligned}
&A_1(14, 4, 5), \\
&A_2(-5, -3, 2), \\
9. \quad &A_3(-2, -6, -3), \\
&A_4(-2, 2, -1).
\end{aligned}$$

$$\begin{aligned}
&A_1(1, 2, 0), \\
&A_2(3, 0, -3), \\
10. \quad &A_3(5, 2, 6), \\
&A_4(8, 4, -9).
\end{aligned}$$

$$\begin{aligned}
&A_1(2, -1, 2), \\
&A_2(1, 2, -1), \\
11. \quad &A_3(3, 2, 1), \\
&A_4(-4, 2, 5).
\end{aligned}$$

$$\begin{aligned}
&A_1(1, 1, 2), \\
&A_2(-1, 1, 3), \\
12. \quad &A_3(2, -2, 4), \\
&A_4(-1, 0, -2).
\end{aligned}$$

$$\begin{aligned}
&A_1(2, 3, 1), \\
&A_2(4, 1, -2), \\
13. \quad &A_3(6, 3, 7), \\
&A_4(7, 5, -3).
\end{aligned}$$

$$\begin{aligned}
&A_1(1, 1, -1), \\
&A_2(2, 3, 1), \\
14. \quad &A_3(3, 2, 1), \\
&A_4(5, 9, -8).
\end{aligned}$$

$$\begin{aligned}
&A_1(1, 5, -7), \\
&A_2(-3, 6, 3), \\
15. \quad &A_3(-2, 7, 3), \\
&A_4(-4, 8, -12).
\end{aligned}$$

$$\begin{aligned}
&A_1(-3, 4, -7), \\
&A_2(1, 5, -4), \\
16. \quad &A_3(-5, -2, 0), \\
&A_4(2, 5, 4).
\end{aligned}$$

$$\begin{aligned}
&A_1(-1, 2, -3), \\
&A_2(4, -1, 0), \\
17. \quad &A_3(2, 1, -2), \\
&A_4(3, 4, 5).
\end{aligned}$$

$$\begin{aligned}
&A_1(4, -1, 3), \\
&A_2(-2, 1, 0), \\
18. \quad &A_3(0, -5, 1), \\
&A_4(3, 2, -6).
\end{aligned}$$

$$\begin{array}{lll}
A_1(1, -1, 1), & A_1(1, 2, 0), & A_1(1, 0, 2), \\
A_2(-2, 0, 3), & A_2(1, -1, 2), & A_2(1, 2, -1), \\
19. \quad A_3(2, 1, -1), & 20. \quad A_3(0, 1, -1), & 21. \quad A_3(2, -2, 1), \\
A_4(2, -2, -4). & A_4(-3, 0, 1). & A_4(2, 1, 0).
\end{array}$$

$$\begin{array}{lll}
A_1(1, 2, -3), & A_1(3, 10, -1), & A_1(-1, 2, 4), \\
A_2(1, 0, 1), & A_2(-2, 3, -5), & A_2(-1, -2, -4), \\
22. \quad A_3(-2, -1, 6), & 23. \quad A_3(-6, 0, -3), & 24. \quad A_3(3, 0, -1), \\
A_4(0, -5, -4). & A_4(1, -1, 2). & A_4(7, -3, 1).
\end{array}$$

$$\begin{array}{lll}
A_1(0, -3, 1), & A_1(1, 3, 0), & A_1(-2, -1, -1), \\
A_2(-4, 1, 2), & A_2(4, -1, 2), & A_2(0, 3, 2), \\
25. \quad A_3(2, -1, 5), & 26. \quad A_3(3, 0, 1), & 27. \quad A_3(3, 1, -4), \\
A_4(3, 1, -4). & A_4(-4, 3, 5). & A_4(-4, 7, 3).
\end{array}$$

$$\begin{array}{lll}
A_1(-3, -5, 6), & A_1(2, -4, -3), & A_1(1, -1, 2), \\
A_2(2, 1, -4), & A_2(5, -6, 0), & A_2(2, 1, 2), \\
28. \quad A_3(0, -3, -1), & 29. \quad A_3(-1, 3, -3), & 30. \quad A_3(1, 1, 4), \\
A_4(-5, 2, -8). & A_4(-10, -8, 7). & A_4(6, -3, 8).
\end{array}$$

Javoblar. 6.1 $23\frac{1}{3}; 2\sqrt{14}; 6.2 18\frac{2}{3}; 4; 6.3 21,5; \frac{43}{\sqrt{105}}; 6.4 \frac{80}{3}; \frac{10}{\sqrt{22}}; 6.5 190; 2\sqrt{38}; 6.6$

$12; \frac{18\sqrt{27}}{27}; 6.7 46\frac{2}{3}; 4\sqrt{14}; 6.8 83\frac{1}{3}; 5\sqrt{2}; 6.9 112\frac{2}{3}; \sqrt{26}; 6.10 34; 7\frac{2}{7};$

$6.11 11; 3\times\sqrt{\frac{11}{2}}; 6.12 5\frac{5}{6}; \sqrt{\frac{35}{2}}; 6.13 23\frac{1}{3}; 5; 6.14 7\frac{1}{2}; \frac{45\sqrt{17}}{17}; 6.15 17,5; 7;$

$6.16 25\frac{1}{6}; \sqrt{\frac{151}{15}}; 6.17 6\frac{2}{3}; 5\sqrt{2}; 6.18 45\frac{1}{3}; \frac{17\sqrt{5}}{5}; 6.19 5,5; \frac{33}{\sqrt{101}}; 6.20 \frac{19}{6}; \sqrt{\frac{19}{2}};$

$6.21 1\frac{1}{6}; \sqrt{\frac{7}{11}}; 6.22 16; 8\times\sqrt{\frac{2}{3}}; 6.23 45,5; 7; 6.24 24; 4; 6.25 32\frac{1}{3}; \sqrt{\frac{97}{2}};$

$6.26 2,5; 5\times\sqrt{\frac{3}{2}}; 6.27 23\frac{1}{3}; \frac{140}{\sqrt{1021}}; 6.28 \frac{191}{6}; \sqrt{\frac{191}{3}}; 6.29 73; \frac{438}{\sqrt{747}} 6.30 6; 3\sqrt{6}.$

Nuqtadan tekislikkacha bo'lgan masofa

Izlanayotgan masofani uchlari $M_0(x_0, y_0, z_0)$, $M_1(x_1, y_1, z_1)$, $M_2(x_2, y_2, z_2)$, $M_3(x_3, y_3, z_3)$ berilgan tetraedrning uchi $M_0(x_0, y_0, z_0)$ dan $M_1M_2M_3$ yog'iga tushirilgan balandlik orqali topish mumkin, ya'ni

$$d = \frac{|Ax_0 + By_0 + Cz_0 + D|}{\sqrt{A^2 + B^2 + C^2}}$$

Masalani hal qilishning boshqacha ko'rinishi esa, $M_0(x_0, y_0, z_0)$ nuqtadan tekislikkacha bo'lgan d masofa:

$$d = \frac{(\vec{n}, \overrightarrow{M_1M_0})}{|\vec{n}|},$$

bu yerda $|\vec{n}|$ –tekislikning normal vektori

$$|\vec{n}| = [\overrightarrow{M_1M_2}, \overrightarrow{M_1M_3}].$$

$$\overrightarrow{M_1M_2} = \{x_2 - x_1, y_2 - y_1, z_2 - z_1\}, \overrightarrow{M_1M_3} = \{x_3 - x_1, y_3 - y_1, z_3 - z_1\},$$

$$\overrightarrow{M_1M_0} = \{x_0 - x_1, y_0 - y_1, z_0 - z_1\} \text{ vektorlarning koordinatalarini topamiz}$$

va

$$|\vec{n}| = [\overrightarrow{M_1M_2}, \overrightarrow{M_1M_3}] = \begin{vmatrix} i & j & k \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix}$$

ni topamiz.

7–masala. M_1, M_2, M_3 nuqtalardan o'tuvchi M_0 nuqtadan

tekislikkacha bo'lgan masofani toping.

$$\begin{aligned} M_1(2, 3, 1), \\ M_2(4, 1, -2), \\ M_3(6, 3, 7), \\ M_0(-5, -4, 8). \end{aligned}$$

Uch nuqtadan o'tuvchi tekislik tenglamasi

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix} = 0,$$

$$\begin{vmatrix} x - 2 & y - 3 & z - 1 \\ 2 & -2 & -3 \\ 4 & 0 & 6 \end{vmatrix} = 0,$$

$$-12(x - 2) - 24(y - 3) + 8(z - 1) = 0,$$

$$-12x - 24y + 8z + 88 = 0,$$

$$d = \frac{|Ax_0 + By_0 + Cz_0 + D|}{\sqrt{A^2 + B^2 + C^2}},$$

$$d = \frac{|-12 \cdot (-5) - 24 \cdot (-4) + 8 \cdot 8 + 88|}{\sqrt{(-12)^2 + (-24)^2 + 8^2}} = \frac{308}{\sqrt{784}} = \frac{308}{28} = 11.$$

$M_1(-3, 4, -7),$	$M_1(-1, 2, -3),$	$M_1(-3, -1, 1),$
$M_2(1, 5, -4),$	$M_2(4, -1, 0),$	$M_2(-9, 1, -2),$
^{1.} $M_3(-5, -2, 0),$	^{2.} $M_3(2, 1, -2),$	^{3.} $M_3(3, -5, 4),$
$M_0(-12, 7, -1).$	$M_0(1, -6, -5).$	$M_0(-7, 0, -1).$

$M_1(1, -1, 1),$	$M_1(1, 2, 0),$	$M_1(1, 0, 2),$
$M_2(-2, 0, 3),$	$M_2(1, -1, 2),$	$M_2(1, 2, -1),$
^{4.} $M_3(2, 1, -1),$	^{5.} $M_3(0, 1, -1),$	^{6.} $M_3(2, -2, 1),$
$M_0(-2, 4, 2).$	$M_0(2, -1, 4).$	$M_0(-5, -9, 1).$

$M_1(1, 2, -3),$	$M_1(3, 10, -1),$	$M_1(-1, 2, 4),$
$M_2(1, 0, 1),$	$M_2(-2, 3, -5),$	$M_2(-1, -2, -4),$
^{7.} $M_3(-2, -1, 6),$	^{8.} $M_3(-6, 0, -3),$	^{9.} $M_3(3, 0, -1),$
$M_0(3, -2, -9).$	$M_0(-6, 7, -10).$	$M_0(-2, 3, 5).$
$M_1(0, -3, 1),$	$M_1(1, 3, 0),$	$M_1(-2, -1, -1),$
$M_2(-4, 1, 2),$	$M_2(4, -1, 2),$	$M_2(0, 3, 2),$
^{10.} $M_3(2, -1, 5),$	^{11.} $M_3(3, 0, 1),$	^{12.} $M_3(3, 1, -4),$
$M_0(-3, 4, -5).$	$M_0(4, 3, 0).$	$M_0(-21, 20, -16).$
$M_1(-3, -5, 6),$	$M_1(2, -4, -3),$	$M_1(1, -1, 2),$
$M_2(2, 1, -4),$	$M_2(5, -6, 0),$	$M_2(2, 1, 2),$
^{13.} $M_3(0, -3, -1),$	^{14.} $M_3(-1, 3, -3),$	^{15.} $M_3(1, 1, 4),$
$M_0(3, 6, 68).$	$M_0(2, -10, 8).$	$M_0(-3, 2, 7).$
$M_1(1, 3, 6),$	$M_1(-4, 2, 6),$	$M_1(7, 2, 4),$
$M_2(2, 2, 1),$	$M_2(2, -3, 0),$	$M_2(7, -1, -2),$
^{16.} $M_3(-1, 0, 1),$	^{17.} $M_3(-10, 5, 8),$	^{18.} $M_3(-5, -2, -1),$
$M_0(5, -4, 5).$	$M_0(-12, 1, 8).$	$M_0(10, 1, 8).$
$M_1(2, 1, 4),$	$M_1(-1, -5, 2),$	$M_1(0, -1, -1),$
$M_2(3, 5, -2),$	$M_2(-6, 0, -3),$	$M_2(-2, 3, 5),$
^{19.} $M_3(-7, -3, 2),$	^{20.} $M_3(3, 6, -3),$	^{21.} $M_3(1, -5, -9),$
$M_0(-3, 1, 8).$	$M_0(10, -8, -7).$	$M_0(-4, -13, 6).$
$M_1(5, 2, 0),$	$M_1(2, -1, -2),$	$M_1(-2, 0, -4),$
$M_2(2, 5, 0),$	$M_2(1, 2, 1),$	$M_2(-1, 7, 1),$
^{22.} $M_3(1, 2, 4),$	^{23.} $M_3(5, 0, -6),$	^{24.} $M_3(4, -8, -4),$
$M_0(-3, -6, -8).$	$M_0(14, -3, 7).$	$M_0(-6, 5, 5).$

$$\begin{array}{lll}
M_1(14, 4, 5), & M_1(1, 2, 0), & M_1(2, -1, 2), \\
M_2(-5, -3, 2), & M_2(3, 0, -3), & M_2(1, 2, -1), \\
^{25.} M_3(-2, -6, -3), & ^{26.} M_3(5, 2, 6) & ^{27.} M_3(3, 2, 1), \\
M_0(-1, -8, 7). & M_0(-13, -8, 16). & M_0(-5, 3, 7).
\end{array}$$

$$\begin{array}{lll}
M_1(1, 1, 2), & M_1(1, 1, -1), & M_1(1, 5, -7), \\
M_2(-1, 1, 3), & M_2(2, 3, 1), & M_2(-3, 6, 3), \\
^{28.} M_3(2, -2, 4), & ^{29.} M_3(3, 2, 1), & ^{30.} M_3(-2, 7, 3), \\
M_0(2, 3, 8). & M_0(-3, -7, 6). & M_0(1, -1, 2).
\end{array}$$

Javoblar. 7.1 $\frac{459}{\sqrt{2265}}$; 7.2 $5\sqrt{2}$; 7.3 0; 7.4 $\frac{9}{\sqrt{101}}$; 7.5 $\frac{1}{\sqrt{38}}$; 7.6 $\sqrt{77}$; 7.7 $2\sqrt{6}$; 7.8 7;

7.9 $\frac{5}{9}$; 7.10 $\frac{90}{\sqrt{194}}$; 7.11 $\sqrt{6}$; 7.12 $\frac{1023}{\sqrt{1021}}$; 7.13 $\sqrt{573}$; 7.14 $\frac{73}{\sqrt{83}}$; 7.15 $\sqrt{6}$; 7.16 $2\sqrt{14}$; 7.17

4; 7.18 3; 7.19 4; 7.20 $2\sqrt{38}$; 7.21 $2\sqrt{45}$; 7.22 $8\sqrt{3}$; 7.23 $4\sqrt{14}$; 7.24 $\frac{23\sqrt{2}}{5}$;

7.25 $3\sqrt{\frac{13}{2}}$; 7.26 $19\frac{1}{7}$; 7.27 $2\sqrt{22}$; 7.28 $7\sqrt{\frac{7}{10}}$; 7.29 $\frac{45}{\sqrt{17}}$; 7.30 7.

Normal vektori berilgan tekislik tenglamasi

8–masala. \overline{BC} vektorga perpendikulyar bo'lgan A nuqtadan o'tuvchi tekislik tenglamasini yozing.

$$\begin{array}{l}
A(0, -2, 8), \\
B(4, 3, 2), \\
C(1, 4, 3). \\
\overline{BC} = \{-3, 1, 1\}.
\end{array}$$

\overline{BC} izlanayotgan tekislikka perpendikulyar bo'lganligidan, uni vektorning normal sifatida olish mumkin, u holda

$$- 3(x - 0) + (y + 2) + (z - 8) = 0,$$

$$- 3x + y + z - 6 = 0.$$

$$\begin{aligned} & A(1, 0, -2), \\ 1. & B(2, -1, 3), \\ & C(0, -3, 2). \end{aligned}$$

$$\begin{aligned} & A(-1, 3, 4), \\ 2. & B(-1, 5, 0), \\ & C(2, 6, 1). \end{aligned}$$

$$\begin{aligned} & A(4, -2, 0), \\ 3. & B(1, -1, -5), \\ & C(-2, 1, -3). \end{aligned}$$

$$\begin{aligned} & A(-8, 0, 7), \\ 4. & B(-3, 2, 4), \\ & C(-1, 4, 5). \end{aligned}$$

$$\begin{aligned} & A(7, -5, 1), \\ 5. & B(5, -1, -3), \\ & C(3, 0, -4). \end{aligned}$$

$$\begin{aligned} & A(-3, 5, -2), \\ 6. & B(-4, 0, 3), \\ & C(-3, 2, 5). \end{aligned}$$

$$\begin{aligned} & A(1, -1, 8), \\ 7. & B(-4, -3, 10), \\ & C(-1, -1, 7). \end{aligned}$$

$$\begin{aligned} & A(-2, 0, -5), \\ 8. & B(2, 7, -3), \\ & C(1, 10, -1). \end{aligned}$$

$$\begin{aligned} & A(1, 9, -4), \\ 9. & B(5, 7, 1), \\ & C(3, 5, 0). \end{aligned}$$

$$\begin{aligned} & A(-7, 0, 3), \\ 10. & B(1, -5, -4), \\ & C(2, -3, 0). \end{aligned}$$

$$\begin{aligned} & A(0, -3, 5), \\ 11. & B(-7, 2, 6), \\ & C(-3, 2, 4). \end{aligned}$$

$$\begin{aligned} & A(5, -1, 2), \\ 12. & B(2, -4, 3), \\ & C(4, -1, 3). \end{aligned}$$

$$\begin{aligned} & A(-3, 7, 2), \\ 13. & B(3, 5, 1), \\ & C(4, 5, 3). \end{aligned}$$

$$\begin{aligned} & A(1, -1, 5), \\ 14. & B(0, 7, 8), \\ & C(-1, 3, 8). \end{aligned}$$

$$\begin{aligned} & A(-10, 0, 9), \\ 15. & B(12, 4, 11), \\ & C(8, 5, 15). \end{aligned}$$

$$\begin{aligned} & A(3, -3, -6), \\ 16. & B(1, 9, -5), \\ & C(6, 6, -4). \end{aligned}$$

$$\begin{aligned} & A(2, 1, 7), \\ 17. & B(9, 0, 2), \\ & C(9, 2, 3). \end{aligned}$$

$$\begin{aligned} & A(-7, 1, -4), \\ 18. & B(8, 11, -3), \\ & C(9, 9, -1). \end{aligned}$$

$$\begin{aligned} & A(1, 0, -6), \\ 19. & B(-7, 2, 1), \\ & C(-9, 6, 1). \end{aligned}$$

$$\begin{aligned} & A(-3, 1, 0), \\ 20. & B(6, 3, 3), \\ & C(9, 4, -2). \end{aligned}$$

$$\begin{aligned} & A(-4, -2, 5), \\ 21. & B(3, -3, -7), \\ & C(9, 3, -7). \end{aligned}$$

$$\begin{array}{lll} A(0, -8, 10), & A(1, -5, -2), & A(0, 7, -9), \\ 22. B(-5, 5, 7), & 23. B(6, -2, 1), & 24. B(-1, 8, -11), \\ C(-8, 0, 4). & C(2, -2, -2). & C(-4, 3, -12). \end{array}$$

$$\begin{array}{lll} A(-3, -1, 7), & A(5, 3, 1), & A(-1, 2, -2), \\ 25. B(0, 2, -6), & 26. B(0, 0, -3), & 27. B(13, 14, 1), \\ C(2, 3, -5). & C(5, -1, 0). & C(14, 15, 2). \end{array}$$

$$\begin{array}{lll} A(7, -5, 0), & A(-3, 6, 4), & A(2, 5, -3), \\ 28. B(8, 3, -1), & 29. B(8, -3, 5), & 30. B(7, 8, -1), \\ C(8, 5, 1). & C(0, -3, 7). & C(9, 7, 4). \end{array}$$

Javoblar. 8.1 $2x + 2y + z = 0$; 8.2 $3x + y + z - 4 = 0$; 8.3 $-3x + 2y + 2z + 16 = 0$;
 8.4 $-4x + 2y + z - 39 = 0$; 8.5 $-2x + y - z + 20 = 0$; 8.6 $x + 2y + 2z - 9 = 0$;
 8.7 $3x + 2y - 3z + 23 = 0$; 8.8 $-x + 3y + 2z + 8 = 0$; 8.9 $-2x - 2y - z - 16 = 0$;
 8.10 $x + 2y + 4z - 5 = 0$; 8.11 $2x - z + 5 = 0$; 8.12 $2x + 3y - 7 = 0$; 8.13 $x + 2z - 1 = 0$;
 8.14 $x + 4y + 3 = 0$; 8.15 $-4x + y + 4z - 76 = 0$; 8.16 $5x - 3y + z - 18 = 0$;
 8.17 $2y + z - 9 = 0$; 8.18 $x - 2y + 2z + 17 = 0$; 8.19 $-x + 2y + 1 = 0$;
 8.20 $3x + y - 5z + 8 = 0$; 8.21 $x + y + 6 = 0$; 8.22 $3x + 5y + 3z + 10 = 0$;
 8.23 $4x + 3z + 2 = 0$; 8.24 $-3x - 5y - z + 26 = 0$; 8.25 $2x + y + z = 0$;
 8.26 $5x - y + 3z - 19 = 0$; 8.27 $x + y + z + 1 = 0$; 8.28 $y + z + 5 = 0$; 8.29 $x + z - 1 = 0$; 8.30
 $2x - y + 5z + 16 = 0$.

Tekisliklar orasidagi burchak

Fazoda tekisliklar

$$A_1x + B_1y + C_1z + D_1 = 0 \quad \text{va} \quad A_2x + B_2y + C_2z + D_2 = 0$$

tenglamalar bilan berilgan bo'lsin. Ular orasidagi j burchak ushbu formula bilan hisoblanadi:

$$\cos j = \frac{\vec{n}_1 \times \vec{n}_2}{|\vec{n}_1| |\vec{n}_2|},$$

bunda $\vec{n}_1 = \{A_1, B_1, C_1\}$, $\vec{n}_2 = \{A_2, B_2, C_2\}$ – mos ravishda berilgan tekisliklarning normal vektorlari.

9–masala. Tekisliklar orasidagi burchakni toping.

$$6x + 2y - 4z + 17 = 0,$$

$$9x + 3y - 6z - 4 = 0.$$

$$\vec{n}_1 = \{6, 2, -4\},$$

$$\vec{n}_2 = \{9, 3, -6\}.$$

$$\begin{aligned} \cos j &= \frac{6 \times 9 + 2 \times 3 + (-4) \times (-6)}{\sqrt{6^2 + 2^2 + (-4)^2} \times \sqrt{9^2 + 3^2 + (-6)^2}} = \\ &= \frac{84}{\sqrt{56} \times \sqrt{126}} = \frac{84}{\sqrt{7056}} = \frac{84}{84} = 1, \\ j &= \arccos 1 = 0. \end{aligned}$$

$$\begin{array}{lll} 1. \quad \begin{array}{l} x - 3y + 5 = 0, \\ 2x - y + 5z - 16 = 0. \end{array} & 2. \quad \begin{array}{l} x - 3y + z - 1 = 0, \\ x + z - 1 = 0. \end{array} & 3. \quad \begin{array}{l} 4x - 5y + 3z - 1 = 0, \\ x - 4y - z + 9 = 0. \end{array} \end{array}$$

$$\begin{array}{lll} 4. \quad \begin{array}{l} 3x - y + 2z + 15 = 0, \\ 5x + 9y - 3z - 1 = 0. \end{array} & 5. \quad \begin{array}{l} x - y\sqrt{2} + z - 1 = 0, \\ x + y\sqrt{2} - z + 36 = 0. \end{array} & 6. \quad \begin{array}{l} 3y - z = 0, \\ 2y + z = 0. \end{array} \end{array}$$

$$\begin{array}{lll} 7. \quad \begin{array}{l} 6x + 3y - 2z = 0, \\ x + 2y + 6z - 12 = 0. \end{array} & 8. \quad \begin{array}{l} x + 2y + 2z - 3 = 0, \\ 16x + 12y - 15z - 1 = 0. \end{array} & 9. \quad \begin{array}{l} 2x - y + 5z + 16, \\ x + 2y + 3z + 8 = 0. \end{array} \end{array}$$

$$\begin{array}{lll} 10. \quad \begin{array}{l} 2x + 2y + z - 1 = 0, \\ x + z - 1 = 0. \end{array} & 11. \quad \begin{array}{l} 3x + y + z - 4 = 0, \\ y + z + 5 = 0. \end{array} & 12. \quad \begin{array}{l} 3x - 2y - 2z - 16 = 0, \\ x + y - 3z - 7 = 0. \end{array} \end{array}$$

$$\begin{array}{lll} 13. \quad \begin{array}{l} 2x + 2y + z + 9 = 0, \\ x - y + 3z - 1 = 0. \end{array} & 14. \quad \begin{array}{l} x + 2y + 2z - 3 = 0, \\ 2x - y + 2z + 5 = 0. \end{array} & 15. \quad \begin{array}{l} 3x + 2y - 3z - 1 = 0, \\ x + y + z - 7 = 0. \end{array} \end{array}$$

$$\begin{array}{lll} 16. \quad x - 3y - 2z - 8 = 0, & 17. \quad 3x - 2y + 3z + 23 = 0, & 18. \quad x + y + 3z - 7 = 0, \\ & & y + z - 1 = 0. \\ & & y + z + 5 = 0. \end{array}$$

$$\begin{array}{lll} 19. \quad x - 2y + 2z + 17 = 0, & 20. \quad x - 2y - 1 = 0, & 21. \quad 2x - z + 5 = 0, \\ & & 2x + 3y - 7 = 0. \\ & & x + y + 6 = 0. \end{array}$$

$$\begin{array}{lll} 22. \quad 5x + 3y + z - 18 = 0, & 23. \quad 4x + 3z - 2 = 0, & 24. \quad x + 4y - z + 1 = 0, \\ & & 2x + y + 4z - 3 = 0. \\ & & x + 2y + 2z + 5 = 0. \end{array}$$

$$\begin{array}{lll} 25. \quad 2y + z - 9 = 0, & 26. \quad 2x - 6y + 14z - 1 = 0, & 27. \quad x - y + 7z - 1 = 0, \\ & & 2x - 2y - 5 = 0. \\ & & 5x - 15y + 35z - 3 = 0. \\ & & x - y + 2z - 1 = 0. \end{array}$$

$$\begin{array}{lll} 28. \quad 3x - y - 5 = 0, & 29. \quad x + y + z\sqrt{2} - 3 = 0, & 30. \quad x + 2y - 2z - 7 = 0, \\ & & x + y - 35 = 0. \\ & & x - y + z\sqrt{2} - 1 = 0. \\ & & 2x + y - 3 = 0. \end{array}$$

Javoblar. 9.1 $\arccos \frac{\sqrt{3}}{6} \approx 73^\circ 13' 17''$; 9.2 $\arccos \sqrt{\frac{2}{11}} \approx 64^\circ 45' 38''$

9.3 $\arccos 0,7 \approx 45^\circ 34' 23''$; 9.4 90° ; 9.5 60° ; 9.6 45° ; 9.7 90° ; 9.8 $\arccos \frac{2}{15} \approx 82^\circ$;

9.9 $\arccos \sqrt{\frac{15}{28}} \approx 42^\circ 57' 7''$; 9.10 $\arccos \frac{\sqrt{2}}{2} = 45^\circ$; 9.11 $\arccos \sqrt{\frac{2}{11}} \approx 64^\circ 45' 38''$

9.12 $\arccos \frac{7}{\sqrt{187}} \approx 59^\circ 12' 37''$; 9.13 $\arccos \frac{1}{\sqrt{11}} \approx 72^\circ 27' 56''$; 9.14 $\arccos \frac{4}{9} \approx 63^\circ 36' 44''$

9.15 $\arccos \frac{2}{\sqrt{66}} \approx 75^\circ 44' 54''$; 9.16 $\arccos 0 = 90^\circ$; 9.17 $\arccos \frac{1}{\sqrt{44}} \approx 81^\circ 19' 45''$

9.18 $\arccos 2\sqrt{\frac{2}{11}} = 31^\circ 28' 56''$; 9.19 $\arccos \frac{\sqrt{5}}{3} = 41^\circ 48' 37''$

9.20 $\arccos \frac{3}{\sqrt{10}} = 18^\circ 26' 56''$; 9.21 $\arccos \frac{4}{\sqrt{65}} \approx 60^\circ 15' 18''$; 9.22 $\arccos \frac{\sqrt{7}}{5} = 58^\circ 37' 7''$

9.23 $\arccos \frac{2}{3} = 48^\circ 11' 23''$; 9.24 $\arccos \frac{\sqrt{2}}{3\sqrt{42}} = 84^\circ 54' 41''$; 9.25 90° ; 9.26 1 ;

9.27 $\arccos \sqrt{\frac{2}{102}} = 78^\circ 34' 42''$; 9.28 45° ; 9.29 60° 9.30 45° .

Bir xil uzoqlikda yotgan nuqtaning koordinatalari

10–masala. B va C nuqlardan bir xil uzoqlikda yotgan A nuqtaning koordinatalarini toping.

$$A(x, 0, 0), \quad B(1, 2, 3), \quad C(2, 6, 10).$$

$$AB = \sqrt{(1-x)^2 + 2^2 + 3^2} = \sqrt{x^2 - 2x + 14},$$

$$AC = \sqrt{(2-x)^2 + 6^2 + 10^2} = \sqrt{x^2 - 4x + 140}.$$

Shartga ko'ra $AB = AC$ ekanligidan

$$\sqrt{x^2 - 2x + 14} = \sqrt{x^2 - 4x + 140},$$

$$x^2 - 2x + 14 = x^2 - 4x + 140,$$

$$2x = 126,$$

$$x = 63$$

Demak, $A(63, 0, 0)$.

$A(0, 0, z),$	$A(0, 0, z),$	$A(0, 0, z),$
1. $B(5, 1, 0),$	2. $B(3, 3, 1),$	3. $B(3, 1, 3),$
$C(0, 2, 3).$	$C(4, 1, 2).$	$C(1, 4, 2).$
$A(0, 0, z),$	$A(0, 0, z),$	$A(0, 0, z),$
4. $B(-1, -1, -6),$	5. $B(-13, 4, 6),$	6. $B(-5, -5, 6),$
$C(2, 3, 5).$	$C(10, -9, 5).$	$C(-7, 6, 2).$
$A(0, 0, z),$	$A(0, 0, z),$	$A(0, 0, z),$
7. $B(-18, 1, 0),$	8. $B(10, 0, -2),$	9. $B(-6, 7, 5),$
$C(15, -10, 2).$	$C(9, -2, 1).$	$C(8, -4, 3).$

- | | | |
|---------------------|---------------------|----------------------|
| $A(0, 0, z),$ | $A(0, 0, z),$ | $A(0, y, 0),$ |
| 10. $B(6, -7, 1),$ | 11. $B(7, 0, -15),$ | 12. $B(3, 0, 3),$ |
| $C(-1, 2, 5).$ | $C(2, 10, -12).$ | $C(0, 2, 4).$ |
| $A(0, y, 0),$ | $A(0, y, 0),$ | $A(0, y, 0),$ |
| 13. $B(1, 6, 4),$ | 14. $B(-2, 8, 10),$ | 15. $B(-2, -4, 6),$ |
| $C(5, 7, 1).$ | $C(6, 11, -2).$ | $C(7, 2, 5).$ |
| $A(0, y, 0),$ | $A(0, y, 0),$ | $A(0, y, 0),$ |
| 16. $B(2, 2, 4),$ | 17. $B(0, -4, 1),$ | 18. $B(0, 5, -9),$ |
| $C(0, 4, 2).$ | $C(1, -3, 5).$ | $C(-1, 0, 5).$ |
| $A(0, y, 0),$ | $A(0, y, 0),$ | $A(0, y, 0),$ |
| 19. $B(-2, 4, -6),$ | 20. $B(7, 3, -4),$ | 21. $B(0, -2, 4),$ |
| $C(8, 5, 1).$ | $C(1, 5, 7).$ | $C(-4, 0, 4).$ |
| $A(x, 0, 0),$ | $A(x, 0, 0),$ | $A(x, 0, 0),$ |
| 22. $B(0, 1, 3),$ | 23. $B(4, 0, 5),$ | 24. $B(8, 1, -7),$ |
| $C(2, 0, 4).$ | $C(5, 4, 2).$ | $C(10, -2, 1).$ |
| $A(x, 0, 0),$ | $A(x, 0, 0),$ | $A(x, 0, 0),$ |
| 25. $B(3, 5, 6),$ | 26. $B(4, 5, -2),$ | 27. $B(-2, 0, 6),$ |
| $C(1, 2, 3).$ | $C(2, 3, 4).$ | $C(0, -2, -4).$ |
| $A(x, 0, 0),$ | $A(x, 0, 0),$ | $A(x, 0, 0),$ |
| 28. $B(1, 5, 9),$ | 29. $B(4, 6, 8),$ | 30. $B(-2, -4, -6),$ |
| $C(3, 7, 11).$ | $C(2, 4, 6).$ | $C(-1, -2, -3).$ |

Javoblar. 10.1 $A(0; 0; -2\frac{1}{6})$; 10.2 $A(0; 0; 1)$; 10.3 $A(0; 0; -1)$; 10.4 $A(0; 0; 0)$;

10.5 $A(0; 0; 7,5)$; 10.6 $A(0; 0; -\frac{3}{8})$; 10.7 $A(0; 0; 1)$; 10.8 $A(0; 0; -3)$; 10.9 $A(0; 0; 5,25)$;

- 10.10 $A(0; 0; -7)$; 10.11 $A(0; 0; -4\frac{1}{3})$; 10.12 $A(0; \frac{1}{2}; 0)$; 10.13 $A(0; 11; 0)$;
 10.14 $A(0; -\frac{7}{6}; 0)$; 10.15 $A(0; 1\frac{5}{6}; 0)$; 10.16 $A(0; -1; 0)$; 10.17 $A(0; 9; 0)$;
 10.18 $A(0; 8; 0)$; 10.19 $A(0; 17; 0)$; 10.20 $A(0; \frac{1}{4}; 0)$; 10.21 $A(0; 3; 0)$;
 10.22 $A(2,5; 0; 0)$; 10.23 $A(2; 0; 0)$; 10.24 $A(-2,25; 0; 0)$; 10.25 $A(14; 0; 0)$;
 10.26 $A(4; 0; 0)$; 10.27 $A(-5; 0; 0)$; 10.28 $A(18; 0; 0)$; 10.29 $A(15; 0; 0)$;
 10.30 $A(-21; 0; 0)$.

11–masala. Markazi koordinata boshida bo'lgan k –gomotetiya koeffitsiyenti bo'lsin. A nuqta a tekislikning obrazi(aksi)ga tegishliligini tekshiring.

$$\begin{aligned} A(1, 1, 1), \\ a : 7x - 6y + z - 5 = 0, \\ k = -2. \end{aligned}$$

Markazi koordinata boshida bo'lgan a tekislik o'xshash akslantirishlarga ko'ra a tekislikka o'tadi.

$$\begin{aligned} a : Ax + By + Cz + D = 0, \\ a : Ax + By + Cz + k \times D = 0, \\ a' : 7x - 6y + z + 10 = 0. \\ A(1, 1, 1) \rightarrow 7 - 6 + 1 + 10 = 0, \\ 12 \neq 0. \end{aligned}$$

Shunday qilib, A nuqta a tekislikning obrazi (aksi)ga tegishli bo'lmas ekan.

$$\begin{aligned} A(1, 2, -1), & \quad A(2, 1, 2), \\ 1. a : 2x + 3y + z - 1 = 0, & \quad 2. a : x - 2y + z + 1 = 0, \\ k = 2. & \quad k = -2. \end{aligned}$$

$$A(-1, 1, 1),$$

$$3. \mathbf{a} : 3x - y + 2z + 4 = 0,$$

$$k = \frac{1}{2}.$$

$$A_{\text{Cl}}^{\text{æ}}, \frac{1}{3}, -2\frac{\ddot{\text{o}}}{\text{ø}}$$

$$5. \mathbf{a} : x - 3y + z + 6 = 0,$$

$$k = \frac{1}{3}.$$

$$A(2, 0, -1),$$

$$7. \mathbf{a} : x - 3y + 5z - 1 = 0,$$

$$k = -1.$$

$$A(2, -5, 4),$$

$$9. \mathbf{a} : 5x + 2y - z + 3 = 0,$$

$$k = \frac{4}{3}.$$

$$A(-2, 3, -3),$$

$$11. \mathbf{a} : 3x + 2y - z - 2 = 0,$$

$$k = \frac{3}{2}.$$

$$A(0, 1, -1),$$

$$13. \mathbf{a} : 6x - 5y + 3z - 4 = 0,$$

$$k = -\frac{3}{4}.$$

$$A(-2, 4, 1),$$

$$4. \mathbf{a} : 3x + y + 2z + 2 = 0,$$

$$k = 3.$$

$$A_{\text{C}}^{\text{æ}}, \frac{1}{3}, 1\frac{\ddot{\text{o}}}{\text{ø}}$$

$$6. \mathbf{a} : 2x - 3y + 3z - 2 = 0,$$

$$k = 1,5.$$

$$A(1, -2, 1),$$

$$8. \mathbf{a} : 5x + y - z + 6 = 0,$$

$$k = 2/3.$$

$$A(2, -3, 1),$$

$$10. \mathbf{a} : x + y - 2z + 2 = 0,$$

$$k = \frac{5}{2}.$$

$$A_{\text{C}}^{\text{æ}}, \frac{1}{3}, 1\frac{\ddot{\text{o}}}{\text{ø}}$$

$$12. \mathbf{a} : 4x - 3y + 5z - 10 = 0,$$

$$k = \frac{1}{2}.$$

$$A(2, 3, -2),$$

$$14. \mathbf{a} : 3x - 2y + 4z - 6 = 0,$$

$$k = -\frac{4}{3}.$$

$$A(-2, -1, 1),$$

$$15. \mathbf{a} : x - 2y + 6z - 10 = 0,$$

$$k = \frac{3}{5}.$$

$$A\left(-\frac{1}{3}, 1, 1\right),$$

$$17. \mathbf{a} : 3x - y + 5z - 6 = 0,$$

$$k = \frac{5}{6}.$$

$$A(-1, 2, 3),$$

$$19. \mathbf{a} : x - 3y + z + 2 = 0,$$

$$k = 2,5.$$

$$A(3, 5, 2),$$

$$21. \mathbf{a} : 5x - 3y + z - 4 = 0,$$

$$k = \frac{1}{2}.$$

$$A(-1, 1, -2),$$

$$23. \mathbf{a} : 4x - y + 3z - 6 = 0,$$

$$k = -\frac{5}{3}.$$

$$A(-3, -2, 4),$$

$$25. \mathbf{a} : 2x - 3y + z - 5 = 0,$$

$$k = -\frac{4}{5}.$$

$$A(5, 0, -1),$$

$$16. \mathbf{a} : 2x - y + 3z - 1 = 0,$$

$$k = 3.$$

$$A(2, 5, 1),$$

$$18. \mathbf{a} : 5x - 2y + z - 3 = 0,$$

$$k = \frac{1}{3}.$$

$$A(4, 3, 1),$$

$$20. \mathbf{a} : 3x - 4y + 5z - 6 = 0,$$

$$k = \frac{5}{6}.$$

$$A(4, 0, -3),$$

$$22. \mathbf{a} : 7x - y + 3z - 1 = 0,$$

$$k = 3.$$

$$A(2, -5, -1),$$

$$24. \mathbf{a} : 5x + 2y - 3z - 9 = 0,$$

$$k = \frac{1}{3}.$$

$$A(5, 0, -6),$$

$$26. \mathbf{a} : 6x - y - z + 7 = 0,$$

$$k = \frac{2}{7}.$$

$$A(1, 2, 2),$$

$$27. a : 3x - z + 5 = 0,$$

$$k = -\frac{1}{5}.$$

$$A(3, 2, 4),$$

$$28. a : 2x - 3y + z - 6 = 0,$$

$$k = \frac{2}{3}.$$

$$A(7, 0, -1),$$

$$29. a : x - y - z - 1 = 0,$$

$$k = 4.$$

$$A(0, 3, -1),$$

$$30. a : 2x - y + 3z - 1 = 0,$$

$$k = 2.$$

Javoblar. 11.1 yo'q; 11.2 ha; 11.3 ha; 11.4 yo'q; 11.5 ha; 11.6 ha; 11.7 yo'q; 11.8 yo'q; 11.9 ha; 11.10 yo'q; 11.11 ha; 11.12 ha; 11.13 yo'q; 11.14 ha; 11.15 ha; 11.16 yo'q; 11.17 ha; 11.18 ha; 11.19 yo'q; 11.20 ha; 11.21 ha; 11.22 yo'q; 11.23 yo'q; 11.24 ha; 11.25 yo'q; 11.26 yo'q; 11.27 ha; 11.28 yo'q; 11.29 yo'q; 11.30 yo'q.

To'g'ri chiziqning kanonik tenglamasi

12–masala. To'g'ri chiziqning kanonik tenglamasini yozing.

$$x - 3y + 2z + 2 = 0,$$

$$x + 3y + z + 14 = 0.$$

$$\overline{S} = \overline{n_1}, \overline{n_2} = \begin{vmatrix} i & j & k \\ 1 & -3 & 2 \\ 1 & 3 & 1 \end{vmatrix} = -9i + j + 6k.$$

$$\overline{S} = \{-9, 1, 6\}.$$

(x_0, y_0, z_0) to'g'ri chiziqdan o'tuvchi biror nuqtaning koordinasini topamiz.

z ning koordinatasiga $z = 0$ qiymatni beramiz

$$\begin{cases} x - 3y + 2 = 0, \\ x + 3y + 14 = 0 \end{cases} \Rightarrow \begin{cases} x - 3y + 2 = 0, \\ 6y = -12 \end{cases} \Rightarrow \begin{cases} x = -8, \\ y = -2 \end{cases}$$

Shunday qilib, izlanayotgan nuqtaning koordinatasi $(-8, -2, 0)$.

To'g'ri chiziqli tenglamasi

$$\frac{x+8}{-9} = \frac{y+2}{1} = \frac{z}{6}.$$

$$\begin{array}{l} 2x + y + z - 2 = 0, \\ 1. \quad 2x - y - 3z + 6 = 0. \end{array}$$

$$\begin{array}{l} x - 2y + z - 4 = 0, \\ 2. \quad 2x + 2y - z - 8 = 0. \end{array}$$

$$\begin{array}{l} x + y + z - 2 = 0, \\ 3. \quad x - y - 2z + 2 = 0. \end{array}$$

$$\begin{array}{l} 2x + 3y + z + 6 = 0, \\ 4. \quad x - 3y - 2z + 3 = 0. \end{array}$$

$$\begin{array}{l} 3x + y - z - 6 = 0, \\ 5. \quad 3x - y + 2z = 0. \end{array}$$

$$\begin{array}{l} x + 5y + 2z + 11 = 0, \\ 6. \quad x - y - z - 1 = 0. \end{array}$$

$$\begin{array}{l} 3x + 4y - 2z + 1 = 0, \\ 7. \quad 2x - 4y + 3z + 4 = 0. \end{array}$$

$$\begin{array}{l} 5x + y - 3z + 4 = 0, \\ 8. \quad x - y + 2z + 2 = 0. \end{array}$$

$$\begin{array}{l} x - y - z - 2 = 0, \\ 9. \quad x - 2y + z + 4 = 0. \end{array}$$

$$\begin{array}{l} 4x + y - 3z + 2 = 0, \\ 10. \quad 2x - y + z - 8 = 0. \end{array}$$

$$\begin{array}{l} 3x + 3y - 2z - 1 = 0, \\ 11. \quad 2x - 3y + z + 6 = 0. \end{array}$$

$$\begin{array}{l} 6x - 7y - 4z - 2 = 0, \\ 12. \quad x + 7y - z - 5 = 0. \end{array}$$

$$\begin{array}{l} 8x - y - 3z - 1 = 0, \\ 13. \quad x + y + z + 10 = 0. \end{array}$$

$$\begin{array}{l} 6x - 5y - 4z + 8 = 0, \\ 14. \quad 6x + 5y + 3z + 4 = 0. \end{array}$$

$$\begin{array}{l} x + 5y - z - 5 = 0, \\ 15. \quad 2x - 5y + 2z + 5 = 0. \end{array}$$

$$\begin{array}{l} 2x - 3y + z + 6 = 0, \\ 16. \quad x - 3y - 2z + 3 = 0. \end{array}$$

$$\begin{array}{l} 5x + y + 2z + 4 = 0, \\ 17. \quad x - y - 3z + 2 = 0. \end{array}$$

$$\begin{array}{l} 4x + y + z + 2 = 0, \\ 18. \quad 2x - y - 3z - 8 = 0. \end{array}$$

$$\begin{array}{l} 2x + y - 3z - 2 = 0, \\ 19. \quad 2x - y + z + 6 = 0. \end{array}$$

$$\begin{array}{l} x + 5y - z + 11 = 0, \\ 20. \quad x - y + 2z - 1 = 0. \end{array}$$

$$\begin{array}{l} x + y - 2z - 2 = 0, \\ 21. \quad x - y + z + 2 = 0. \end{array}$$

$$\begin{array}{l} x - y + z - 2 = 0, \\ 22. \quad x - 2y - z + 4 = 0. \end{array}$$

$$\begin{array}{l} 6x - 7y - z - 2 = 0, \\ 23. \quad x + 7y - 4z - 5 = 0. \end{array}$$

$$\begin{array}{l} x + 5y + 2z - 5 = 0, \\ 24. \quad 2x - 5y - z + 5 = 0. \end{array}$$

$$\begin{array}{l} x - 3y + z + 2 = 0, \\ 25. \quad x + 3y + 2z + 14 = 0. \end{array}$$

$$\begin{array}{l} 2x + 3y - 2z + 6 = 0, \\ 26. \quad x - 3y + z + 3 = 0. \end{array}$$

$$\begin{array}{l} 3x + 4y + 3z + 1 = 0, \\ 27. \quad 2x - 4y - 2z + 4 = 0. \end{array}$$

$$\begin{array}{lll} 3x + 3y + z - 1 = 0, & 6x - 5y + 3z + 8 = 0, & 2x - 3y - 2z + 6 = 0, \\ 28. \quad 2x - 3y - 2z + 6 = 0. & 29. \quad 6x + 5y - 4z + 4 = 0. & 30. \quad x - 3y + z + 3 = 0. \end{array}$$

Javoblar. 12.1 $1 - x = \frac{y - 4}{4} = -\frac{z}{2}$; 12.2 $\frac{x - 4}{-1} = \frac{y}{3} = \frac{z}{6}$; 12.3 $\frac{x}{-1} = \frac{y - 2}{3} = \frac{z}{-2}$;

12.4 $\frac{x + 3}{-3} = \frac{y}{5} = \frac{z}{-9}$; 12.5 $x - 1\frac{1}{3} = \frac{y}{-9} = \frac{z + 2}{-6}$; 12.6 $\frac{x + 6}{-3} = \frac{y + 1}{3} = \frac{z}{-6}$;

12.7 $\frac{x + 1}{-4} = \frac{y - \frac{1}{2}}{-13} = \frac{z}{-20}$; 12.8 $\frac{x + 1}{-1} = \frac{y - 1}{-13} = \frac{z}{-6}$; 12.9 $\frac{x - 8}{-3} = \frac{y - 6}{-2} = \frac{z}{-1}$;

12.10 $\frac{x - 1}{-2} = \frac{y + 6}{-10} = \frac{z}{-6}$; 12.11 $\frac{x + 1}{-3} = \frac{y - \frac{1}{3}}{-7} = \frac{z}{-15}$; 12.12 $\frac{x}{35} = \frac{y - \frac{4}{7}}{2} = \frac{z}{49}$;

12.13 $\frac{x + 1}{2} = \frac{y + 9}{-11} = \frac{z}{9}$; 12.14 $\frac{x + 1}{5} = \frac{y - 0,4}{-42} = \frac{z}{60}$; 12.15 $\frac{x}{5} = \frac{y - 1}{-4} = \frac{z}{-15}$;

12.16 $\frac{x + 3}{9} = \frac{y}{5} = \frac{z}{-3}$; 12.17 $\frac{x + 1}{-1} = \frac{y - 1}{17} = \frac{z}{-6}$; 12.18 $\frac{x - 1}{-2} = \frac{y + 6}{14} = \frac{z}{-6}$;

12.19 $\frac{x + 2}{-2} = \frac{y - 4}{-8} = \frac{z}{-4}$; 12.20 $\frac{x + 1}{9} = \frac{y + 2}{-3} = \frac{z}{-6}$; 12.21 $\frac{x}{-1} = \frac{y - 2}{-3} = \frac{z}{-2}$;

12.22 $\frac{x - 8}{3} = \frac{y - 6}{2} = \frac{z}{-1}$; 12.23 $\frac{x - 1}{35} = \frac{y - \frac{4}{7}}{23} = \frac{z}{49}$; 12.24 $\frac{x}{5} = \frac{y - 1}{5} = \frac{z}{-15}$;

12.25 $\frac{x + 8}{-9} = \frac{y + 2}{-1} = \frac{z}{6}$; 12.26 $\frac{x + 3}{-3} = \frac{y}{-4} = \frac{z}{-9}$; 12.27 $\frac{x + 1}{4} = \frac{y + \frac{1}{2}}{12} = \frac{z}{-20}$;

12.28 $\frac{x + 1}{-3} = \frac{y - \frac{4}{3}}{8} = \frac{z}{-15}$; 12.29 $\frac{x + 1}{5} = \frac{y - 0,4}{42} = \frac{z}{60}$; 12.30 $\frac{x + 3}{9} = \frac{y}{4} = \frac{z}{3}$.

To'g'ri chiziq va tekislikning kesishish nuqtasi

13–masala. To'g'ri chiziq va tekislikning kesishish nuqtasini toping.

$$\begin{array}{l} \frac{x + 2}{-1} = \frac{y - 1}{1} = \frac{z + 3}{2}, \\ x + 2y - z - 2 = 0, \end{array}$$

$$\begin{cases} x = -t - 2, \\ y = t + 1, \\ z = 2t - 3. \end{cases}$$

Tekislik tenglamasiga olib borib qo'yamiz

$$\begin{aligned} (-t - 2) + 2(t + 1) - (2t - 3) - 2 &= 0, \\ -t - 2 + 2t + 2 - 2t + 3 - 2 &= 0, \\ -t + 1 &= 0, \\ t &= 1. \end{aligned}$$

Shunday qilib, izlanayotgan nuqtaning koordinasi $(-3, 2, -1)$.

$$\begin{aligned} 1. \quad \frac{x-2}{-1} = \frac{y-3}{-1} = \frac{z+1}{4}, \quad 2. \quad \frac{x+1}{3} = \frac{y-3}{-4} = \frac{z+1}{5}, \quad 3. \quad \frac{x-1}{-1} = \frac{y+5}{4} = \frac{z-1}{2}, \\ x+2y+3z-14=0. \quad x+2y-5z+20=0. \quad x-3y+7z-24=0. \end{aligned}$$

$$\begin{aligned} 4. \quad \frac{x-1}{1} = \frac{y}{0} = \frac{z+3}{2}, \quad 5. \quad \frac{x-5}{1} = \frac{y-3}{-1} = \frac{z-2}{0}, \quad 6. \quad \frac{x+1}{-3} = \frac{y+2}{2} = \frac{z-3}{-2}, \\ 2x-y+4z=0. \quad 3x+y-5z-12=0. \quad x+3y-5z+9=0. \end{aligned}$$

$$\begin{aligned} 7. \quad \frac{x-1}{-2} = \frac{y-2}{1} = \frac{z+1}{-1}, \quad 8. \quad \frac{x-1}{2} = \frac{y-2}{0} = \frac{z-4}{1}, \quad 9. \quad \frac{x+2}{-1} = \frac{y-1}{1} = \frac{z+4}{-1}, \\ x-2y+5z+17=0. \quad x-2y+4z-19=0. \quad 2x-y+3z+23=0. \end{aligned}$$

$$\begin{aligned} 10. \quad \frac{x+2}{1} = \frac{y-2}{0} = \frac{z+3}{0}, \quad 11. \quad \frac{x-1}{2} = \frac{y-1}{-1} = \frac{z+2}{3}, \quad 12. \quad \frac{x-1}{1} = \frac{y+1}{0} = \frac{z-1}{-1}, \\ 2x-3y-5z-7=0. \quad 4x+2y-z-11=0. \quad 3x-2y-4z-8=0. \end{aligned}$$

$$\begin{aligned} 13. \quad \frac{x+3}{1} = \frac{y-2}{-5} = \frac{z+2}{3}, \quad 14. \quad \frac{x-2}{2} = \frac{y-2}{-1} = \frac{z-4}{3}, \quad 15. \quad \frac{x-3}{-1} = \frac{y-4}{5} = \frac{z-4}{2}, \\ 5x-y+4z+3=0. \quad x+3y+5z-42=0. \quad 7x+y+4z-47=0. \end{aligned}$$

$$16. \frac{x+3}{2} = \frac{y-1}{3} = \frac{z-1}{5}, \quad 17. \frac{x-3}{2} = \frac{y+1}{3} = \frac{z+3}{2}, \quad 18. \frac{x-5}{-2} = \frac{y-2}{0} = \frac{z+4}{-1},$$

$$2x+3y+7z-52=0. \quad 3x+4y+7z-16=0. \quad 2x-5y+4z+24=0.$$

$$19. \frac{x-1}{8} = \frac{y-8}{-5} = \frac{z+5}{12}, \quad 20. \frac{x-3}{1} = \frac{y-1}{-1} = \frac{z+5}{0}, \quad 21. \frac{x-5}{-1} = \frac{y+3}{5} = \frac{z-1}{2},$$

$$x-2y-3z+18=0. \quad x+7y+3z+11=0. \quad 3x+7y-5z-11=0.$$

$$22. \frac{x-1}{7} = \frac{y-2}{1} = \frac{z-6}{-1}, \quad 23. \frac{x-3}{1} = \frac{y+2}{-1} = \frac{z-8}{0}, \quad 24. \frac{x+1}{-2} = \frac{y}{0} = \frac{z+1}{3},$$

$$4x+y-6z-5=0. \quad 5x+9y+4z-25=0. \quad x+4y+13z-23=0.$$

$$25. \frac{x-1}{6} = \frac{y-3}{1} = \frac{z+5}{3}, \quad 26. \frac{x-2}{4} = \frac{y-1}{-3} = \frac{z+3}{-2}, \quad 27. \frac{x-1}{2} = \frac{y+2}{-5} = \frac{z-3}{-2},$$

$$3x-2y+5z-3=0. \quad 3x-y+4z=0. \quad x+2y-5z+16=0.$$

$$28. \frac{x-1}{1} = \frac{y-3}{0} = \frac{z+2}{-2}, \quad 29. \frac{x+3}{0} = \frac{y-2}{-3} = \frac{z+5}{11}, \quad 30. \frac{x-7}{3} = \frac{y-3}{1} = \frac{z+1}{-2},$$

$$3x-7y-2z+7=0. \quad 5x+7y+9z-32=0. \quad 2x+y+7z-3=0.$$

Javoblar. 13.1 (1; 2; 3); 13.2 (2; -1; 4); 13.3 (0; -1; 3); 13.4 (2; 0; -1); 13.5 (7; 1; 2);

13.6 (-4; 0; 1); 13.7 (7; -1; $-5\frac{1}{5}$); 13.8 (3; 2; 5); 13.9 (-3; 2; -5); 13.10 (-1; 2; -3);

13.11 (3; 0; 1); 13.12 (2; -1; 0); 13.13 (-2; -3; 1); 13.14 (4; 1; 7); 13.15 (2; 9; 6);

13.16 (-1; 4; 6); 13.17 (5; 2; -1); 13.18 (3; 2; -5); 13.19 (9; 3; 7); 13.20 (4; 0; -5);

13.21 (4; 2; 3); 13.22 (8; 3; 5); 13.23 (4; -3; 8); 13.24 (-3; 0; 2); 13.25 (7; 4; -2);

13.26 (6; -2; -5); 13.27 (3; -7; 1); 13.28 (2; 3; -4); 13.29 (-3; -1; 6); 13.30

(10; 4; -3).

To'g'ri chiziqqa nisbatan nuqtalarning simmetrikligi

14-masala. To'g'ri chiziqqa nisbatan, M nuqtaga simmetrik bo'lgan M' nuqtani toping.

$$M(3, 3, 3),$$

$$\frac{x-1}{-1} = \frac{y-1,5}{0} = \frac{z-3}{1}.$$

$$-1 \cdot (x-3) + 0 \cdot (y-3) + 1 \cdot (z-3) = 0,$$

$$-x + z = 0.$$

To'g'ri chiziq va tekislikning kesishish nuqtasini topamiz

$$\frac{x-1}{-1} = \frac{y-1,5}{0} = \frac{z-3}{1} \quad \text{B} \quad \begin{cases} x = -t + 1, \\ y = 1,5, \\ z = t + 3. \end{cases}$$

$$-(-t+1) + (t+3) = 0,$$

$$2t + 2 = 0,$$

$$t = -1.$$

$M_0(2; 1,5; 2)$ – kesishish nuqtasining koordinatasi

Bundan,

$$x_{M_0} = \frac{x_M + x_{M_0}}{2} \quad \text{B} \quad x_{M'} = 2x_{M_0} - x_M = 2 \cdot 2 - 3 = 1,$$

$$y_{M_0} = \frac{y_M + y_{M_0}}{2} \quad \text{B} \quad y_{M'} = 2y_{M_0} - y_M = 2 \cdot 1,5 - 3 = 0,$$

$$z_{M_0} = \frac{z_M + z_{M_0}}{2} \quad \text{B} \quad z_{M'} = 2z_{M_0} - z_M = 2 \cdot 2 - 3 = 1.$$

Demak, $M'(1, 0, 1)$ – izlanayotgan nuqta.

To'g'ri chiziqqa (1–15 variantlar uchun) yoki tekislikka (16–30 variantlar uchun) nisbatan M nuqtaga simmetrik bo'lgan M' nuqtani toping

$$M(0, -3, -2),$$

$$1. \quad \frac{x-1}{1} = \frac{y+1,5}{-1} = \frac{z}{1}.$$

$$M(2, -1, 1),$$

$$2. \quad \frac{x-4,5}{1} = \frac{y+3}{-0,5} = \frac{z-2}{1}.$$

$$M(1, 1, 1),$$

$$3. \frac{x-2}{1} = \frac{y+1,5}{-2} = \frac{z-1}{1}.$$

$$M(1, 0, -1),$$

$$5. \frac{x-3,5}{2} = \frac{y-1,5}{2} = \frac{z}{0}.$$

$$M(-2, -3, 0),$$

$$7. \frac{x+0,5}{1} = \frac{y+1,5}{0} = \frac{z-0,5}{1}.$$

$$M(0, 2, 1),$$

$$9. \frac{x-1,5}{2} = \frac{y}{-1} = \frac{z-2}{1}.$$

$$M(-1, 2, 0),$$

$$11. \frac{x+0,5}{1} = \frac{y+0,75}{-0,2} = \frac{z-2}{2}.$$

$$M(-1, 0, 1),$$

$$13. \frac{x+0,5}{0} = \frac{y-1}{0} = \frac{z-4}{2}.$$

$$M(1, 0, 1),$$

$$15. 4x+6y+4z-25=0.$$

$$M(0, 2, 1),$$

$$17. 2x+4y-3=0.$$

$$M(-1, 2, 0),$$

$$19. 4x-5y-z-7=0.$$

$$M(1, 2, 3),$$

$$4. \frac{x-0,5}{0} = \frac{y+1,5}{-1} = \frac{z-1,5}{1}.$$

$$M(2, 1, 0),$$

$$6. \frac{x-2}{0} = \frac{y+1,5}{-1} = \frac{z+0,5}{1}.$$

$$M(-1, 0, -1),$$

$$8. \frac{x}{-1} = \frac{y-1,5}{0} = \frac{z-2}{1}.$$

$$M(3, -3, -1),$$

$$10. \frac{x-6}{5} = \frac{y-3,5}{4} = \frac{z+0,5}{0}.$$

$$M(2, -2, -3),$$

$$12. \frac{x-1}{-1} = \frac{y+0,5}{0} = \frac{z+1,5}{0}.$$

$$M(0, -3, -2),$$

$$14. \frac{x-0,5}{0} = \frac{y+1,5}{-1} = \frac{z-1,5}{1}.$$

$$M(-1, 0, -1),$$

$$16. 2x+6y-2z+11=0.$$

$$M(2, 1, 0),$$

$$18. y+z+2=0.$$

$$M(2, -1, 1),$$

$$20. x-y+2z-2=0.$$

$$^{21.} \begin{matrix} M(1, 1, 1), \\ x + 4y + 3z + 5 = 0. \end{matrix}$$

$$^{22.} \begin{matrix} M(1, 2, 3), \\ 2x + 10y + 10z - 1 = 0. \end{matrix}$$

$$^{23.} \begin{matrix} M(0, -3, -2), \\ 2x + 10y + 10z - 1 = 0. \end{matrix}$$

$$^{24.} \begin{matrix} M(1, 0, -1), \\ 2y + 4z - 1 = 0. \end{matrix}$$

$$^{25.} \begin{matrix} M(3, -3, -1), \\ 2x - 4y - 4z - 13 = 0. \end{matrix}$$

$$^{26.} \begin{matrix} M(-2, -3, 0), \\ x + 5y + 4 = 0. \end{matrix}$$

$$^{27.} \begin{matrix} M(2, -2, -3), \\ y + z + 2 = 0. \end{matrix}$$

$$^{28.} \begin{matrix} M(-1, 0, 1), \\ 2x + 4y - 3 = 0. \end{matrix}$$

$$^{29.} \begin{matrix} M(3, 3, 3), \\ 8x + 6y + 8z - 22 = 0. \end{matrix}$$

$$^{30.} \begin{matrix} M(-2, 0, 3), \\ 2x - 2y + 10z + 1 = 0. \end{matrix}$$

Javoblar. 14.1 $M \notin 1; 1; 1$); 14.2 $M \notin 3; -3; -1$); 14.3 $M \notin 1; 0; -1$); 14.4 $M \notin 0; -3; -2$);
 14.5 $M \notin 2; -1; 1$); 14.6 $M \notin 2; -2; -3$); 14.7 $M \notin -1; 0; -1$); 14.8 $M \notin 3; 3; 3$);
 14.9 $M \notin -1; 0; 1$); 14.10 $M \notin -1; 2; 0$); 14.11 $M \notin -2; -3; 0$); 14.12 $M \notin 2; 1; 0$);
 14.13 $M \notin 0; 2; 1$); 14.14 $M \notin 1; 2; 3$); 14.15 $M \notin 3; 3; 3$); 14.16 $M \notin -2; -3; 0$);
 14.17 $M \notin -1; 0; 1$); 14.18 $M \notin 2; -2; -3$); 14.19 $M \notin 3; -3; -1$); 14.20 $M \notin 1; 0; -1$); 14.21
 $M \notin 0; -3; -2$); 14.22 $M \notin 0; -3; -2$); 14.23 $M \notin 1; 2; 3$); 14.24 $M \notin 1; 1; 1$);
 14.25 $M \notin 2; -1; 1$); 14.26 $M \notin -1; -2; 0$); 14.27 $M \notin 2; 1; 0$); 14.28 $M \notin 0; 2; 1$);
 14.29 $M \notin -1; 0; -1$); 14.30 $M \notin -3; 1; -2$).

II BOB. Limitlar nazariyasi

Limitlar nazariyasi bo'limida, siz sonli ketma-ketlik tushunchasi, nuqtada son va funksiyaning uzluksizligi, turli limitlarni hisoblashning maxsus yo'llari bilan tanishasiz.

Ta'rif. Agar a nuqtaning ixtiyoriy $(a - \epsilon, a + \epsilon)$ atrofi " $\epsilon > 0$ olinganda ham $\{x_n\}$ ketma-ketlikning biror hadidan boshlab, keyingi

barcha hadlari shu trofga tegishli bo'lsa, a son $\{x_n\}$ ketma – ketlikning limiti deyiladi va

$$\lim_{n \rightarrow \infty} x_n = a$$

kabi belgilanadi.

Ketma-ketlikning ti'rifini quyidagicha ta'riflash ham mumkin.

Ta'rif. Agar " $\epsilon > 0$ son olinganda ham shunday natural n_0 son ($n_0 \in \mathbb{N}$) topilsaki, barcha $n > n_0$ lar uchun

$$|x_n - a| < \epsilon$$

tengsizlik bajarilsa, a son $\{x_n\}$ ketma–ketlikning limiti deyiladi.

1–masala. $\lim_{n \rightarrow \infty} a_n = a$ ekanligini isbotlang ($N(\epsilon)$ ko'rsating).

$$a_n = \frac{3n - 2}{2n - 1}, \quad a = \frac{3}{2}.$$

limitning ta'rifiga ko'ra:

$$\epsilon > 0: \exists N(\epsilon): \forall n: n \geq N(\epsilon): |a_n - a| < \epsilon$$

$$\left| \frac{3n - 2}{2n - 1} - \frac{3}{2} \right| < \epsilon;$$

$$\left| \frac{2(3n - 2) - 3(2n - 1)}{2(2n - 1)} \right| < \epsilon;$$

$$\left| \frac{6n - 4 - 6n + 3}{2(2n - 1)} \right| < \epsilon;$$

$$\left| \frac{-1}{2(2n - 1)} \right| < \epsilon;$$

$$\left| \frac{1}{2(2n - 1)} \right| < \epsilon;$$

$$\forall n \geq N: \frac{1}{2(2n - 1)} > 0 \text{ ekanligidan,}$$

$$\frac{1}{2(2n-1)} < \epsilon;$$

$$2n-1 > \frac{1}{2\epsilon};$$

$$n > \frac{1}{2\epsilon} + \frac{1}{2};$$

$$N(\epsilon) = \frac{1}{2\epsilon} + \frac{1}{2} + 1 = \frac{3}{2\epsilon} + \frac{1}{2} = \frac{1+6\epsilon}{4\epsilon}$$

" $n > N(\epsilon)$ da $|a_n - a| < \epsilon$ tengsizlik bajarilishidan,

$$\lim_{n \rightarrow \infty} \frac{3n-2}{2n-1} = \frac{3}{2}$$

kelib chiqadi.

$$1. a_n = \frac{4n-1}{2n+1}, a = 2.$$

$$2. a_n = \frac{7n+4}{2n+1}, a = \frac{7}{2}.$$

$$3. a_n = \frac{2n-5}{3n+1}, a = \frac{2}{3}.$$

$$4. a_n = \frac{7n-1}{n+1}, a = 7.$$

$$5. a_n = \frac{4n^2+1}{3n^2+2}, a = \frac{4}{3}.$$

$$6. a_n = \frac{9-n^3}{1+2n^3}, a = -\frac{1}{2}.$$

$$7. a_n = \frac{4n-3}{2n+1}, a = 2.$$

$$8. a_n = \frac{1-2n^2}{2+4n^2}, a = -\frac{1}{2}.$$

$$9. a_n = -\frac{5n}{n+1}, a = -5.$$

$$10. a_n = \frac{n+1}{1-2n}, a = -\frac{1}{2}.$$

$$11. a_n = \frac{2n+1}{3n-5}, a = \frac{2}{3}.$$

$$12. a_n = \frac{1-2n^2}{n^2+3}, a = -2.$$

$$13. a_n = \frac{3n^2}{2-n^2}, a = -3.$$

$$14. a_n = \frac{n}{3n-1}, a = \frac{1}{3}.$$

$$15. a_n = \frac{3n^3}{n^3 - 1}, \quad a = 3.$$

$$16. a_n = \frac{4 + 2n}{1 - 3n}, \quad a = -\frac{2}{3}.$$

$$17. a_n = \frac{5n + 15}{6 - n}, \quad a = -5.$$

$$18. a_n = \frac{3 - n^2}{4 + 2n^2}, \quad a = -\frac{1}{2}.$$

$$19. a_n = \frac{2n - 1}{2 - 3n}, \quad a = -\frac{2}{3}.$$

$$20. a_n = \frac{3n - 1}{5n + 1}, \quad a = \frac{3}{5}.$$

$$21. a_n = \frac{4n - 3}{2n + 1}, \quad a = 2.$$

$$22. a_n = \frac{1 - 2n^2}{2 + 4n^2}, \quad a = -\frac{1}{2}.$$

$$23. a_n = \frac{5n + 1}{10n - 3}, \quad a = \frac{1}{2}.$$

$$24. a_n = \frac{2 - 2n}{3 + 4n}, \quad a = -\frac{1}{2}.$$

$$25. a_n = \frac{23 - 4n}{2 - n}, \quad a = 4.$$

$$26. a_n = \frac{1 + 3n}{6 - n}, \quad a = -3.$$

$$27. a_n = \frac{2n + 3}{n + 5}, \quad a = 2.$$

$$28. a_n = \frac{3n^2 + 2}{4n^2 - 1}, \quad a = \frac{3}{4}.$$

$$29. a_n = \frac{2 - 3n^2}{4 + 5n^2}, \quad a = -\frac{3}{5}.$$

$$30. a_n = \frac{2n^3}{n^3 - 2}, \quad a = 2.$$

Javoblar. 1.1 $\frac{3 + e^{-n}}{2e^{-n}}$; 1.2 $\frac{1 + 2e^{-n}}{4e^{-n}}$; 1.3 $\frac{17 + 6e^{-n}}{9e^{-n}}$; 1.4 $\frac{8e^{-n}}{e^{-n}}$; 1.5 $\frac{1}{3} \left| \frac{5}{3e^{-n}} - 2 \right| e^{-n} + 1$;

1.6 $\frac{1}{2} \sqrt{\frac{1}{2e^{-n}} - 1} e^{-n} + 1$; 1.7 $\frac{5 + e^{-n}}{2e^{-n}}$; 1.8 $\frac{1}{2} \left| \frac{1}{e^{-n}} - 1 \right| e^{-n} + 1$; 1.9 $\frac{5e^{-n}}{e^{-n}}$; 1.10 $3 \times \frac{1}{4e^{-n}} + \frac{1}{2} e^{-n}$;

1.11 $2 + \frac{13 + 6e^{-n}}{9e^{-n}}$; 1.12 $\frac{1}{e^{-n}} \sqrt{\frac{7}{e^{-n}} - 3} e^{-n} + 1$; 1.13 $\frac{1}{e^{-n}} \sqrt{\frac{6}{e^{-n}} + 2} e^{-n} + 1$; 1.14 $\frac{1 + 12e^{-n}}{9e^{-n}}$;

1.15 $\frac{1}{e^{-n}} \sqrt{1 + \frac{3}{e^{-n}}} e^{-n} + 1$; 1.16 $\frac{14 + 12e^{-n}}{9e^{-n}}$; 1.17 $7 + \frac{45e^{-n}}{e^{-n}}$; 1.18 $\frac{1}{e^{-n}} \sqrt{\frac{5}{2e^{-n}} - 2} e^{-n} + 1$; 1.19 $\frac{1 + 15e^{-n}}{9e^{-n}}$;

1.20 $\frac{8 + 20e^{-n}}{25e^{-n}}$; 1.21 $\frac{5 + e^{-n}}{2e^{-n}}$; 1.22 $\frac{1}{2} \left| \frac{1}{e^{-n}} - 1 \right| e^{-n} + 1$; 1.23 $\frac{5 + 14e^{-n}}{20e^{-n}}$; 1.24 $\frac{7 + 2e^{-n}}{8e^{-n}}$;

$$1.25 \quad 3 + \frac{1}{e} \sqrt[5]{\frac{5}{e}}; \quad 1.26 \quad 7 + \frac{1}{e} \sqrt[9]{\frac{9}{e}}; \quad 1.27 \quad \frac{1}{e} \sqrt[7]{\frac{7}{e}} - 4; \quad 1.28 \quad \frac{1}{e} \sqrt[2]{\left| \frac{11}{4e} + 1 \right|} + 1; \quad 1.29 \quad \frac{1}{e} \sqrt[5]{\frac{1}{5} \left| \frac{22}{5e} - 4 \right|} + 1;$$

$$1.30 \quad \frac{1}{e} \sqrt[3]{2 + \frac{4}{e}} + 1.$$

Sonli ketma-ketlikning limiti

2–masala. Sonli ketma-ketlikning limitini hisoblang.

$$\lim_{n \rightarrow \infty} \frac{(6-n)^2 - (6+n)^2}{(6+n)^2 - (1-n)^2} = \lim_{n \rightarrow \infty} \frac{(36 - 12n + n^2) - (36 + 12n + n^2)}{(36 + 12n + n^2) - (1 - 2n + n^2)} =$$

$$= \lim_{n \rightarrow \infty} \frac{-24n}{14n + 35} = \lim_{n \rightarrow \infty} \frac{-24}{14 + 35/n} = -\frac{24}{14} = -\frac{12}{7}.$$

$$1. \lim_{n \rightarrow \infty} \frac{(3-n)^2 + (3+n)^2}{(3-n)^2 - (3+n)^2}.$$

$$2. \lim_{n \rightarrow \infty} \frac{(3-n)^4 + (2-n)^4}{(1-n)^4 - (1+n)^4}.$$

$$3. \lim_{n \rightarrow \infty} \frac{(3-n)^4 - (2-n)^4}{(1-n)^3 - (1+n)^3}.$$

$$4. \lim_{n \rightarrow \infty} \frac{(1-n)^4 - (1+n)^4}{(1+n)^3 - (1-n)^3}.$$

$$5. \lim_{n \rightarrow \infty} \frac{(n+1)^3 - (n+1)^2}{(n-1)^3 - (n+1)^3}.$$

$$6. \lim_{n \rightarrow \infty} \frac{(1+2n)^3 - 8n^3}{(1+2n)^2 + 4n^2}.$$

$$7. \lim_{n \rightarrow \infty} \frac{(3-4n)^2}{(n-3)^3 - (n+3)^3}.$$

$$8. \lim_{n \rightarrow \infty} \frac{(3-n)^3}{(n+1)^2 - (n+1)^3}.$$

$$9. \lim_{n \rightarrow \infty} \frac{(n+1)^2 + (n-1)^2 - (n+2)^3}{(4-n)^3}.$$

$$10. \lim_{n \rightarrow \infty} \frac{2(n+1)^2 + (n-2)^3}{n^2 + 2n - 3}.$$

$$11. \lim_{n \rightarrow \infty} \frac{(n+1)^3 + (n+2)^3}{(n+4)^3 + (n+5)^3}.$$

$$12. \lim_{n \rightarrow \infty} \frac{(n+3)^3 + (n+4)^3}{(n+3)^4 - (n+4)^4}.$$

$$13. \lim_{n \rightarrow \infty} \frac{(n+1)^4 - (n-1)^4}{(n+1)^3 + (n-1)^3}.$$

$$14. \lim_{n \rightarrow \infty} \frac{8n^3 - 2n}{(n+1)^4 - (n-1)^4}.$$

$$15. \lim_{n \rightarrow \infty} \frac{(n+6)^3 - (n+1)^3}{(2n+3)^2 + (n+4)^2}.$$

$$16. \lim_{n \rightarrow \infty} \frac{(2n-3)^3 - (n+5)^3}{(3n-1)^3 + (2n+3)^3}.$$

$$17. \lim_{n \in \mathbb{N}} \frac{(n+10)^2 + (3n+1)^2}{(n+6)^3 - (n+1)^3}.$$

$$18. \lim_{n \in \mathbb{N}} \frac{(2n+1)^3 + (3n+2)^3}{(2n+3)^3 - (n-7)^3}.$$

$$19. \lim_{n \in \mathbb{N}} \frac{(n+7)^3 - (n+2)^3}{(3n+2)^2 + (4n+1)^2}.$$

$$20. \lim_{n \in \mathbb{N}} \frac{(2n+1)^3 - (2n+3)^3}{(2n+1)^2 + (2n+3)^2}.$$

$$21. \lim_{n \in \mathbb{N}} \frac{n^3 - (n-1)^3}{(n+1)^4 - n^4}.$$

$$22. \lim_{n \in \mathbb{N}} \frac{(n+2)^4 - (n-2)^4}{(n+5)^2 + (n-5)^2}.$$

$$23. \lim_{n \in \mathbb{N}} \frac{(n+1)^4 - (n-1)^4}{(n+1)^3 + (n-1)^3}.$$

$$24. \lim_{n \in \mathbb{N}} \frac{(n+1)^3 - (n-1)^3}{(n+1)^2 - (n-1)^2}.$$

$$25. \lim_{n \in \mathbb{N}} \frac{(n+1)^3 - (n-1)^3}{(n+1)^2 - (n-1)^2}.$$

$$26. \lim_{n \in \mathbb{N}} \frac{(n+2)^3 + (n-2)^3}{n^4 + 2n^2 - 1}.$$

$$27. \lim_{n \in \mathbb{N}} \frac{(n+1)^3 + (n-1)^2}{n^3 - 3n}.$$

$$28. \lim_{n \in \mathbb{N}} \frac{(n+1)^3 + (n-1)^3}{n^3 + 1}.$$

$$29. \lim_{n \in \mathbb{N}} \frac{(n+2)^2 - (n-2)^2}{(n+3)^2}.$$

$$30. \lim_{n \in \mathbb{N}} \frac{(2n+1)^2 - (n+1)^2}{n^2 + n + 1}.$$

Javoblar. 2.1 - ∞ ; 2.2 $\frac{1}{2}$; 2.3 2; 2.4 - 4; 2.5 - ∞ ; 2.6 1,5; 2.7 - $\frac{8}{9}$; 2.8 1; 2.9 0; 2.10 $+\infty$;

2.11 1; 2.12 - $\frac{1}{2}$; 2.13 4; 2.14 1; 2.15 3; 2.16 $\frac{1}{5}$; 2.17 $\frac{2}{3}$; 2.18 5; 2.19 $\frac{3}{5}$; 2.20 - 3;

2.21 0; 2.22 $+\infty$; 2.23 4; 2.24 $+\infty$; 2.25 3; 2.26 0; 2.27 2; 2.28 2; 2.29 0; 2.30 3.

Sonli ketma-ketlikning limiti

3-masala. Sonli ketma-ketlikning limitini hisoblang.

$$\lim_{n \in \mathbb{N}} \frac{\sqrt[3]{n^2 - 1} + 7n^3}{\sqrt[4]{n^{12} + n + 1} - n} = \lim_{n \in \mathbb{N}} \frac{\sqrt[3]{\frac{1}{n^7} - \frac{1}{n^9}} + 7}{\sqrt[4]{1 + \frac{1}{n^{11}} + \frac{1}{n^{12}} - \frac{1}{n^2}}} = 7.$$

$$1. \lim_{n \in \mathbb{N}} \frac{n^3 \sqrt{5n^2} + \sqrt[4]{9n^8 + 1}}{(n + \sqrt{n}) \sqrt{7 - n + n^2}}.$$

$$2. \lim_{n \in \mathbb{N}} \frac{\sqrt{n-1} + \sqrt{n^2 + 1}}{\sqrt[3]{3n^3 + 3} + \sqrt[4]{n^5 + 1}}.$$

$$3. \lim_{n^{\circledast} \nexists} \frac{\sqrt{n^3+1} - \sqrt{n-1}}{\sqrt[3]{n^3+1} - \sqrt{n-1}}.$$

$$4. \lim_{n^{\circledast} \nexists} \frac{\sqrt{3n-1} + \sqrt[3]{125n^3+n}}{\sqrt[3]{n} - n}.$$

$$5. \lim_{n^{\circledast} \nexists} \frac{n^5\sqrt{n} - \sqrt[3]{27n^6+n^2}}{(n + \sqrt[4]{n})\sqrt{9+n^2}}.$$

$$6. \lim_{n^{\circledast} \nexists} \frac{\sqrt{n+2} - \sqrt{n^2+2}}{\sqrt[4]{4n^4+1} - \sqrt[3]{n^4-1}}.$$

$$7. \lim_{n^{\circledast} \nexists} \frac{\sqrt{n^4+2} + \sqrt{n-2}}{\sqrt[4]{n^4+2} + \sqrt{n-2}}.$$

$$8. \lim_{n^{\circledast} \nexists} \frac{6n^3 - \sqrt{n^5+1}}{\sqrt{4n^6+3} - n}.$$

$$9. \lim_{n^{\circledast} \nexists} \frac{\sqrt{5n+2} - \sqrt[5]{8n^3+5}}{\sqrt[4]{n+7} - n}.$$

$$10. \lim_{n^{\circledast} \nexists} \frac{n^4\sqrt{3n+1} + \sqrt{81n^4 - n^2 + 1}}{(n + \sqrt[3]{n})\sqrt{5 - n + n^2}}.$$

$$11. \lim_{n^{\circledast} \nexists} \frac{\sqrt{n+3} - \sqrt{n^2-3}}{\sqrt[3]{n^5-4} - \sqrt[4]{n^4+1}}.$$

$$12. \lim_{n^{\circledast} \nexists} \frac{\sqrt{n^3+3} - \sqrt{n-3}}{\sqrt[5]{n^5+3} + \sqrt{n-3}}.$$

$$13. \lim_{n^{\circledast} \nexists} \frac{\sqrt[3]{n} - 9n^2}{3n - \sqrt[4]{9n^8+1}}.$$

$$14. \lim_{n^{\circledast} \nexists} \frac{\sqrt{4n+1} - \sqrt[3]{27n^3+4}}{\sqrt[4]{n} - \sqrt[3]{n^5+n}}.$$

$$15. \lim_{n^{\circledast} \nexists} \frac{n^3\sqrt{7n} - \sqrt[4]{81n^8-1}}{(n + 4\sqrt{n})\sqrt{n^2-5}}.$$

$$16. \lim_{n^{\circledast} \nexists} \frac{\sqrt[3]{n^3-7} + \sqrt[3]{n^2+4}}{\sqrt[4]{n^5+5} + \sqrt{n}}.$$

$$17. \lim_{n^{\circledast} \nexists} \frac{\sqrt{n^6+4} + \sqrt{n-4}}{\sqrt[5]{n^6+6} - \sqrt{n-6}}.$$

$$18. \lim_{n^{\circledast} \nexists} \frac{4n^2 - \sqrt[4]{n^3}}{\sqrt[3]{n^6+n^3+1} - 5n}.$$

$$19. \lim_{n^{\circledast} \nexists} \frac{\sqrt{n+3} - \sqrt[3]{8n^3+3}}{\sqrt[4]{n+4} - \sqrt[5]{n^5+5}}.$$

$$20. \lim_{n^{\circledast} \nexists} \frac{n^4\sqrt{11n} + \sqrt{25n^4-81}}{(n - 7\sqrt{n})\sqrt{n^2-n+1}}.$$

$$21. \lim_{n^{\circledast} \nexists} \frac{\sqrt[3]{n^2} - \sqrt{n^2+5}}{\sqrt[5]{n^7} - \sqrt{n+1}}.$$

$$22. \lim_{n^{\circledast} \nexists} \frac{\sqrt{n^7+5} - \sqrt{n-5}}{\sqrt[7]{n^7+5} + \sqrt{n-5}}.$$

$$23. \lim_{n^{\circledast} \nexists} \frac{\sqrt[3]{n^2+2} - 5n^2}{n - \sqrt{n^4-n+1}}.$$

$$24. \lim_{n^{\circledast} \nexists} \frac{\sqrt{n+2} - \sqrt[3]{n^3+2}}{\sqrt[7]{n+2} - \sqrt[5]{n^5+2}}.$$

$$25. \lim_{n^{\circledast} \nexists} \frac{n\sqrt{71n} - \sqrt[3]{64n^6+9}}{(n - \sqrt[3]{n})\sqrt{11+n^2}}.$$

$$26. \lim_{n^{\circledast} \nexists} \frac{\sqrt{n+6} - \sqrt{n^2-5}}{\sqrt[3]{n^3+3} + \sqrt[4]{n^3+1}}.$$

$$27. \lim_{n \in \mathbb{N}} \frac{\sqrt{n^8 + 6} - \sqrt{n - 6}}{\sqrt[8]{n^8 + 6} + \sqrt{n - 6}}.$$

$$28. \lim_{n \in \mathbb{N}} \frac{n^2 - \sqrt{n^3 + 1}}{\sqrt[3]{n^6 + 2} - n}.$$

$$29. \lim_{n \in \mathbb{N}} \frac{\sqrt{n+1} - \sqrt[3]{n^3 + 1}}{\sqrt[4]{n+1} - \sqrt[5]{n^5 + 1}}.$$

$$30. \lim_{n \in \mathbb{N}} \frac{n^6 \sqrt{n} + \sqrt[5]{n^{10} + 1}}{(n + \sqrt[4]{n})^3 \sqrt[3]{n^3 - 1}}.$$

Javoblar. 3.1 $\sqrt{3}$; 3.2 $-\infty$; 3.3 ∞ ; 3.4 7; 3.5 5; 3.6 -3 ; 3.7 0; 3.8 ∞ ; 3.9 3; 3.10 2;
3.11 9; 3.12 0; 3.13 ∞ ; 3.14 $3\sqrt{3}$; 3.15 0; 3.16 -3 ; 3.17 0; 3.18 ∞ ; 3.19 4; 3.20 2;
3.21 5; 3.22 0; 3.23 ∞ ; 3.24 5; 3.25 1; 3.26 -4 ; 3.27 -1 ; 3.28 ∞ ; 3.29 1; 3.30 ∞ .

Sonli ketma-ketlikning limiti

4-masala. Sonli ketma-ketlikning limitini hisoblang.

$$\begin{aligned} & \lim_{n \in \mathbb{N}} \frac{\sqrt{(n^2 + 5)(n^4 + 2)} - \sqrt{n^6 - 3n^3 + 5}}{n} = \\ &= \lim_{n \in \mathbb{N}} \frac{(\sqrt{(n^2 + 5)(n^4 + 2)} - \sqrt{n^6 - 3n^3 + 5})(\sqrt{(n^2 + 5)(n^4 + 2)} + \sqrt{n^6 - 3n^3 + 5})}{n(\sqrt{(n^2 + 5)(n^4 + 2)} + \sqrt{n^6 - 3n^3 + 5})} = \\ &= \lim_{n \in \mathbb{N}} \frac{5n^4 + 3n^3 + 2n^2 + 5}{n(\sqrt{(n^2 + 5)(n^4 + 2)} + \sqrt{n^6 - 3n^3 + 5})} = \\ &= \lim_{n \in \mathbb{N}} \frac{\frac{1}{n^4}(5n^4 + 3n^3 + 2n^2 + 5)}{\frac{1}{n^4}n(\sqrt{(n^2 + 5)(n^4 + 2)} + \sqrt{n^6 - 3n^3 + 5})} = \\ &= \lim_{n \in \mathbb{N}} \frac{5 + \frac{3}{n} + \frac{2}{n^2} + \frac{5}{n^4}}{\left(\sqrt{(1 + \frac{5}{n^2})(1 + \frac{2}{n^4})} + \sqrt{1 - \frac{3}{n^3} + \frac{5}{n^6}}\right)} = \frac{5 + 0 + 0 + 0}{\sqrt{(1 + 0)(1 + 0)} + \sqrt{1 - 0 + 0}} = \frac{5}{2}. \end{aligned}$$

$$1. \lim_{n \in \mathbb{N}} n(\sqrt{n^2 + 1} - \sqrt{n^2 - 1}).$$

$$2. \lim_{n \in \mathbb{N}} n(\sqrt{n(n - 2)} - \sqrt{n^2 - 3}).$$

$$3. \lim_{n \in \mathbb{N}} \left(n - \sqrt[3]{n^3 - 5}\right) n \sqrt{n}.$$

$$4. \lim_{n \in \mathbb{N}} (\sqrt{(n^2 + 1)(n^2 - 4)} - \sqrt{n^4 - 9}).$$

5. $\lim_{n \in \mathbb{R}^+} \frac{\sqrt{n^5 - 8} - n\sqrt{n(n^2 + 5)}}{\sqrt{n}}.$
6. $\lim_{n \in \mathbb{R}^+} (\sqrt{n^2 - 3n + 2} - n).$
7. $\lim_{n \in \mathbb{R}^+} \left(n + \sqrt[3]{4 - n^3} \right).$
8. $\lim_{n \in \mathbb{R}^+} (\sqrt{n(n+2)} - \sqrt{n^2 - 2n + 3}).$
9. $\lim_{n \in \mathbb{R}^+} (\sqrt{(n+2)(n+1)} - \sqrt{(n-1)(n+3)}).$
10. $\lim_{n \in \mathbb{R}^+} n^2 (\sqrt{n(n^4 - 1)} - \sqrt{n^5 - 8}).$
11. $\lim_{n \in \mathbb{R}^+} n(\sqrt[3]{5 + 8n^3} - 2n).$
12. $\lim_{n \in \mathbb{R}^+} n^2 (\sqrt[3]{5 + n^3} - \sqrt[3]{3 + n^3}).$
13. $\lim_{n \in \mathbb{R}^+} (\sqrt[3]{(n+2)^2} - \sqrt[3]{(n-3)^2}).$
14. $\lim_{n \in \mathbb{R}^+} \frac{\sqrt{(n+1)^3} - \sqrt{n(n-1)(n-3)}}{\sqrt{n}}.$
15. $\lim_{n \in \mathbb{R}^+} (\sqrt{n^2 + 3n - 2} - \sqrt{n^2 - 3}).$
16. $\lim_{n \in \mathbb{R}^+} \sqrt{n}(\sqrt{n+2} - \sqrt{n-3}).$
17. $\lim_{n \in \mathbb{R}^+} \frac{\sqrt{n(n^5 + 9)} - \sqrt{(n^4 - 1)(n^2 + 5)}}{n}.$
18. $\lim_{n \in \mathbb{R}^+} (\sqrt{n(n+5)} - n).$
19. $\lim_{n \in \mathbb{R}^+} \sqrt{n^3 + 8}(\sqrt{n^3 + 2} - \sqrt{n^3 - 1}).$
20. $\lim_{n \in \mathbb{R}^+} \frac{\sqrt{(n^3 + 1)(n^2 + 3)} - \sqrt{n(n^4 + 2)}}{2\sqrt{n}}.$

$$21. \lim_{n \in \mathbb{N}} (\sqrt{(n^2 + 1)(n^2 + 2)} - \sqrt{(n^2 - 1)(n^2 - 2)}).$$

$$22. \lim_{n \in \mathbb{N}} \frac{\sqrt{(n^5 + 1)(n^2 - 1)} - n\sqrt{n(n^4 + 1)}}{n}.$$

$$23. \lim_{n \in \mathbb{N}} \frac{\sqrt{(n^4 + 1)(n^2 - 1)} - \sqrt{n^6 - 1}}{n}.$$

$$24. \lim_{n \in \mathbb{N}} (n - \sqrt{n(n - 1)}).$$

$$25. \lim_{n \in \mathbb{N}} n^3 (\sqrt[3]{n^2(n^6 + 4)} - \sqrt[3]{(n^8 - 1)}).$$

$$26. \lim_{n \in \mathbb{N}} (n\sqrt{n} - \sqrt{n(n+1)(n+2)}).$$

$$27. \lim_{n \in \mathbb{N}} \sqrt[3]{n} (\sqrt[3]{n^2} - \sqrt[3]{n(n-1)}).$$

$$28. \lim_{n \in \mathbb{N}} \sqrt{n+2} (\sqrt{n+3} - \sqrt{n-4}).$$

$$29. \lim_{n \in \mathbb{N}} n(\sqrt{n^4 + 3} - \sqrt{n^4 - 2}).$$

$$30. \lim_{n \in \mathbb{N}} \sqrt{n(n+1)(n+2)} (\sqrt{n^3 - 3} - \sqrt{n^3 - 2}).$$

Javoblar. 4.1 ∞ ; 4.2 $-\infty$; 4.3 0; 4.4 $-\frac{3}{2}$; 4.5 $-\frac{5}{2}$; 4.6 $-\frac{3}{2}$; 4.7 0; 4.8 2; 4.9 $\frac{1}{2}$; 4.10 $-\infty$;
 4.11 0; 4.12 $\frac{2}{3}$; 4.13 0; 4.14 $\frac{7}{2}$; 4.15 $\frac{3}{2}$; 4.16 $\frac{5}{2}$; 4.17 $-\frac{5}{2}$; 4.18 $\frac{5}{2}$; 4.19 $\frac{3}{2}$; 4.20 $\frac{3}{4}$;
 4.21 3; 4.22 $-\infty$; 4.23 $-\frac{1}{2}$; 4.24 $\frac{1}{2}$; 4.25 0; 4.26 $-\infty$; 4.27 $\frac{1}{3}$; 4.28 $\frac{7}{2}$; 4.29 0; 4.30 $-\frac{1}{2}$.

Sonli ketma-ketlikning limiti

5-masala. Sonli ketma-ketlikning limitini hisoblang.

$$\begin{aligned} \lim_{n \in \mathbb{N}} \frac{(2n+1)! + (2n+2)!}{(2n+3)! - (2n+2)!} &= \lim_{n \in \mathbb{N}} \frac{(2n+1)! + (2n+2)!}{(2n+3) \times (2n+2)! - (2n+2)!} = \\ &= \lim_{n \in \mathbb{N}} \frac{(2n+1)! + (2n+2)!}{(2n+2) \times ((2n+3) - 1)} = \lim_{n \in \mathbb{N}} \frac{(2n+1)! + (2n+2)!}{(2n+2) \times 2n+2} = \end{aligned}$$

$$\begin{aligned}
&= \lim_{n \rightarrow \infty} \frac{(2n+1)!}{(2n+2)!} + \frac{(2n+2)!}{(2n+2)!} \cdot \frac{1}{(2n+2)!} \\
&= \lim_{n \rightarrow \infty} \frac{1}{(2n+2)} + \frac{1}{(2n+2)} = 0 + 0 = 0.
\end{aligned}$$

$$1. \lim_{n \rightarrow \infty} \left(\frac{1}{n^2} + \frac{2}{n^2} + \frac{3}{n^2} + \dots + \frac{n-1}{n^2} \right).$$

$$2. \lim_{n \rightarrow \infty} \frac{(2n+1)! + (2n+2)!}{(2n+3)!}.$$

$$3. \lim_{n \rightarrow \infty} \frac{1+3+5+7+\dots+(2n-1)}{n+1} - \frac{2n+1}{2}.$$

$$4. \lim_{n \rightarrow \infty} \frac{2^{n+1} + 3^{n+1}}{2^n + 3^n}.$$

$$5. \lim_{n \rightarrow \infty} \frac{1+2+3+\dots+n}{\sqrt{9n^4+1}}.$$

$$6. \lim_{n \rightarrow \infty} \frac{1+3+5+\dots+(2n-1)}{1+2+3+\dots+n}.$$

$$7. \lim_{n \rightarrow \infty} \frac{1+3+5+\dots+(2n-1)}{n+3} - n.$$

$$8. \lim_{n \rightarrow \infty} \frac{1+4+7+\dots+(3n-2)}{\sqrt{5n^4+n+1}}.$$

$$9. \lim_{n \rightarrow \infty} \frac{(n+4)! - (n+2)!}{(n+3)!}.$$

$$10. \lim_{n \rightarrow \infty} \frac{(3n-1)! + (3n+1)!}{(3n!)(n-1)}.$$

$$11. \lim_{n \rightarrow \infty} \frac{2^n - 5^{n+1}}{2^{n+1} + 5^{n+2}}.$$

12. $\lim_{n \in \mathbb{N}} \frac{1 + \frac{1}{3} + \frac{1}{3^2} + \dots + \frac{1}{3^n}}{1 + \frac{1}{5} + \frac{1}{5^2} + \dots + \frac{1}{5^n}}.$
13. $\lim_{n \in \mathbb{N}} \frac{1 - 3 + 5 - 7 + 9 - 11 + \dots + (4n - 3) - (4n - 1)}{\sqrt{n^2 + 1} + \sqrt{n^2 + n + 1}}.$
14. $\lim_{n \in \mathbb{N}} \frac{1 - 2 + 3 - 4 + \dots + (2n - 1) - 2n}{n}.$
15. $\lim_{n \in \mathbb{N}} \frac{\sqrt[3]{n^3 + 5} - \sqrt{3n^4 + 2}}{1 + 3 + 5 + \dots + (2n - 1)}.$
16. $\lim_{n \in \mathbb{N}} \frac{3^n - 2^n}{3^{n-1} + 2^n}.$
17. $\lim_{n \in \mathbb{N}} \frac{n + 2}{1 + 2 + 3 + \dots + n} - \frac{2}{3}.$
18. $\lim_{n \in \mathbb{N}} \left(\frac{5}{6} + \frac{13}{36} + \dots + \frac{3^n + 2^n}{6^n} \right).$
19. $\lim_{n \in \mathbb{N}} \frac{2 - 5 + 4 - 7 + \dots + 2n - (2n + 3)}{n + 3}.$
20. $\lim_{n \in \mathbb{N}} \frac{1 + 2 + \dots + n}{n - n^2 + 3}.$
21. $\lim_{n \in \mathbb{N}} \frac{n^2 + \sqrt{n} - 1}{2 + 7 + 12 + \dots + (5n - 3)}.$
22. $\lim_{n \in \mathbb{N}} \left(\frac{3}{4} + \frac{5}{16} + \frac{9}{64} + \dots + \frac{1 + 2^n}{4^n} \right).$
23. $\lim_{n \in \mathbb{N}} \frac{2 + 4 + 6 + \dots + 2n}{1 + 3 + 5 + \dots + (2n - 1)}.$
24. $\lim_{n \in \mathbb{N}} \frac{1 + 5 + 9 + 13 + \dots + (4n - 3)}{n + 1} - \frac{4n + 1}{2}.$

$$25. \lim_{n \in \mathbb{N}} \frac{1 - 2 + 3 - 4 + \dots - 2n}{\sqrt[3]{n^3 + 2n + 2}}.$$

$$26. \lim_{n \in \mathbb{N}} \frac{2^n + 7^n}{2^n - 7^{n-1}}.$$

$$27. \lim_{n \in \mathbb{N}} \frac{n! + (n+2)!}{(n-1)! + (n+2)!}.$$

$$28. \lim_{n \in \mathbb{N}} \frac{3 + 6 + 9 + \dots + 3n}{n^2 + 4}.$$

$$29. \lim_{n \in \mathbb{N}} \left(\frac{7}{10} + \frac{29}{100} + \dots + \frac{2^n + 5^n}{10^n} \right).$$

$$30. \lim_{n \in \mathbb{N}} \frac{2 + 4 + \dots + 2n}{n+3} - n \cdot \frac{0}{0}$$

Javoblar. 5.1 $\frac{1}{2}$; 5.2 0; 5.3 $-\frac{3}{2}$; 5.4 3; 5.5 $\frac{1}{6}$; 5.6 2; 5.7 -3 ; 5.8 $\frac{3}{2\sqrt{5}}$; 5.9 ∞ ; 5.10 3;

5.11 $-\frac{1}{5}$; 5.12 $\frac{6}{5}$; 5.13 -1 ; 5.14 0; 5.15 $-\sqrt{3}$; 5.16 3; 5.17 $-\frac{2}{3}$; 5.18 ∞ ; 5.19 -3 ;

5.20 $-\frac{1}{2}$; 5.21 $\frac{2}{5}$; 5.22 ∞ ; 5.23 1; 5.24 $-\frac{7}{2}$; 5.25 -1 ; 5.26 -7 ; 5.27 1; 5.28 $\frac{3}{2}$; 5.29 ∞ ;

5.30 -2 .

Sonli ketma-ketlikning limiti

6-masala. Sonli ketma-ketlikning limitini hisoblang.

$$\begin{aligned} \lim_{n \in \mathbb{N}} \frac{4n^2 + 4n - 1}{4n^2 + 2n + 3} &= \lim_{n \in \mathbb{N}} \frac{4n^2 + 2n + 3 + 2n - 4}{4n^2 + 2n + 3} = \\ &= \lim_{n \in \mathbb{N}} \left(1 + \frac{2n - 4}{4n^2 + 2n + 3} \right) = \lim_{n \in \mathbb{N}} \left(1 + \frac{1}{\left(\frac{4n^2 + 2n + 3}{2n - 4} \right)} \right) = \end{aligned}$$

$$= \lim_{n \rightarrow \infty} \frac{1}{1 + \left(\frac{4n^2 + 2n + 3}{2n - 4} \right)^{\frac{1}{2n-4}}} =$$

$$= \lim_{n \rightarrow \infty} \frac{1}{1 + \left(\frac{4n^2 + 2n + 3}{2n - 4} \right)^{\frac{1}{2n-4}}} =$$

$$\lim_{t \rightarrow 0} 1 + t^{\frac{1}{t}} = e \text{ ikkinchi ajoyib limitdan foydalangan holda,}$$

$$e^{\lim_{n \rightarrow \infty} \frac{(2n-4)(1-2n)}{4n^2+2n+3}} = e^{\lim_{n \rightarrow \infty} \frac{2n-4n^2-4+8n}{4n^2+2n+3}} = e^{\lim_{n \rightarrow \infty} \frac{-4n^2+10n-4}{4n^2+2n+3}} =$$

$$= e^{\lim_{n \rightarrow \infty} \frac{\frac{1}{n^2}(-4n^2+10n-4)}{\frac{1}{n^2}(4n^2+2n+3)}} = e^{\lim_{n \rightarrow \infty} \frac{-4+\frac{10}{n}-\frac{4}{n^2}}{4+\frac{2}{n}+\frac{3}{n^2}}} = e^{\frac{-4+0-0}{4+0+0}} = e^{-1} = \frac{1}{e}.$$

$$1. \lim_{n \rightarrow \infty} \frac{n^{n+1} - 1}{n - 1}.$$

$$2. \lim_{n \rightarrow \infty} \frac{2n+3}{2n+1}.$$

$$3. \lim_{n \rightarrow \infty} \frac{n^2 - 1}{n^2}.$$

$$4. \lim_{n \rightarrow \infty} \frac{n - 1}{n + 3}.$$

$$5. \lim_{n \rightarrow \infty} \frac{2n^2 + 2}{2n^2 + 1}.$$

$$6. \lim_{n \rightarrow \infty} \frac{3n^2 - 6n + 7}{3n^2 + 20n - 1}.$$

$$7. \lim_{n \rightarrow \infty} \frac{n^2 - 3n + 6}{n^2 + 5n + 1}.$$

$$8. \lim_{n \rightarrow \infty} \frac{n - 10}{n + 1}.$$

$$9. \lim_{n \rightarrow \infty} \frac{6n - 7}{6n + 4}.$$

$$10. \lim_{n \rightarrow \infty} \frac{3n^2 + 4n - 1}{3n^2 + 2n + 7}.$$

$$11. \lim_{n \rightarrow \infty} \frac{n^2 + n + 1}{n^2 + n - 1}.$$

$$12. \lim_{n \rightarrow \infty} \frac{2n^2 + 5n + 7}{2n^2 + 5n + 3}.$$

$$13. \lim_{n \rightarrow \infty} \frac{n - 1}{n + 1} \cdot n^2.$$

$$14. \lim_{n \rightarrow \infty} \frac{5n^2 + 3n - 1}{5n^2 + 3n + 3} \cdot n.$$

$$15. \lim_{n \rightarrow \infty} \frac{3n + 1}{3n - 1} \cdot n^{2n+3}.$$

$$16. \lim_{n \rightarrow \infty} \frac{2n^2 + 7n - 1}{2n^2 + 3n - 1} \cdot n^3.$$

$$17. \lim_{n \rightarrow \infty} \frac{n + 3}{n + 5} \cdot n^{n+4}.$$

$$18. \lim_{n \rightarrow \infty} \frac{n^3 + 1}{n^3 - 1} \cdot n^{2n - n^3}.$$

$$19. \lim_{n \rightarrow \infty} \frac{2n^2 + 21n - 7}{2n^2 + 18n + 9} \cdot n^{2n+1}.$$

$$20. \lim_{n \rightarrow \infty} \frac{10n - 3}{10n - 1} \cdot n^{5n}.$$

$$21. \lim_{n \rightarrow \infty} \frac{3n^2 - 5n}{3n^2 - 5n + 7} \cdot n^{n+1}.$$

$$22. \lim_{n \rightarrow \infty} \frac{n + 3}{n + 1} \cdot n^{n^2}.$$

$$23. \lim_{n \rightarrow \infty} \frac{n^2 - 6n + 5}{n^2 - 5n + 5} \cdot n^{3n+2}.$$

$$24. \lim_{n \rightarrow \infty} \frac{n + 4}{n + 2} \cdot n^n.$$

$$25. \lim_{n \rightarrow \infty} \frac{7n^2 + 18n - 15}{7n^2 + 11n + 15} \cdot n^{n+2}.$$

$$26. \lim_{n \rightarrow \infty} \frac{2n - 1}{2n + 1} \cdot n^{n+1}.$$

$$27. \lim_{n \rightarrow \infty} \frac{n^3 + n + 1}{n^3 + 2} \cdot n^{2n^2}.$$

$$28. \lim_{n \rightarrow \infty} \frac{13n + 3}{13n - 10} \cdot n^{n-3}.$$

$$29. \lim_{n \rightarrow \infty} \frac{2n^2 + 2n + 3}{2n^2 + 2n + 1} \cdot n^{3n^2 - 7}.$$

$$30. \lim_{n \rightarrow \infty} \frac{n + 5}{n - 7} \cdot n^{n/6+1}.$$

Javoblar. 6.1 e^2 ; 6.2 e ; 6.3 0; 6.4 e^{-4} ; 6.5 \sqrt{e} ; 6.6 $e^{\frac{26}{3}}$; 6.7 e^{-4} ; 6.8 e^{-33} ; 6.9 $e^{-\frac{11}{2}}$;
 6.10 $e^{\frac{4}{3}}$; 6.11 e^{-2} ; 6.12 1; 6.13 0; 6.14 $e^{-\frac{4}{5}}$; 6.15 $e^{\frac{4}{3}}$; 6.16 0; 6.17 e^{-2} ; 6.18 e^{-2} ; 6.19 e^3 ;
 6.20 $\frac{1}{e}$; 6.21 1; 6.22 0; 6.23 e^{-3} ; 6.24 e^2 ; 6.25 e ; 6.26 $\frac{1}{e}$; 6.27 e^2 ; 6.28 e ; 6.29 e^3 ; 6.30 e^2 .

7-masala. Isbotlang ($d(e)$ toping):

$$\lim_{x \rightarrow \frac{1}{3}} \frac{15x^2 - 2x - 1}{x - \frac{1}{3}} = 8$$

Koshining funksiya limiti ta'rifiga ko'ra: Agar " $\epsilon > 0$ " son olinganda ham shunday $\delta(\epsilon) > 0$ topilsaki, " $x \in M$ " uchun

$$(0 < |x - a| < \delta(\epsilon)) \Rightarrow (|f(x) - A| < \epsilon)$$

tengsizlik bajarilsa, $A \in R$ soni $f(x)$ funksiyaning A nuqtadagi limiti deyiladi:

$$"\epsilon > 0 : \exists \delta(\epsilon) > 0 : " x \in M : (0 < |x - a| < \delta(\epsilon)) \Rightarrow (|f(x) - A| < \epsilon).$$

Natijad , $x \rightarrow \frac{1}{3}$ da

$$\begin{aligned} \left| \frac{15x^2 - 2x - 1}{x - \frac{1}{3}} - 8 \right| &= \left| \frac{(15x + 3)(x - \frac{1}{3})}{x - \frac{1}{3}} - 8 \right| = \\ &= |15x + 3 - 8| = |15 - x| = 15 \left| x - \frac{1}{3} \right| < \epsilon \end{aligned}$$

yoki

$$\left| x - \frac{1}{3} \right| < \frac{\epsilon}{15}$$

Bu erda $\delta(\epsilon) = \frac{\epsilon}{15}$.

Shunday qilib, $x \rightarrow \frac{1}{3}$ da funksiyaning limiti mavjud va u 8 ga va

$$\delta(\epsilon) = \frac{\epsilon}{15} \text{ teng.}$$

$$\begin{aligned} 1. \quad & f(x) = 5x^2 - 1, \\ & x_0 = 6. \end{aligned}$$

$$\begin{aligned} 2. \quad & f(x) = 4x^2 - 2, \\ & x_0 = 5. \end{aligned}$$

$$\begin{aligned} 3. \quad & f(x) = 3x^2 - 3, \\ & x_0 = 4. \end{aligned}$$

$$\begin{aligned} 4. \quad & f(x) = -2x^2 - 5, \\ & x_0 = 2. \end{aligned}$$

$$\begin{aligned} 5. \quad & f(x) = -3x^2 - 6, \\ & x_0 = 1. \end{aligned}$$

$$\begin{aligned} 6. \quad & f(x) = -4x^2 - 7, \\ & x_0 = 1. \end{aligned}$$

7. $f(x) = -5x^2 - 8,$
 $x_0 = 2.$

9. $f(x) = -4x^2 + 9,$
 $x_0 = 4.$

11. $f(x) = -2x^2 + 7,$
 $x_0 = 6.$

13. $f(x) = 3x^2 + 5,$
 $x_0 = 8.$

15. $f(x) = 5x^2 + 3,$
 $x_0 = 8.$

17. $f(x) = 4x^2 - 1,$
 $x_0 = 6.$

19. $f(x) = 2x^2 - 3,$
 $x_0 = 4.$

21. $f(x) = -3x^2 - 5,$
 $x_0 = 2.$

23. $f(x) = -5x^2 - 7,$
 $x_0 = 1.$

25. $f(x) = -3x^2 - 9,$
 $x_0 = 3.$

27. $f(x) = 2x^2 + 8,$
 $x_0 = 5.$

29. $f(x) = 4x^2 + 6,$
 $x_0 = 7.$

8. $f(x) = -5x^2 - 9,$
 $x_0 = 3.$

10. $f(x) = -3x^2 + 8,$
 $x_0 = 5.$

12. $f(x) = 2x^2 + 6,$
 $x_0 = 7.$

14. $f(x) = 4x^2 + 4,$
 $x_0 = 9.$

16. $f(x) = 5x^2 + 1,$
 $x_0 = 7.$

18. $f(x) = 3x^2 - 2,$
 $x_0 = 5.$

20. $f(x) = -2x^2 - 4,$
 $x_0 = 3.$

22. $f(x) = -4x^2 - 6,$
 $x_0 = 1.$

24. $f(x) = -4x^2 - 8,$
 $x_0 = 2.$

26. $f(x) = -2x^2 + 9,$
 $x_0 = 4.$

28. $f(x) = 3x^2 + 7,$
 $x_0 = 6.$

30. $f(x) = 5x^2 + 5,$
 $x_0 = 8.$

Javoblar. 7.1 $\frac{e}{2}$; 7.2 $\frac{e}{5}$; 7.3 $\frac{e}{3}$; 7.4 $\frac{e}{4}$; 7.5 $\frac{e}{6}$; 7.6 $\frac{e}{6}$; 7.7 $\frac{e}{9}$; 7.8 $\frac{e}{3}$; 7.9 $\frac{e}{3}$; 7.10 $\frac{e}{7}$; 7.11 e ;

7.12 $\frac{e}{2}$; 7.13 $\frac{e}{6}$; 7.14 $\frac{e}{10}$; 7.15 e ; 7.16 e ; 7.17 $\frac{e}{6}$; 7.18 $\frac{e}{6}$; 7.19 $\frac{e}{2}$; 7.20 $\frac{e}{5}$; 7.21 $\frac{e}{2}$;

$$7.22 \frac{e}{2}; 7.23 \frac{e}{2}; 7.24 e; 7.25 \frac{e}{3}; 7.26 \frac{e}{5}; 7.27 \frac{e}{2}; 7.28 \frac{e}{3}; 7.29 \frac{e}{3}; 7.30 \frac{e}{15}.$$

8-masala . $f(x)$ funksiya x_0 nuqtada uzluksiz ekanligi isbotlang ($d(e)$ ni toping.).

$$f(x) = 2x^2 - 4, \quad x_0 = 3.$$

$$|x - x_0| < d(e) \text{ da } |f(x) - f(x_0)| < e,$$

$$|2x^2 - 4 - (2 \times 9 - 4)| = |2x^2 - 18| = 2|x^2 - 9| < e,$$

$$|x^2 - 9| < e/2, \quad |(x - 3)(x + 3)| < e/2 \text{ } \mathbf{B} \quad |x - 3| < e/2 \text{ } \mathbf{B}$$

$$|x - x_0| < d(e) = \frac{e}{2} \text{ } \mathbf{B} \quad |f(x) - f(x_0)| < e \text{ bajariladi.}$$

- | | |
|----------------------------------|----------------------------------|
| 1. $f(x) = 5x^2 - 1, x_0 = 6.$ | 2. $f(x) = 4x^2 - 2, x_0 = 5.$ |
| 3. $f(x) = 3x^2 - 3, x_0 = 4.$ | 4. $f(x) = -2x^2 - 5, x_0 = 2.$ |
| 5. $f(x) = -3x^2 - 6, x_0 = 1.$ | 6. $f(x) = -4x^2 - 7, x_0 = 1.$ |
| 7. $f(x) = -5x^2 - 8, x_0 = 2.$ | 8. $f(x) = -5x^2 - 9, x_0 = 3.$ |
| 9. $f(x) = -4x^2 + 9, x_0 = 4.$ | 10. $f(x) = -3x^2 + 8, x_0 = 5.$ |
| 11. $f(x) = -2x^2 + 7, x_0 = 6.$ | 12. $f(x) = 2x^2 + 6, x_0 = 7.$ |
| 13. $f(x) = 3x^2 + 5, x_0 = 8.$ | 14. $f(x) = 4x^2 + 4, x_0 = 9.$ |
| 15. $f(x) = 5x^2 + 3, x_0 = 8.$ | 16. $f(x) = 5x^2 + 1, x_0 = 7.$ |
| 17. $f(x) = 4x^2 - 1, x_0 = 6.$ | 18. $f(x) = 3x^2 - 2, x_0 = 5.$ |
| 19. $f(x) = 2x^2 - 3, x_0 = 4.$ | 20. $f(x) = -2x^2 - 4, x_0 = 3.$ |
| 21. $f(x) = -3x^2 - 5, x_0 = 2.$ | 22. $f(x) = -4x^2 - 6, x_0 = 1.$ |
| 23. $f(x) = -5x^2 - 7, x_0 = 1.$ | 24. $f(x) = -4x^2 - 8, x_0 = 2.$ |
| 25. $f(x) = -3x^2 - 9, x_0 = 3.$ | 26. $f(x) = -2x^2 + 9, x_0 = 4.$ |
| 27. $f(x) = 2x^2 + 8, x_0 = 5.$ | 28. $f(x) = 3x^2 + 7, x_0 = 6.$ |
| 29. $f(x) = 4x^2 + 6, x_0 = 7.$ | 30. $f(x) = 5x^2 + 5, x_0 = 8.$ |

Javoblar. 8.1 $\frac{e}{5}$; 8.2 $\frac{e}{4}$; 8.3 $\frac{e}{3}$; 8.4 $\frac{e}{2}$; 8.5 $\frac{e}{3}$; 8.6 $\frac{e}{4}$; 8.7 $\frac{e}{5}$; 8.8 $\frac{e}{5}$; 8.9 $\frac{e}{4}$; 8.10 $\frac{e}{3}$; 8.11 $\frac{e}{2}$; 8.12 $\frac{e}{2}$; 8.13 $\frac{e}{3}$; 8.14 $\frac{e}{4}$; 8.15 $\frac{e}{5}$; 8.16 $\frac{e}{5}$; 8.17 $\frac{e}{4}$; 8.18 $\frac{e}{3}$; 8.19 $\frac{e}{2}$; 8.20 $\frac{e}{2}$; 8.21 $\frac{e}{3}$; 8.22 $\frac{e}{4}$; 8.23 $\frac{e}{5}$; 8.24 $\frac{e}{4}$; 8.25 $\frac{e}{3}$; 8.26 $\frac{e}{2}$; 8.27 $\frac{e}{2}$; 8.28 $\frac{e}{3}$; 8.29 $\frac{e}{4}$; 8.30 $\frac{e}{5}$.

Funksiyaning limiti

9-masala . Funksiyaning limitini hisoblang .

$$\begin{aligned}\lim_{x \rightarrow 3} \frac{x^3 - 4x^2 - 3x + 18}{x^3 - 5x^2 + 3x + 9} &= \frac{0}{0} = \lim_{x \rightarrow 3} \frac{(x-3)(x^2 - x - 6)}{(x-3)(x^2 - 2x - 3)} = \\ &= \lim_{x \rightarrow 3} \frac{x^2 - x - 6}{x^2 - 2x - 3} = \frac{0}{0} = \lim_{x \rightarrow 3} \frac{(x-3)(x+2)}{(x-3)(x+1)} = \\ &= \lim_{x \rightarrow 3} \frac{x+2}{x+1} = \frac{3+2}{3+1} = \frac{5}{4} = 1\frac{1}{4}.\end{aligned}$$

$$1. \lim_{x \rightarrow -1} \frac{(x^3 - 2x - 1)(x+1)}{x^4 + 4x^2 - 5}.$$

$$2. \lim_{x \rightarrow -1} \frac{x^3 - 3x - 2}{x + x^2}.$$

$$3. \lim_{x \rightarrow -1} \frac{(x^2 + 3x + 2)^2}{x^3 + 2x^2 - x - 2}.$$

$$4. \lim_{x \rightarrow 1} \frac{(2x^2 - x - 1)^2}{x^3 + 2x^2 - x - 2}.$$

$$5. \lim_{x \rightarrow -3} \frac{(x^2 + 2x - 3)^2}{x^3 + 4x^2 + 3x}.$$

$$6. \lim_{x \rightarrow -1} \frac{(x^3 - 2x - 1)(x+1)}{x^4 + 4x^2 - 5}.$$

$$7. \lim_{x \rightarrow 0} \frac{(1+x^3) - (1+3x)}{x + x^5}.$$

$$8. \lim_{x \rightarrow 1} \frac{x^2 - 2x + 1}{2x^2 - x - 1}.$$

$$9. \lim_{x \rightarrow -1} \frac{x^3 - 3x - 2}{x^2 - x - 2}.$$

$$10. \lim_{x \rightarrow -1} \frac{x^3 + 5x^2 + 7x + 3}{x^3 + 4x^2 + 5x + 2}.$$

$$11. \lim_{x \rightarrow 1} \frac{x^3 - 3x + 2}{x^3 - x^2 - x + 1}.$$

$$12. \lim_{x \rightarrow 1} \frac{x^3 + x^2 - 5x + 3}{x^3 - x^2 - x + 1}.$$

$$13. \lim_{x \rightarrow -1} \frac{x^3 + 4x^2 + 5x + 2}{x^3 - 3x - 2}.$$

$$14. \lim_{x \rightarrow 1} \frac{x^4 - 1}{2x^4 - x^2 - 1}.$$

$$15. \lim_{x \rightarrow -2} \frac{x^3 + 5x^2 + 8x + 4}{x^3 + 3x^2 - 4}.$$

$$16. \lim_{x \rightarrow 2} \frac{x^3 - 5x^2 + 8x - 4}{x^3 - 3x^2 + 4}.$$

$$17. \lim_{x \rightarrow 2} \frac{x^3 - 6x^2 + 12x - 8}{x^3 - 3x^2 + 4}.$$

$$18. \lim_{x \rightarrow -2} \frac{x^3 + 5x^2 + 8x + 4}{x^3 + 7x^2 + 16x + 12}.$$

$$19. \lim_{x \rightarrow -1} \frac{x^3 - 3x - 2}{(x^2 - x - 2)^2}.$$

$$20. \lim_{x \rightarrow 2} \frac{x^3 - 3x - 2}{x - 2}.$$

$$21. \lim_{x \rightarrow -1} \frac{x^3 - 3x - 2}{x^2 + 2x + 1}.$$

$$22. \lim_{x \rightarrow 1} \frac{x^2 - 2x + 1}{x^3 - x^2 - x + 1}.$$

$$23. \lim_{x \rightarrow 1} \frac{x^4 - 1}{2x^4 - x^2 - 1}.$$

$$24. \lim_{x \rightarrow -1} \frac{x^2 + 3x + 2}{x^3 + 2x^2 - x - 2}.$$

$$25. \lim_{x \rightarrow 1} \frac{2x^2 - x - 1}{x^3 + 2x^2 - x - 2}.$$

$$26. \lim_{x \rightarrow -3} \frac{x^2 + 2x - 3}{x^3 + 4x^2 + 3x}.$$

$$27. \lim_{x \rightarrow -1} \frac{x^3 - 2x - 1}{x^4 + 2x + 1}.$$

$$28. \lim_{x \rightarrow 0} \frac{(1+x)^3 - (1+3x)}{x^2 + x^5}.$$

$$29. \lim_{x \rightarrow 1} \frac{x^2 - 1}{2x^2 - x - 1}.$$

$$30. \lim_{x \rightarrow -3} \frac{x^3 + 7x^2 + 15x + 9}{x^3 + 8x^2 + 21x + 18}.$$

Javoblar. 9.1 0; 9.2 0; 9.3 0; 9.4 0; 9.5 0; 9.6 0; 9.7 0; 9.8 0; 9.9 0; 9.10 2; 9.11 $\frac{3}{2}$; 9.12 2;

9.13 $-\frac{1}{3}$; 9.14 $\frac{2}{3}$; 9.15 $\frac{1}{3}$; 9.16 $\frac{1}{3}$; 9.17 0; 9.18 - 1; 9.19 $-\frac{1}{3}$; 9.20 9; 9.21 - 3; 9.22 $\frac{1}{2}$;

9.23 $\frac{2}{3}$; 9.24 $-\frac{1}{2}$; 9.25 $\frac{1}{2}$; 9.26 $-\frac{2}{3}$; 9.27 $-\frac{1}{2}$; 9.28 3; 9.29 $\frac{2}{3}$; 9.30 2.

Funksiyaning limiti

10-masala. Funksiyaning limitini hisoblang .

$$\begin{aligned}
& \lim_{x \rightarrow 3} \frac{\sqrt{x+13} - 2\sqrt{x+1}}{\sqrt[3]{x^2 - 9}} = \lim_{x \rightarrow 3} \frac{(\sqrt{x+13} - 2\sqrt{x+1})(\sqrt{x+13} + 2\sqrt{x+1})}{\sqrt[3]{x^2 - 9}(\sqrt{x+13} + 2\sqrt{x+1})} = \\
& = \lim_{x \rightarrow 3} \frac{x+13 - 4(x+1)}{\sqrt[3]{x^2 - 9}(\sqrt{x+13} + 2\sqrt{x+1})} = \\
& = \lim_{x \rightarrow 3} \frac{-3x+9}{\sqrt[3]{(x-3)(x+3)}(\sqrt{x+13} + 2\sqrt{x+1})} = \\
& = \lim_{x \rightarrow 3} \frac{-3(x-3)}{\sqrt[3]{(x-3)(x+3)}(\sqrt{x+13} + 2\sqrt{x+1})} = \\
& = \lim_{x \rightarrow 3} \frac{-3\sqrt[3]{(x-3)^3}}{\sqrt[3]{(x-3)(x+3)}(\sqrt{x+13} + 2\sqrt{x+1})} = \\
& = \lim_{x \rightarrow 3} \frac{-3\sqrt[3]{(x-3)^2}}{\sqrt[3]{(x+3)}(\sqrt{x+13} + 2\sqrt{x+1})} = \\
& = \lim_{x \rightarrow 3} \frac{-3\sqrt[3]{(3-3)^2}}{\sqrt[3]{(3+3)}(\sqrt{3+13} + 2\sqrt{3+1})} = \\
& = \lim_{x \rightarrow 3} \frac{-3\sqrt[3]{0^2}}{\sqrt[3]{6}(\sqrt{16} + 2\sqrt{4})} = 0.
\end{aligned}$$

$$1. \lim_{x \rightarrow 4} \frac{\sqrt{1+2x} - 3}{\sqrt{x} - 2}.$$

$$2. \lim_{x \rightarrow -8} \frac{\sqrt{1-x} - 3}{2 + \sqrt[3]{x}}.$$

$$3. \lim_{x \rightarrow 1} \frac{\sqrt{x-1}}{\sqrt[3]{x^2-1}}.$$

$$4. \lim_{x \rightarrow 3} \frac{\sqrt{x+13} - 2\sqrt{x+1}}{x^2 - 9}.$$

$$5. \lim_{x \rightarrow -2} \frac{\sqrt[3]{x-6} + 2}{x^3 + 8}.$$

$$6. \lim_{x \rightarrow 16} \frac{\sqrt[4]{x} - 2}{\sqrt{x} - 4}.$$

$$7. \lim_{x \rightarrow 8} \frac{\sqrt{9+2x} - 5}{\sqrt[3]{x} - 2}.$$

$$8. \lim_{x \rightarrow 0} \frac{\sqrt{1-2x+x^2} - (1+x)}{x}.$$

$$9. \lim_{x \rightarrow 0} \frac{\sqrt[3]{8+3x+x^2} - 2}{x+x^2}.$$

$$10. \lim_{x \rightarrow 0} \frac{\sqrt[3]{27+x} - \sqrt[3]{27-x}}{x + 2\sqrt[3]{x^4}}.$$

$$11. \lim_{x \rightarrow 1} \frac{\sqrt[3]{x} - 1}{\sqrt{1+x} - \sqrt{2x}}.$$

$$12. \lim_{x \rightarrow 0} \frac{\sqrt{1+x} - \sqrt{1-x}}{\sqrt[3]{1+x} - \sqrt[3]{1-x}}.$$

$$13. \lim_{x \rightarrow 2} \frac{\sqrt[3]{4x} - 2}{\sqrt{2+x} - \sqrt{2x}}.$$

$$14. \lim_{x \rightarrow 1} \frac{\sqrt{x} - 1}{x^2 - 1}.$$

$$15. \lim_{x \rightarrow 3} \frac{\sqrt[3]{9x} - 3}{\sqrt{3+x} - \sqrt{2x}}.$$

$$16. \lim_{x \rightarrow -2} \frac{\sqrt[3]{x-6} + 2}{x+2}.$$

$$17. \lim_{x \rightarrow 4} \frac{\sqrt[3]{16x} - 4}{\sqrt{4+x} - \sqrt{2x}}.$$

$$18. \lim_{x \rightarrow 8} \frac{\sqrt{9+2x} - 5}{\sqrt[3]{x^2} - 4}.$$

$$19. \lim_{x \rightarrow \frac{1}{2}} \frac{\sqrt[3]{\frac{x}{4}} - \frac{1}{2}}{\sqrt{\frac{1}{2}+x} - \sqrt{2x}}.$$

$$20. \lim_{x \rightarrow \frac{1}{3}} \frac{\sqrt[3]{\frac{x}{9}} - \frac{1}{3}}{\sqrt{\frac{1}{3}+x} - \sqrt{2x}}.$$

$$21. \lim_{x \rightarrow \frac{1}{4}} \frac{\sqrt[3]{\frac{x}{16}} - \frac{1}{4}}{\sqrt{\frac{1}{4}+x} - \sqrt{2x}}.$$

$$22. \lim_{x \rightarrow 0} \frac{\sqrt{1+x} - \sqrt{1-x}}{\sqrt[3]{x}}.$$

$$23. \lim_{x \rightarrow 0} \frac{\sqrt[3]{27+x} - \sqrt[3]{27-x}}{\sqrt[3]{x^2} + \sqrt[5]{x}}.$$

$$24. \lim_{x \rightarrow 0} \frac{\sqrt[3]{8+3x} - x^2 - 2}{\sqrt[3]{x^2} + x^3}.$$

$$25. \lim_{x \rightarrow 0} \frac{\sqrt{1+2x+3x^2} - (1+x)}{\sqrt[3]{x}}.$$

$$26. \lim_{x \rightarrow 8} \frac{\sqrt{9+2x} - 5}{\sqrt[3]{x} - 2}.$$

$$27. \lim_{x \rightarrow 16} \frac{\sqrt[4]{x} - 2}{\sqrt[3]{(\sqrt{x} - 4)^2}}.$$

$$28. \lim_{x \rightarrow -2} \frac{\sqrt[3]{x-6} + 2}{\sqrt[3]{x^3} + 8}.$$

$$29. \lim_{x \rightarrow 4} \frac{\sqrt{x} - 2}{\sqrt[3]{x^2} - 16}.$$

$$30. \lim_{x \rightarrow -8} \frac{10 - x - 6\sqrt{1-x}}{2 + \sqrt[3]{x}}.$$

Javoblar. 10.1 $1\frac{1}{3}$; 10.2 -2 ; 10.3 0 ; 10.4 $-\frac{1}{16}$; 10.5 $\frac{1}{144}$; 10.6 $\frac{1}{4}$; 10.7 $2\frac{2}{5}$; 10.8 -2 ;
 10.9 $\frac{1}{4}$; 10.10 $\frac{2}{27}$; 10.11 $-\frac{2\sqrt{2}}{3}$; 10.12 $\frac{3}{2}$; 10.13 $-1\frac{1}{3}$; 10.14 $\frac{1}{4}$; 10.15 $-\frac{2\sqrt{6}}{3}$; 10.16 $\frac{1}{12}$;
 10.17 $-\frac{4\sqrt{2}}{3}$; 10.18 $0,6$; 10.19 $-\frac{2}{3}$; 10.20 $-\frac{2}{3}\sqrt{\frac{2}{3}}$; 10.21 $-\frac{2}{3}\sqrt{\frac{1}{2}}$; 10.22 0 ; 10.23 0 ;
 10.24 0 ; 10.25 0 ; 10.26 $2\frac{2}{5}$; 10.27 0 ; 10.28 0 ; 10.29 0 ; 10.30 0 .

Funksiyaning limiti

11-masala. Funksiyaning limitini hisoblang .

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\ln(1-3x)}{\sqrt{8x+4}-2} &= \frac{0}{0} = \frac{1}{2} \lim_{x \rightarrow 0} \frac{\ln(1-3x)}{\sqrt{2x+1}-1} = \frac{1}{2} \lim_{x \rightarrow 0} \frac{\frac{-3}{1-3x}}{\frac{1}{\sqrt{2x+1}}} = \\ &= -\frac{1}{2} \lim_{x \rightarrow 0} \frac{3\sqrt{2x+1}}{1-3x} = -\frac{1}{2} \times \frac{3}{1} = -\frac{3}{2}. \end{aligned}$$

$$1. \lim_{x \rightarrow 0} \frac{\ln(1+\sin x)}{\sin 4(x-p)}.$$

$$2. \lim_{x \rightarrow 0} \frac{1-\cos 10(x+p)}{e^{x^2}-1}.$$

$$3. \lim_{x \rightarrow 0} \frac{3x^2-5x}{\sin 3x}.$$

$$4. \lim_{x \rightarrow 0} \frac{1-\cos 2x}{\cos 7x-\cos 3x}.$$

$$5. \lim_{x \rightarrow 0} \frac{4x}{\operatorname{tg}(p(2+x))}.$$

$$6. \lim_{x \rightarrow 0} \frac{2x}{\operatorname{tg}(2p(x+1/2))}.$$

$$7. \lim_{x \rightarrow 0} \frac{1-\cos^3 x}{4x^2}.$$

$$8. \lim_{x \rightarrow 0} \frac{\arcsin 3x}{\sqrt{2+x}-\sqrt{2}}.$$

$$9. \lim_{x \rightarrow 0} \frac{2^{x+1}-2}{\ln(1+4x)}.$$

$$10. \lim_{x \rightarrow 0} \frac{\operatorname{arctg} 2x}{\sin(2p(x+10))}.$$

$$11. \lim_{x \rightarrow 0} \frac{\ln(1-7x)}{\sin(p(x+7))}.$$

$$12. \lim_{x \rightarrow 0} \frac{\cos(x+5p/2)\operatorname{tg} x}{\arcsin 2x^2}.$$

$$13. \lim_{x \rightarrow 0} \frac{1-\sqrt{3x+1}}{\cos(p(x+1/2))}.$$

$$14. \lim_{x \rightarrow 0} \frac{\sin 7x}{x^2+px}.$$

$$15. \lim_{x \rightarrow 0} \frac{\sqrt{4+x}-2}{3\operatorname{arctg} x}.$$

$$16. \lim_{x \rightarrow 0} \frac{2\sin(p(x+1))}{\ln(1+2x)}.$$

$$17. \lim_{x \rightarrow 0} \frac{\cos 2x-\cos x}{1-\cos x}.$$

$$18. \lim_{x \rightarrow 0} \frac{\sqrt{1+x}-1}{\sin(p(x+2))}.$$

$$19. \lim_{x \rightarrow 0} \frac{\sin(5(x+p))}{e^{3x}-1}.$$

$$20. \lim_{x \rightarrow 0} \frac{1-\sqrt{\cos x}}{x \sin x}.$$

$$21. \lim_{x \rightarrow 0} \frac{\arcsin 2x}{\ln(e-x)-1}.$$

$$22. \lim_{x \rightarrow 0} \frac{e^{4x}-1}{\sin(p(x/2+1))}.$$

$$23. \lim_{x \rightarrow 0} \frac{1 + \cos(x - p)}{(e^{3x} - 1)^2}.$$

$$24. \lim_{x \rightarrow 0} \frac{\sin^2 x - \operatorname{tg}^2 x}{x^4}.$$

$$25. \lim_{x \rightarrow 0} \frac{\arcsin 2x}{\ln(e - x) - 1}.$$

$$26. \lim_{x \rightarrow 0} \frac{\operatorname{tg} x - \sin x}{x(1 - \cos 2x)}.$$

$$27. \lim_{x \rightarrow 0} \frac{\ln(x^2 + 1)}{2 - \sqrt{2x^2 + 4}}.$$

$$28. \lim_{x \rightarrow 0} \frac{\operatorname{tg}(p(1 + x/2))}{\ln(x + 1)}.$$

$$29. \lim_{x \rightarrow 0} \frac{e^{4px} - 1}{\sqrt[3]{8 + 24x} - 2}.$$

$$30. \lim_{x \rightarrow 0} \frac{x \sin 2x}{1 + \cos(x - 3p)}.$$

Javoblar. 11.1 $\frac{1}{4}$; 11.2 50; 11.3 $-1\frac{2}{3}$; 11.4 $-\frac{1}{10}$; 11.5 $\frac{4}{p}$; 11.6 $\frac{1}{p}$; 11.7 $\frac{3}{8}$; 11.8 $6\sqrt{2}$;

11.9 $\frac{\ln 2}{2}$; 11.10 $\frac{1}{p}$; 11.11 $\frac{7}{p}$; 11.12 $-\frac{1}{2}$; 11.13 $-\frac{3}{2}$; 11.14 $\frac{3}{p}$; 11.15 $\frac{7}{p}$; 11.16 $\frac{1}{12}$;

11.17 $-p$; 11.18 -3 ; 11.19 $\frac{1}{2p}$; 11.20 $-1\frac{2}{3}$; 11.21 $\frac{1}{4}$; 11.22 $-\frac{2}{3}$; 11.23 $-\frac{8}{p}$; 11.24 $\frac{1}{18}$; 11.25

-1 ; 11.26 $-2e$; 11.27 $\frac{1}{4}$; 11.28 2 ; 11.29 $\frac{p}{2}$; 11.30 $2p$.

Funksiyaning limiti

12-masala. Funksiyaning limitini hisoblang.

$$\lim_{x \rightarrow p} \frac{\cos 3x - \cos x}{\operatorname{tg}^2 2x} = \lim_{x \rightarrow p} \frac{-2 \sin \frac{3x+x}{2} \sin \frac{3x-x}{2}}{\operatorname{tg}^2 2x} =$$

$$= \lim_{x \rightarrow p} \frac{-2 \sin 2x \sin x}{\operatorname{tg}^2 2x} \stackrel{\substack{éx = y + p \\ éx \rightarrow p}}{=} \lim_{y \rightarrow 0} \frac{-2 \sin 2y \sin y}{\operatorname{tg}^2 2y} \stackrel{y \rightarrow 0}{=} \lim_{y \rightarrow 0} \frac{-2 \sin 2(y+p) \sin(y+p)}{\operatorname{tg}^2 2(y+p)} =$$

$$= \lim_{y \rightarrow 0} \frac{-2 \sin 2(y+p) \sin(y+p)}{\operatorname{tg}^2 2(y+p)} = \lim_{y \rightarrow 0} \frac{2 \sin(2y+2p) \sin y}{\operatorname{tg}^2(2y+2p)} =$$

$$\stackrel{\substack{é \sin y \approx y, \text{ при } y \rightarrow 0 \\ é \sin 2y \approx 2y, \text{ при } y \rightarrow 0 \ (2y \rightarrow 0)}}{=} \lim_{y \rightarrow 0} \frac{2 \times 2y \times y}{(2y)^2} = \lim_{y \rightarrow 0} \frac{4y^2}{4y^2} = \lim_{y \rightarrow 0} 1 = 1.$$

$$1. \lim_{x \rightarrow 1} \frac{x^2 - 1}{\ln x}.$$

$$3. \lim_{x \rightarrow p} \frac{1 + \cos 3x}{\sin^2 7x}.$$

$$5. \lim_{x \rightarrow 1} \frac{1 + \cos px}{\operatorname{tg}^2 px}.$$

$$7. \lim_{x \rightarrow p} \frac{\sin^2 x - \operatorname{tg}^2 x}{(x - p)^4}.$$

$$9. \lim_{x \rightarrow p} \frac{\cos 5x - \cos 3x}{\sin^2 x}.$$

$$11. \lim_{x \rightarrow 2} \frac{\sin 7px}{\sin 8px}.$$

$$13. \lim_{x \rightarrow 1} \frac{\sqrt{x^2 - 3x + 3} - 1}{\sin px}.$$

$$15. \lim_{x \rightarrow 1} \frac{3^{5x-3} - 3^{2x^2}}{\operatorname{tg} px}.$$

$$17. \lim_{x \rightarrow \frac{p}{2}} \frac{\ln 2x - \ln p}{\sin(\frac{5x}{2}) \cos x}.$$

$$19. \lim_{x \rightarrow p} \frac{e^p - e^x}{\sin 5x - \sin 3x}.$$

$$21. \lim_{x \rightarrow 2} \frac{1 - 2^{4-x^2}}{2(\sqrt{2x} - \sqrt{3x^2 - 5x + 2})}.$$

$$23. \lim_{x \rightarrow -2} \frac{\operatorname{tg} px}{x + 2}.$$

$$25. \lim_{x \rightarrow \frac{p}{3}} \frac{1 - 2 \cos x}{p - 3x}.$$

$$27. \lim_{x \rightarrow 1} \frac{1 - x^2}{\sin px}.$$

$$2. \lim_{x \rightarrow 1} \frac{\sqrt{x^2 - x + 1} - 1}{\ln x}.$$

$$4. \lim_{x \rightarrow \frac{p}{4}} \frac{1 - \sin 2x}{(p - 4x)^2}.$$

$$6. \lim_{x \rightarrow \frac{p}{2}} \frac{\operatorname{tg} 3x}{\operatorname{tg} x}.$$

$$8. \lim_{x \rightarrow 1} \frac{\sqrt{x^2 - x + 1} - 1}{\operatorname{tg} px}.$$

$$10. \lim_{x \rightarrow 2p} \frac{\sin 7x - \sin 3x}{e^{x^2} - e^{4p^2}}.$$

$$12. \lim_{x \rightarrow 2} \frac{\ln(5 - 2x)}{\sqrt{10 - 3x} - 2}.$$

$$14. \lim_{x \rightarrow p} \frac{x^2 - p^2}{\sin x}.$$

$$16. \lim_{x \rightarrow 4} \frac{2^x - 16}{\sin px}.$$

$$18. \lim_{x \rightarrow \frac{p}{4}} \frac{\ln \operatorname{tg} x}{\cos 2x}.$$

$$20. \lim_{x \rightarrow 2} \frac{\ln(9 - 2x^2)}{\sin 2px}.$$

$$22. \lim_{x \rightarrow 1} \frac{\sqrt[3]{x} - 1}{\sqrt[4]{x} - 1}.$$

$$24. \lim_{x \rightarrow p} \frac{1 - \sin(x/2)}{p - x}.$$

$$26. \lim_{x \rightarrow 2} \frac{\operatorname{arctg}(x^2 - 2x)}{\sin 3px}.$$

$$28. \lim_{x \rightarrow 1} \frac{\cos(px/2)}{1 - \sqrt{x}}.$$

29. $\lim_{x \rightarrow 1} \frac{3 - \sqrt{10 - x}}{\sin 3\pi x}$.

30. $\lim_{x \rightarrow p} \frac{\sin 5x}{\tan 3x}$.

Javoblar. 12.1 2; 12.2 $\frac{1}{2}$; 12.3 $\frac{9}{98}$; 12.4 $\frac{1}{8}$; 12.5 $\frac{1}{2}$; 12.6 $\frac{1}{3}$; 12.7 - 1; 12.8 $\frac{1}{2p}$; 12.9 8;

12.10 $\frac{1}{p > e^{4p^2}}$; 12.11 $\frac{7}{8}$; 12.12 $2\frac{2}{3}$; 12.13 $\frac{1}{2p}$; 12.14 - $2p$; 12.15 $\frac{9\ln 3}{p}$; 12.16 $\frac{16\ln 2}{p}$;

$$\mathbf{12.17} \frac{2\sqrt{2}}{\rho}; \mathbf{12.18} -1; \mathbf{12.19} \frac{e^{\rho}}{2}; \mathbf{12.20} -\frac{4}{\rho}; \mathbf{12.21} -\frac{8\ln 2}{5}; \mathbf{12.22} \frac{4}{3}; \mathbf{12.23} \rho;$$

12.24 0; 12.25 $-\frac{\sqrt{3}}{3}$; 12.26 $\frac{2}{3\rho}$; 12.27 $\frac{2}{\rho}$; 12.28 ρ ; 12.29 $-\frac{1}{18\rho}$; 12.30 $-\frac{5}{3}$.

Funksiyaning limiti

13–masala. Funksiyaning limitini hisoblang .

$$\begin{aligned}
& \lim_{x \rightarrow p} \frac{\sin^{-1}\left(\frac{x^2 - p^2}{2}\right)}{2\sqrt{\sin x + 1}} = \lim_{y \rightarrow 0} \frac{\sin^{-1}\left(\frac{(y+p)^2 - p^2}{2}\right)}{2\sqrt{\sin(y+p) + 1}} \\
&= \lim_{y \rightarrow 0} \frac{\sin^{-1}\left(\frac{y^2 + 2yp + p^2 - p^2}{2}\right)}{2\sqrt{\sin y + 1}} = \lim_{y \rightarrow 0} \frac{\sin^{-1}\left(\frac{y^2 + 2yp}{2}\right)}{2\sqrt{\sin y + 1}} \\
&= \lim_{y \rightarrow 0} \frac{\sin^{-1}\left(y \times \frac{y+2p}{p}\right)}{2\sqrt{\sin y + 1}} = \lim_{y \rightarrow 0} \frac{\sin^{-1}\left(y \times \frac{y+2p}{p}\right)}{2\sqrt{1 - \sin y}} \\
&= \lim_{y \rightarrow 0} \frac{\sin^{-1}\left(y \times \frac{y+2p}{p}\right)}{2(e^{\ln 2(\sqrt{1 - \sin y} - 1)}} = \lim_{y \rightarrow 0} \frac{\sin^{-1}\left(y \times \frac{y+2p}{p}\right)}{2(e^{\ln 2(\sqrt{1 - \sin y} - 1)} - 1)} \\
&= \lim_{y \rightarrow 0} \frac{-y \times \frac{y+2p}{p}}{2\ln 2(\sqrt{1 - \sin y} - 1)} = \lim_{y \rightarrow 0} \frac{-y \times \frac{y+2p}{p}(\sqrt{1 - \sin y} + 1)}{2\ln 2(\sqrt{1 - \sin y} - 1)(\sqrt{1 - \sin y} + 1)} =
\end{aligned}$$

$$\begin{aligned}
&= \lim_{y \rightarrow 0} \frac{y \times \frac{y+2p}{p} (\sqrt{1-\sin y} + 1)}{2 \ln 2(1 - \sin y - 1)} = \lim_{y \rightarrow 0} \frac{y \times \frac{y+2p}{p} (\sqrt{1-\sin y} + 1)}{2 \ln 2 \sin y} = \\
&= [\sin y \gg y] = \lim_{y \rightarrow 0} \frac{y \times \frac{y+2p}{p} (\sqrt{1-\sin y} + 1)}{2 \ln 2 y} = \\
&= \lim_{y \rightarrow 0} \frac{\frac{y+2p}{p} (\sqrt{1-\sin y} + 1)}{2 \ln 2} = \frac{0+2p}{p} \frac{(\sqrt{1-\sin 0} + 1)}{2 \ln 2} = \\
&= \frac{2(\sqrt{1}+1)}{2 \ln 2} = \frac{2}{\ln 2}.
\end{aligned}$$

$$1. \lim_{x \rightarrow p/2} \frac{2^{\cos x} - 1}{\ln \sin x}.$$

$$2. \lim_{x \rightarrow \frac{1}{2}} \frac{(2x-1)^2}{e^{\sin px} - e^{-\sin 3px}}.$$

$$3. \lim_{x \rightarrow 2} \frac{\ln(x - \sqrt[3]{2x-3})}{\sin(px/2) - \sin((x-1)p)}.$$

$$4. \lim_{x \rightarrow 2} \frac{\operatorname{tg} x - \operatorname{tg} 2}{\sin \ln(x-1)}.$$

$$5. \lim_{x \rightarrow \frac{p}{2}} \frac{e^{\operatorname{tg} 2x} - e^{-\sin 2x}}{\sin x - 1}.$$

$$6. \lim_{x \rightarrow \frac{p}{6}} \frac{\ln \sin 3x}{(6x-p)^2}.$$

$$7. \lim_{x \rightarrow 3} \frac{\sin(\sqrt{2x^2-3x-5} - \sqrt{1+x})}{\ln(x-1) - \ln(x+1) + \ln 2}.$$

$$8. \lim_{x \rightarrow 2p} \frac{(x-2p)^2}{\lg(\cos x - 1)}.$$

$$9. \lim_{x \rightarrow \frac{1}{2}} \frac{\ln(4x-1)}{\sqrt{1-\cos px} - 1}.$$

$$10. \lim_{x \rightarrow -2} \frac{\arcsin \frac{x+2}{2}}{3^{\sqrt{2+x+x^2}} - 9}.$$

$$11. \lim_{x \rightarrow 3} \frac{2^{\sin px} - 1}{\ln(x^3 - 6x - 8)}.$$

$$12. \lim_{x \rightarrow p} \frac{\ln \cos 2x}{(1-p/x)^2}.$$

$$13. \lim_{x \rightarrow 2} \frac{\operatorname{tg} \ln(3x-5)}{e^{x+3} - e^{x^2+1}}.$$

$$14. \lim_{x \rightarrow 2p} \frac{\ln \cos x}{3^{\sin 2x} - 1}.$$

$$15. \lim_{x \rightarrow 1} \frac{\sqrt[3]{1+\ln^2 x} - 1}{1 + \cos px}.$$

$$16. \lim_{x \rightarrow p} \frac{\cos \frac{x}{2}}{e^{\sin x} - e^{\sin 4x}}.$$

$$17. \lim_{x \rightarrow 3} \frac{\ln(2x - 5)}{e^{\sin px} - 1}.$$

$$18. \lim_{x \rightarrow \frac{\pi}{3}} \frac{e^{\sin^2 6x} - e^{\sin^2 3x}}{\log_3 \cos 6x}.$$

$$19. \lim_{x \rightarrow \frac{\pi}{2}} \frac{e^{\sin 2x} - e^{\lg 2x}}{\ln(2x/p)}.$$

$$20. \lim_{x \rightarrow -2} \frac{\operatorname{tg}(e^{x+2} - e^{x^2-4})}{\operatorname{tg} x + \operatorname{tg} 2}.$$

$$21. \lim_{x \rightarrow 1} \frac{\sqrt{2^x + 7} - \sqrt{2^{x+1} + 5}}{x^3 - 1}.$$

$$22. \lim_{x \rightarrow \pi} \frac{\ln(2 + \cos x)}{(3^{\sin x} - 1)^2}.$$

$$23. \lim_{x \rightarrow \pi} \frac{(x^3 - \pi^3) \sin 5x}{e^{\sin^2 x} - 1}.$$

$$24. \lim_{x \rightarrow -1} \frac{\operatorname{tg}(x+1)}{e^{\sqrt[3]{x^3 - 4x^2 + 6}} - e}.$$

$$25. \lim_{x \rightarrow \pi} \frac{\ln \cos 2x}{\ln \cos 4x}.$$

$$26. \lim_{x \rightarrow \frac{\pi}{2}} \frac{\ln \sin x}{(2x - \pi)^2}.$$

$$27. \lim_{x \rightarrow a} \frac{a^{x^2 - a^2} - 1}{\lg \ln(x/a)}.$$

$$28. \lim_{x \rightarrow -3} \frac{\sin(e^{\frac{\sqrt[3]{1-x^2}}{2}} - e^{\sqrt[3]{x+2}})}{\operatorname{arctg}(x+3)}.$$

$$29. \lim_{x \rightarrow ap} \frac{\ln(\cos(x/a) + 2)}{a^{a^2 p^2 / x^2 - ap/x} - a^{ap/x - 1}}. \quad 30. \lim_{x \rightarrow \pi} \frac{\operatorname{tg}(3^{p/x} - 3)}{3^{\cos(3x/2)} - 1}.$$

Javoblar. 13.1 - $2 \ln 2$; 13.2 $\frac{1}{e^{\pi} p^2}$; 13.3 $\frac{2}{3p}$; 13.4 $\frac{1}{\cos^2(2)}$; 13.5 0; 13.6 $-\frac{1}{8}$; 13.7 8;

13.8 $\frac{8}{p}$; 13.9 1; 13.10 $-\frac{\ln 3 + 1}{27 \ln^2 3}$; 13.11 $-\frac{p \ln 2}{21}$; 13.12 $-2p^2$; 13.13 $-e^{-5}$; 13.14 0;

13.15 $\frac{2}{3p^2}$; 13.16 $\frac{1}{10}$; 13.17 $-\frac{2}{p}$; 13.18 $-\frac{3}{2} \ln 3$; 13.19 $-2p$; 13.20 $5 \cos^2 2$; 13.21 $-\frac{\ln 2}{9}$;

13.22 $\frac{1}{2 \ln^2 3}$; 13.23 $-15p^2$; 13.24 $\frac{3}{11e}$; 13.25 $\frac{1}{4}$; 13.26 $-\frac{1}{8}$; 13.27 $2a^2 > \ln a$; **13.28**

$-\frac{1}{12e}$; 13.29 $\frac{p^2}{2 \ln a}$; 13.30 $-\frac{2}{p}$.

Funksiyaning limiti

14-masala. Funksiyaning limitini hisoblang .

$$\begin{aligned}
& \lim_{x \rightarrow 0} \frac{2^{3x} - 3^{5x}}{\sin 7x - 2x} = \lim_{x \rightarrow 0} \frac{(8^x - 1) - (243^x - 1)}{\sin 7x - 2x} = \\
& = \lim_{x \rightarrow 0} \frac{((e^{\ln 8})^x - 1) - ((e^{\ln 243})^x - 1)}{\sin 7x - 2x} = \\
& = \lim_{x \rightarrow 0} \frac{(e^{x \ln 8} - 1) - (e^{x \ln 243} - 1)}{\sin 7x - 2x} = \\
& = \lim_{x \rightarrow 0} \frac{\frac{1}{x}(e^{x \ln 8} - 1) - (e^{x \ln 243} - 1)}{\frac{1}{x}(\sin 7x - 2x)} = \\
& = \lim_{x \rightarrow 0} \frac{\frac{1}{x}(e^{x \ln 8} - 1) - (e^{x \ln 243} - 1)}{\lim_{x \rightarrow 0} \frac{1}{x}(\sin 7x - 2x)} = \\
& = \frac{\lim_{x \rightarrow 0} \frac{e^{x \ln 8} - 1}{x} - \lim_{x \rightarrow 0} \frac{e^{x \ln 243} - 1}{x}}{\lim_{x \rightarrow 0} \frac{\sin 7x}{x} - \lim_{x \rightarrow 0} \frac{2x}{x}} = \frac{e^{x \ln 8} - 1 \gg x \ln 8}{e^{x \ln 243} - 1 \gg x \ln 243} \frac{\hat{e}}{\hat{e}} = \\
& = \frac{\lim_{x \rightarrow 0} \frac{x \ln 8}{x} - \lim_{x \rightarrow 0} \frac{x \ln 243}{x}}{\lim_{x \rightarrow 0} \frac{7x}{x} - \lim_{x \rightarrow 0} \frac{2x}{x}} = \frac{\lim_{x \rightarrow 0} \ln 8 - \lim_{x \rightarrow 0} \ln 243}{\lim_{x \rightarrow 0} 7 - \lim_{x \rightarrow 0} 2} = \\
& = \frac{\ln 8 - \ln 243}{7 - 2} = \frac{1}{5} \ln \frac{8}{243} = \frac{1}{5} \ln \frac{2^3}{3^5}.
\end{aligned}$$

$$\begin{aligned}
& 1. \lim_{x \rightarrow 0} \frac{7^{2x} - 5^{3x}}{2x - \arctg 3x}. \quad 2. \lim_{x \rightarrow 0} \frac{e^{3x} - e^{-2x}}{2 \arcsin x - \sin x}. \quad 3. \lim_{x \rightarrow 0} \frac{6^{2x} - 7^{-2x}}{\sin 3x - 2x}. \\
& 4. \lim_{x \rightarrow 0} \frac{e^{5x} - e^{3x}}{\sin 2x - \sin x}. \quad 5. \lim_{x \rightarrow 0} \frac{3^{2x} - 5^{3x}}{\arctg x + x^3}. \quad 6. \lim_{x \rightarrow 0} \frac{e^{2x} - e^{3x}}{\arctg x - x^2}. \\
& 7. \lim_{x \rightarrow 0} \frac{3^{5x} - 2^x}{x - \sin 9x}. \quad 8. \lim_{x \rightarrow 0} \frac{e^{4x} - e^{-2x}}{2 \arctg x - \sin x}. \quad 9. \lim_{x \rightarrow 0} \frac{12^x - 5^{-3x}}{2 \arcsin x - x}. \\
& 10. \lim_{x \rightarrow 0} \frac{e^{7x} - e^{-2x}}{\sin x - 2x}. \quad 11. \lim_{x \rightarrow 0} \frac{3^{5x} - 2^{7x}}{\arcsin 2x - x}. \quad 12. \lim_{x \rightarrow 0} \frac{e^{5x} - e^x}{\arcsin x + x^3}.
\end{aligned}$$

$$\begin{aligned}
13. \lim_{x \rightarrow 0} \frac{4^x - 2^{7x}}{\operatorname{tg} 3x - x}. & \quad 14. \lim_{x \rightarrow 0} \frac{e^x - e^{-x}}{\operatorname{tg} 2x - \sin x}. & \quad 15. \lim_{x \rightarrow 0} \frac{10^{2x} - 7^{-x}}{2\operatorname{tg} x - \operatorname{arctg} x}. \\
16. \lim_{x \rightarrow 0} \frac{e^{2x} - e^x}{\sin 3x - \sin 5x}. & \quad 17. \lim_{x \rightarrow 0} \frac{7^{3x} - 3^{2x}}{\operatorname{tg} x + x^3}. & \quad 18. \lim_{x \rightarrow 0} \frac{e^{4x} - e^{2x}}{2\operatorname{tg} x - \sin x}. \\
19. \lim_{x \rightarrow 0} \frac{3^{2x} - 7^x}{\arcsin x - 5x}. & \quad 20. \lim_{x \rightarrow 0} \frac{e^{2x} - e^{-5x}}{2\sin x - \operatorname{tg} x}. & \quad 21. \lim_{x \rightarrow 0} \frac{4^{5x} - 9^{-2x}}{\sin x - \operatorname{tg} x^3}. \\
22. \lim_{x \rightarrow 0} \frac{e^{3x} - e^{2x}}{\sin 3x - \operatorname{tg} 2x}. & \quad 23. \lim_{x \rightarrow 0} \frac{5^{2x} - 2^{3x}}{\sin x + \sin x^2}. & \quad 24. \lim_{x \rightarrow 0} \frac{e^x - e^{3x}}{\sin 3x - \operatorname{tg} 2x}. \\
25. \lim_{x \rightarrow 0} \frac{9^x - 2^{3x}}{\operatorname{arctg} 2x - 7x}. & \quad 26. \lim_{x \rightarrow 0} \frac{e^x - e^{-2x}}{x + \sin x^2}. & \quad 27. \lim_{x \rightarrow 0} \frac{3^{5x} - 2^{-7x}}{2x - \operatorname{tg} x}. \\
28. \lim_{x \rightarrow 0} \frac{e^{2x} - e^x}{\sin 2x - \sin x}. & \quad 29. \lim_{x \rightarrow 0} \frac{e^{2x} - e^x}{x + \operatorname{tg} x^2}. & \quad 30. \lim_{x \rightarrow 0} \frac{2^{3x} - 3^{2x}}{x + \arcsin x^3}.
\end{aligned}$$

Javoblar. 14.1 $\ln \frac{125}{49}$; 14.2 5; 14.3 $\ln(6^2 \times 7^2)$; 14.4 2; 14.5 $\ln \frac{9}{125}$; 14.6 - 1; 14.7 $\frac{1}{8} \ln \frac{2}{243}$;
14.8 6; 14.9 $\ln(12 \times 5^3)$; 14.10 - 9; 14.11 $\ln \frac{2^5}{2^7}$; 14.12 4; 14.13 $\ln \sqrt{\frac{1}{2^5}}$; 14.14 2;
14.15 $\ln 700$; 14.16 $-\frac{1}{2}$; 14.17 $\ln \frac{7^3}{3^2}$; 14.18 2; 14.19 $\ln \sqrt{\frac{7}{9}}$; 14.20 7; 14.21 $\ln(2^{10} \times 9^2)$; 14.22
1; 14.23 $\ln \frac{25}{8}$; 14.24 - 2; 14.25 $-\frac{1}{5} \ln \frac{9}{8}$; 14.26 3; 14.27 $\ln(3^5 \times 2^7)$; 14.28 1;
14.29 1; 14.30 $\ln \frac{2^3}{3^2}$.

Funksiyaning limiti

15–masala. Funksiyaning limitini hisoblang .

$$\begin{aligned}
\lim_{x \rightarrow 1} \frac{e^x - e}{\sin(x^2 - 1)} &= \lim_{y \rightarrow 0} \frac{e^{y+1} - e}{\sin((y+1)^2 - 1)} = \lim_{y \rightarrow 0} \frac{e(e^y - 1)}{\sin(y^2 + 2y + 1 - 1)} = \\
&= \lim_{y \rightarrow 0} \frac{e(e^y - 1)}{\sin(y^2 + 2y)} = \lim_{y \rightarrow 0} \frac{e \cdot y}{y^2 + 2y} = \lim_{y \rightarrow 0} \frac{e}{y + 2} = \frac{e}{0 + 2} = \frac{e}{2}.
\end{aligned}$$

$$1. \lim_{x \rightarrow 0} \frac{e^x + e^{-x} - 2}{\sin^2 x}.$$

$$2. \lim_{x \rightarrow 0} \frac{1 + x \sin x - \cos 2x}{\sin^2 x}.$$

$$3. \lim_{x \rightarrow -1} \frac{x^3 + 1}{\sin(x + 1)}.$$

$$4. \lim_{x \rightarrow a} \frac{\operatorname{tg} x - \operatorname{tga}}{\ln x - \ln a}.$$

$$5. \lim_{x \rightarrow 0} \frac{\sqrt{1 + \operatorname{tg} x} - \sqrt{1 - \sin x}}{x^3}.$$

$$6. \lim_{x \rightarrow 0} \frac{e^{ax} - e^{-bx}}{\sin ax - \sin bx}.$$

$$7. \lim_{x \rightarrow 0} \frac{\sqrt{1 + x \sin x} - 1}{e^{x^2} - 1}.$$

$$8. \lim_{x \rightarrow 0} \frac{x^2(e^x - e^{-x})}{e^{x^3+1} - e}.$$

$$9. \lim_{x \rightarrow \rho/3} \frac{1 - 2 \cos x}{\sin(\rho - 3x)}.$$

$$10. \lim_{x \rightarrow 1} \frac{1 - x^2}{\sin \rho x}.$$

$$11. \lim_{x \rightarrow \rho/4} \frac{\sin x - \cos x}{\ln \operatorname{tg} x}.$$

$$12. \lim_{x \rightarrow b} \frac{a^x - a^b}{x - b}.$$

$$13. \lim_{x \rightarrow 0} \frac{1 - \cos 2x + \operatorname{tg}^2 x}{x \sin 3x}.$$

$$14. \lim_{x \rightarrow 0} \frac{\sin 2x - 2 \sin x}{x \ln \cos 5x}.$$

$$15. \lim_{h \rightarrow 0} \frac{\ln(x+h) + \ln(x-h) + 2 \ln x}{h^2}.$$

$$16. \lim_{x \rightarrow 1} \frac{1 - x}{\log_2 x}.$$

$$17. \lim_{x \rightarrow 0} \frac{e^{\sin 2x} - e^{\sin x}}{\operatorname{tg} x}.$$

$$18. \lim_{x \rightarrow 1} \frac{2^x - 2}{\ln x}.$$

$$19. \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin(x-h)}{h}. \quad 20. \lim_{x \rightarrow 0} \frac{\sqrt{x+2} - \sqrt{2}}{\sin 3x}.$$

$$21. \lim_{h \rightarrow 0} \frac{a^{x+h} + a^{x-h} - 2a^x}{h^2}. \quad 22. \lim_{x \rightarrow 0} \frac{1 - \sqrt{\cos x}}{1 - \cos \sqrt{x}}.$$

$$23. \lim_{x \rightarrow 3} \frac{\sqrt[3]{5+x} - 2}{\sin \pi x}. \quad 24. \lim_{x \rightarrow \pi/6} \frac{2 \sin^2 x + \sin x - 1}{2 \sin^2 x - 3 \sin x + 1}.$$

$$25. \lim_{x \rightarrow 10} \frac{\lg x - 1}{\sqrt{x-9} - 1}. \quad 26. \lim_{x \rightarrow 0} \frac{3^{x+1} - 3}{\ln(1 + x\sqrt{1 + xe^x})}.$$

$$27. \lim_{x \rightarrow 0} \frac{\sqrt{\cos x} - 1}{\sin^2 2x}. \quad 28. \lim_{x \rightarrow 0} \frac{\sin bx - \sin ax}{\ln(\operatorname{tg}(\pi/4 + ax))}.$$

$$29. \lim_{x \rightarrow \pi/2} \frac{1 - \sin^3 x}{\cos^2 x}. \quad 30. \lim_{x \rightarrow 3} \frac{\log_3 x - 1}{\operatorname{tg} \pi x}.$$

Javoblar. 15.1 1; 15.2 3; 15.3 3; 15.4 $\frac{a}{\cos^2 a}$; 15.5 $\frac{1}{4}$; 15.6 1; 15.7 $\frac{1}{2}$; 15.8 $\frac{2}{e}$; 15.9 $-\frac{\sqrt{3}}{3}$;

15.10 $\frac{2}{\rho}$; 15.11 $\frac{\sqrt{2}}{2}$; 15.12 $a^b > \ln a$; 15.13 1; 15.14 $\frac{2}{25}$; 15.15 $-\frac{1}{x^2}$; 15.16 $-\ln 2$;

15.17 1; 15.18 $2 \ln 2$; 15.19 $2 \cos x$; 15.20 $\frac{1}{6\sqrt{2}}$; 15.21 0; 15.22 0; 15.23 $-\frac{1}{12\rho}$;

15.24 -3 ; 15.25 $\frac{1}{5 \ln 10}$; 15.26 $3 \ln 3$; 15.27 $-\frac{1}{16}$; 15.28 $\frac{b-a}{2a}$; 15.29 $\frac{3}{2}$; 15.30 $\frac{1}{3\rho \ln 3}$.

Funksiyaning limiti

16-masala. Funksiyaning limitini hisoblang .

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{e^{1+x^2} - 2^x}{1+x^2} \cdot \frac{1}{\sin^3 x} &= \lim_{x \rightarrow 0} \left(e^{\frac{\ln(1+x^2) \cdot 2^x}{1+x^2}} \right) \frac{1}{\sin^3 x} = \\ &= \lim_{x \rightarrow 0} e^{\frac{1}{\sin^3 x} \ln \frac{1+x^2 \cdot 2^x}{1+x^2}} = \exp \left[\lim_{x \rightarrow 0} \frac{1}{\sin^3 x} \ln \frac{1+x^2 \cdot 2^x}{1+x^2} \right] = \end{aligned}$$

$$\begin{aligned}
&= \exp \left\{ \lim_{x \rightarrow 0} \frac{1}{\sin^3 x} \ln \frac{1 + x^2 \times 5^x - x^2 \times 5^x + x^2 \times 2^x}{1 + x^2 \times 5^x} \right\} = \\
&= \exp \left\{ \lim_{x \rightarrow 0} \frac{1}{\sin^3 x} \ln \frac{1 - x^2 \times 5^x + x^2 \times 2^x}{1 + x^2 \times 5^x} \right\} = \\
&= \exp \left\{ \lim_{x \rightarrow 0} \frac{1}{\sin^3 x} \ln \frac{1 - x^2 \frac{5^x + 2^x}{1 + x^2 \times 5^x}}{1} \right\} = \\
&= \frac{e^{x \ln 5} - 1}{e^{x \ln 2} - 1} \gg x \ln 5 = \exp \left\{ \lim_{x \rightarrow 0} \frac{1}{\sin^3 x} \ln \frac{1 - x^2 \frac{5^x + 2^x}{1 + x^2 \times 5^x}}{1} \right\} = \\
&= \exp \left\{ \lim_{x \rightarrow 0} - \frac{x \ln 5}{x(1 + x^2 \times 5^x)} + \lim_{x \rightarrow 0} \frac{x \ln 2}{x(1 + x^2 \times 5^x)} \right\} = \\
&= \exp \left\{ \lim_{x \rightarrow 0} - \frac{\ln 5}{1 + x^2 \times 5^x} + \lim_{x \rightarrow 0} \frac{\ln 2}{1 + x^2 \times 5^x} \right\} = \\
&= \exp \left\{ - \frac{\ln 5}{1 + 0 \times 5^0} + \frac{\ln 2}{1 + 0 \times 5^0} \right\} = \exp \{- \ln 5 + \ln 2\} = \\
&= \exp \left\{ - \ln \frac{5}{2} \right\} = \exp \left\{ \ln \frac{2}{5} \right\} = e^{\ln \frac{2}{5}} = \frac{2}{5}.
\end{aligned}$$

$$1. \lim_{x \rightarrow 0} (1 - \ln(1 + x^3))^{\frac{3}{(x^2 \arcsin x)}}.$$

$$2. \lim_{x \rightarrow 0} (\cos \sqrt{x})^{\frac{1}{x}}.$$

$$3. \lim_{x \rightarrow 0} \frac{1 + x \times 2^x}{1 + x \times 3^x}.$$

$$4. \lim_{x \rightarrow 0} (2 - 3^{\arctg^2 \sqrt{x}})^{\frac{2}{\sin x}}.$$

$$5. \lim_{x \rightarrow 0} \frac{1 + \sin x \cos ax}{1 + \sin x \cos bx}.$$

$$6. \lim_{x \rightarrow 0} (5 - \frac{4}{\cos x})^{\frac{1}{\sin^2 3x}}.$$

$$7. \lim_{x \rightarrow 0} (1 - \ln(1 + \sqrt[3]{x}))^{\frac{x}{\sin^4 \sqrt[3]{x}}}.$$

$$8. \lim_{x \rightarrow 0} (2 - e^{\arcsin^2 \sqrt{x}})^{\frac{3}{x}}.$$

$$9. \lim_{x \rightarrow 0} (\cos px)^{\frac{1}{(x \sin px)}}.$$

$$10. \lim_{x \rightarrow 0} (1 + \sin^2 3x)^{\frac{1}{\ln \cos x}}.$$

11. $\lim_{x \rightarrow 0} (tg(\frac{\rho}{4} - x))^{ctgx}$.
12. $\lim_{x \rightarrow 0} (1 - x \sin^2 x)^{\frac{1}{\ln(1+\rho x^3)}}$.
13. $\lim_{x \rightarrow 0} (2 - 5^{\arcsin x^3})^{\frac{\cos ec^2 x}{x}}$.
14. $\lim_{x \rightarrow 0} (2 - \cos 3x)^{\frac{1}{\ln(1+x^2)}}$.
15. $\lim_{x \rightarrow 0} (2 - e^{\sin x})^{ctg \rho x}$.
16. $\lim_{x \rightarrow 0} (\cos x)^{\frac{1}{\ln(1+\sin^2 x)}}$.
17. $\lim_{x \rightarrow 0} (2 - e^{x^2})^{\frac{1}{\ln(1+tg^2(\rho \frac{x}{3}))}}$.
18. $\lim_{x \rightarrow 0} (3 - 2 \cos x)^{-\cos ec^2 x}$.
19. $\lim_{x \rightarrow 0} (2 - 3^{\sin^2 x})^{\frac{1}{\ln \cos x}}$.
20. $\lim_{x \rightarrow 0} x^2 \sqrt{2 - \cos x}$.
21. $\lim_{x \rightarrow 0} (6 - \frac{5}{\cos x})^{ctg^2 x}$.
22. $\lim_{x \rightarrow 0} (3 - \frac{2}{\cos x})^{\cos ec^2 x}$.
23. $\lim_{x \rightarrow 0} (\frac{1 + \sin x \cos 2x}{1 + \sin x \cos 3x})^{\frac{1}{\sin x^3}}$.
24. $\lim_{x \rightarrow 0} (2 - e^{x^2})^{\frac{1}{(1 - \cos \rho x)}}$.
25. $\lim_{x \rightarrow 0} (1 + \ln \frac{1}{3} \arctg^6 \sqrt{x})^{\frac{1}{x^3}}$.
26. $\lim_{x \rightarrow 0} \frac{1 + tg x \cos 2x}{1 + tg x \cos 5x} x^{\frac{1}{x^3}}$.
27. $\lim_{x \rightarrow 0} (\frac{1 + x3^x}{1 + x7^x})^{\frac{1}{tg^2 x}}$.
28. $\lim_{x \rightarrow 0} (1 + tg^2 x)^{\frac{1}{\ln(1+3x^2)}}$.
29. $\lim_{x \rightarrow 0} (1 - \ln \cos x)^{\frac{1}{tg^2 x}}$.
30. $\lim_{x \rightarrow 0} (1 - \sin^2 \frac{x}{2})^{\frac{1}{\ln(1+tg^2 3x)}}$.

Javoblar. 16.1 e^{-3} ; 16.2 $\frac{1}{\sqrt{e}}$; 16.3 $\frac{2}{3}$; 16.4 $\frac{1}{9}$; 16.5 $e^{\frac{b^2 - a^2}{2}}$; 16.6 $e^{-\frac{2}{9}}$; 16.7 e^{-1} ; 16.8 e^{-3} ;
 16.9 $e^{-\frac{\rho}{2}}$; 16.10 e^{-18} ; 16.11 e^{-2} ; 16.12 $e^{-\frac{1}{\rho}}$; 16.13 $\frac{1}{5}$; 16.14 $e^{\frac{9}{2}}$; 16.15 $e^{-\frac{1}{\rho}}$; 16.16 $e^{-\frac{1}{2}}$;
 16.17 $e^{-\frac{9}{\rho^2}}$; 16.18 e^{-1} ; 16.19 9; 16.20 \sqrt{e} ; 16.21 $e^{-\frac{5}{2}}$; 16.22 e^{-1} ; 16.23 $e^{\frac{5}{2}}$; 16.24 $e^{-\frac{2}{\rho^2}}$;
 16.25 $\frac{1}{3}$; 16.26 $e^{\frac{21}{2}}$; 16.27 $\frac{3}{7}$; 16.28 $e^{\frac{1}{3}}$; 16.29 $e^{\frac{1}{2}}$; 16.30 $e^{-\frac{1}{36}}$.

Funksiyaning limiti

17-masala. Funksiyaning limitini hisoblang .

$$\lim_{x \rightarrow 0} \frac{x^3 + 4\sqrt[3]{x+2}}{x^3 + 9\sqrt[3]{x}} = \lim_{x \rightarrow 0} \frac{0^3 + 4\sqrt[3]{0+2}}{0^3 + 9\sqrt[3]{0}} = \frac{4\sqrt[3]{2}}{9\sqrt[3]{0}} = \sqrt{\frac{4}{9}} = \frac{2}{3}.$$

$$1. \lim_{x \rightarrow 0} \frac{\sin 2x}{x}.$$

$$2. \lim_{x \rightarrow 0} \frac{2 + x}{3 - x}.$$

$$3. \lim_{x \rightarrow 0} \frac{\sin 4x}{x}.$$

$$4. \lim_{x \rightarrow 0} \frac{e^{3x} - 1}{x}.$$

$$5. \lim_{x \rightarrow 0} (\cos x)^{x+3}.$$

$$6. \lim_{x \rightarrow 0} \frac{x^2 + 4}{x + 2}.$$

$$7. \lim_{x \rightarrow 0} \frac{\ln(1+x)}{6x}.$$

$$8. \lim_{x \rightarrow 0} \frac{e^{4x} - 1}{x}.$$

$$9. \lim_{x \rightarrow 0} \frac{e^{x^3} - 1}{x^2}.$$

$$10. \lim_{x \rightarrow 0} \frac{x + 2}{e^{\cos x} + 4}.$$

$$11. \lim_{x \rightarrow 0} \frac{\sin 6x}{2x}.$$

$$12. \lim_{x \rightarrow 0} \frac{e^{x^2} - 1}{x^2}.$$

$$13. \lim_{x \rightarrow 0} \frac{\sin 2x}{\sin 3x}.$$

$$14. \lim_{x \rightarrow 0} \frac{\operatorname{ctg}(x + \frac{\pi}{3})}{3}.$$

$$15. \lim_{x \rightarrow 0} \frac{x^3 + 8}{3x^2 + 10}.$$

$$16. \lim_{x \rightarrow 0} (\sin(x+2))^{\frac{3}{3+x}}.$$

$$17. \lim_{x \rightarrow 0} \frac{e^{2x} - 1}{x}.$$

$$18. \lim_{x \rightarrow 0} \frac{x^4 + 5}{x + 10}.$$

$$19. \lim_{x \rightarrow 0} \frac{1 + 8\cos^2 x}{12x + 1}.$$

$$20. \lim_{x \rightarrow 0} \frac{x^3 + 1}{x^3 + 8}.$$

$$21. \lim_{x \rightarrow 0} \frac{\ln(1+x^2)}{x^2}.$$

$$22. \lim_{x \rightarrow 0} (\cos \frac{x}{p})^{1+x}.$$

$$23. \lim_{x \rightarrow 0} \frac{\arcsin x}{x}.$$

$$24. \lim_{x \rightarrow 0} \frac{\operatorname{arctg} 3x}{x}.$$

$$25. \lim_{x \rightarrow 0} (e^x + x)^{\cos x^4}.$$

$$26. \lim_{x \rightarrow 0} \frac{\sin 5x^2}{\sin x}.$$

$$27. \lim_{x \rightarrow 0} \operatorname{ctg}(\frac{\pi}{4} - x).$$

$$28. \lim_{x \rightarrow 0} \frac{5 \operatorname{tg}^2 x}{\cos x} = 0. \quad 29. \lim_{x \rightarrow 0} \frac{1+8x}{2+11x} = \frac{1}{2}. \quad 30. \lim_{x \rightarrow 0} \frac{\arcsin^2 x}{\arcsin^2 4x} = \frac{1}{16}.$$

Javoblar. 17.1 2; 17.2 1; 17.3 4; 17.4 $\sqrt{3}$; 17.5 1; 17.6 8; 17.7 1; 17.8 16; 17.9 0; 17.10 $\frac{1}{2}$;

17.11 9; 17.12 1; 17.13 1; 17.14 3; 17.15 0,64; 17.16 $\sin 2$; 17.17 $2 \ln 2$; 17.18 $\frac{1}{4}$; 17.19

8; 17.20 $\frac{1}{64}$; 17.21 1; 17.22 1; 17.23 1; 17.24 9; 17.25 1; 17.26 0; 17.27 1;

17.28 1; 17.29 $\frac{1}{2}$; 17.30 $\frac{1}{16}$.

Funksiyaning limiti

18-masala. Funksiyaning limitini hisoblang .

$$\begin{aligned} \lim_{x \rightarrow 1} \frac{2x-1}{x} \frac{\ln(3+2x)}{\ln(2-x)} &= \lim_{x \rightarrow 1} e^{\ln \frac{2x-1}{x} \frac{\ln(3+2x)}{\ln(2-x)}} = \lim_{x \rightarrow 1} e^{\frac{\ln(3+2x)}{\ln(2-x)} \ln \frac{2x-1}{x}} = \\ &= \exp \left[\lim_{x \rightarrow 1} \frac{\ln(3+2x)}{\ln(2-x)} \ln \frac{2x-1}{x} \right] \quad \begin{matrix} \text{é} \\ \text{é} \end{matrix} \begin{matrix} x = y+1 \\ x \rightarrow 1 \end{matrix} \quad \begin{matrix} \text{ü} \\ \text{ü} \end{matrix} \begin{matrix} y = x-1 \\ y \rightarrow 0 \end{matrix} = \\ &= \exp \left[\lim_{x \rightarrow 1} \frac{\ln(3+2(y+1))}{\ln(2-(y+1))} \ln \frac{2(y+1)-1}{y+1} \right] = \\ &= \exp \left[\lim_{x \rightarrow 1} \frac{\ln(5+2y)}{\ln(1-y)} \ln \frac{y+1}{y+1} \right] = \\ &= \exp \left[\lim_{x \rightarrow 1} \frac{\ln(5+2y)}{\ln(1-y)} \ln \left(1 + \frac{y}{y+1} \right) \right] = \\ &= \exp \left[\ln \left(1 + \frac{y}{y+1} \right) \right] = \frac{y}{y+1}, \text{ при } y \rightarrow 0 \quad \frac{y}{y+1} \rightarrow 0 \\ &\ln(1-y) \approx -y, \text{ при } y \rightarrow 0 \quad (-y \rightarrow 0) \end{aligned}$$

$$\begin{aligned}
 &= \exp \left\{ \lim_{x \rightarrow 1} \frac{\ln(5+2y)}{y} \right\} = \exp \left\{ \lim_{x \rightarrow 1} \frac{\ln(5+2y)}{y+1} \right\} \\
 &= \exp \left\{ \frac{\ln(5+2 \cdot 0)}{0+1} \right\} = \exp \{-\ln 5\} = \exp \left\{ \ln \frac{1}{5} \right\} = e^{\ln \frac{1}{5}} = \frac{1}{5}.
 \end{aligned}$$

$$1. \lim_{x \rightarrow 1} \frac{x^3 - 1}{x + 1}.$$

$$2. \lim_{x \rightarrow a} \frac{\sin x}{\sin a}.$$

$$3. \lim_{x \rightarrow 1} \frac{x^2 - 1}{x}.$$

$$4. \lim_{x \rightarrow 2} \frac{\cos x}{\cos 2}.$$

$$5. \lim_{x \rightarrow 8} \frac{x^2 - 7}{x + 1}.$$

$$6. \lim_{x \rightarrow p/4} (\operatorname{tg} x)^{1/(\cos(3p/4 - x))}.$$

$$7. \lim_{x \rightarrow 1} \frac{x^2 - 1}{x}.$$

$$8. \lim_{x \rightarrow a} (2 - x/a)^{\operatorname{tg} \frac{px}{2a}}.$$

$$9. \lim_{x \rightarrow 2p} (\cos x)^{\operatorname{ctg} 2x / \sin 3x}.$$

$$10. \lim_{x \rightarrow 2p} (\cos x)^{1/\sin^2 2x}.$$

$$11. \lim_{x \rightarrow 3} \frac{x^6 - x}{3}.$$

$$12. \lim_{x \rightarrow 4p} (\cos x)^{\operatorname{ctg} x / \sin 4x}.$$

$$13. \lim_{x \rightarrow 1} (3 - 2x)^{\operatorname{tg}(px/2)}.$$

$$14. \lim_{x \rightarrow 4p} (\cos x)^{\frac{5}{\operatorname{tg} 5x \sin 2x}}.$$

$$15. \lim_{x \rightarrow 3} \frac{x^9 - 2x}{3}.$$

$$16. \lim_{x \rightarrow p/2} (\sin x)^{6 \operatorname{tg} x \operatorname{tg} 3x}.$$

$$17. \lim_{x \rightarrow 1} (2e^{x-1} - 1)^{x/(x-1)}.$$

$$18. \lim_{x \rightarrow p/2} \operatorname{ctg} \frac{x}{2}.$$

$$19. \lim_{x \rightarrow 1} (2e^{x-1} - 1)^{\frac{3x-1}{x-1}}.$$

$$20. \lim_{x \rightarrow p/2} (1 + \cos 3x)^{\sec x}.$$

$$21. \lim_{x \rightarrow 2} (2e^{x-2} - 1)^{\frac{3x+2}{x-2}}.$$

$$22. \lim_{x \rightarrow 1} \frac{\sin(x-1)}{x-1}.$$

$$23. \lim_{x \rightarrow 1} \frac{e^{2-x} - e^2}{x - 1}.$$

$$24. \lim_{x \rightarrow p/2} \left(\operatorname{ctg} \frac{x}{2} \right)^{1/\cos x}.$$

$$25. \lim_{x \rightarrow 1} (2 - x)^{\frac{\sin(px/2)}{\ln(2-x)}}.$$

$$26. \lim_{x \rightarrow 3} \frac{\sin x}{\sin 3}.$$

$$27. \lim_{x \rightarrow 1} \frac{e^{x+1} - e^2}{2x - 2}.$$

$$28. \lim_{x \rightarrow p/2} (\sin x)^{\frac{18 \sin x}{\operatorname{ctg} x}}.$$

$$29. \lim_{x \rightarrow 1} \frac{e^{\ln(x+1)} - e^{\ln(2-x)}}{x - 1}.$$

$$30. \lim_{x \rightarrow p} \left(\operatorname{ctg} \frac{x}{4} \right)^{1/\cos(x/2)}.$$

Javoblar. 18.1 e^3 ; 18.2 $e^{\operatorname{ctg} a}$; 18.3 e^3 ; 18.4 $e^{-\operatorname{tg} 2}$; 18.5 $e^{\frac{4}{3}}$; 18.6 e^2 ; 18.7 e^5 ; 18.8 $e^{\frac{2}{p}}$; 18.9 $e^{-\frac{1}{12}}$; 18.10 $e^{-\frac{1}{8}}$; 18.11 $e^{\frac{2}{p}}$; 18.12 $e^{-\frac{1}{8}}$; 18.13 $e^{\frac{4}{p}}$; 18.14 $e^{-\frac{1}{4}}$; 18.15 $e^{\frac{4}{p}}$; 18.16 e^{-1} ; 18.17 e^2 ; 18.18 e ; 18.19 e^4 ; 18.20 e^{-3} ; 18.21 e^{16} ; 18.22 e^{-1} ; 18.23 e^2 ; 18.24 e ; 18.25 e ; 18.26 $e^{\operatorname{ctg} 3}$; 18.27 $\sqrt{3}$; 18.28 1 ; 18.29 2 ; 18.30 e .

Funksiyaning limiti

19–masala. Funksiyaning limitini hisoblang.

$$\begin{aligned} &= \lim_{x \rightarrow 1} \frac{e^{2x} - e^2}{x - 1} = \lim_{x \rightarrow 1} \frac{e^2(e^{2x-2} - 1)}{x - 1} = \\ &\lim_{x \rightarrow 1} \frac{e^{2x} - e^2}{x - 1} = \lim_{x \rightarrow 1} \frac{e^{2x} - e^2}{x - 1} \cdot \frac{\lim_{x \rightarrow 1} x+1}{\lim_{x \rightarrow 1} x+1} = \end{aligned}$$

Cheksiz kichiklarning ekvivalentlik munosabatidan foydalanib:

$$x \rightarrow 1 \quad (2x - 2 \rightarrow 0) \quad \text{da} \quad e^{2x-2} - 1 \sim 2x - 2 \quad \text{ega bo'lamiz.}$$

Natijada:

$$= \lim_{x \rightarrow 1} \frac{e^2(2x - 2)}{x - 1} = \lim_{x \rightarrow 1} 2e^2 = (2e^2)^1 = 4e^2.$$

$$1. \lim_{x \rightarrow e} \frac{\ln x - 1}{e^x - e} \cdot \frac{\sin \frac{p}{2e} x}{\frac{p}{2e}}.$$

$$3. \lim_{x \rightarrow p/4} \frac{\ln \operatorname{tg} x}{e^{1/(x+p/4)} - 1} \cdot \frac{\operatorname{ctg} x}{\frac{1}{x+p/4}}.$$

$$5. \lim_{x \rightarrow 2} \frac{\sin 3px}{e^{\sin^2(x-2)} - 1} \cdot \frac{\sin px}{\sin^2(x-2)}.$$

$$7. \lim_{x \rightarrow 3} (2 - \frac{x}{3})^{\sin px}.$$

$$9. \lim_{x \rightarrow 1} (1 + e^x)^{\frac{\sin px}{1-x}}.$$

$$11. \lim_{x \rightarrow 3} \frac{\arcsin(x-3)}{e^{\sin 3px} - 1} \cdot \frac{x^2 - 8}{\sin 3px}.$$

$$13. \lim_{x \rightarrow 1} \frac{\arctg \frac{x-3/4}{(x-1)^2}}{e^{x+1} - 1} \cdot \frac{x+1}{(x-1)^2}.$$

$$15. \lim_{x \rightarrow a} \frac{\sin x - \sin a}{e^{x^2/a^2} - 1} \cdot \frac{x^2/a^2}{x-a}.$$

$$17. \lim_{x \rightarrow p/4} (\sin x + \cos x)^{1/\operatorname{tg} x}.$$

$$19. \lim_{x \rightarrow 1} (\arcsin x)^{\operatorname{tg} px}.$$

$$21. \lim_{x \rightarrow 1} (\ln^2 ex)^{1/(x^2+1)}.$$

$$23. \lim_{x \rightarrow 1} \frac{x^3 - 1}{e^{1/x^2} - 1} \cdot \frac{1/x^2}{x-1}.$$

$$25. \lim_{x \rightarrow 2} (\cos px)^{\operatorname{tg}(x-2)}.$$

$$27. \lim_{x \rightarrow p/2} (\cos x + 1)^{\sin x}.$$

$$2. \lim_{x \rightarrow p/4} (\operatorname{tg} x)^{\operatorname{ctg} x}.$$

$$4. \lim_{x \rightarrow 2} (\sin x)^{3/(1+x)}.$$

$$6. \lim_{x \rightarrow p/6} (\sin x)^{6x/p}.$$

$$8. \lim_{x \rightarrow 1} \frac{1+x}{e^{(1-x^2)/(1-x)} - 1} \cdot \frac{(1-x^2)/(1-x)}{2+x}.$$

$$10. \lim_{x \rightarrow 1} \frac{\operatorname{tg} 9px}{e^{\sin 4px} - 1} \cdot \frac{x/(x+1)}{\sin 4px}.$$

$$12. \lim_{x \rightarrow p/4} (\sin 2x)^{\frac{x^2 - p^2/16}{x - p/4}}.$$

$$14. \lim_{x \rightarrow p} \frac{\operatorname{ctg} \frac{x}{4}}{e^{\sin(x-p)} - 1} \cdot \frac{\sin(x-p)}{x/4}.$$

$$16. \lim_{x \rightarrow 2} \frac{\sqrt{x+2} - 2}{e^{1/x} - 1} \cdot \frac{1/x}{x^2 - 4}.$$

$$18. \lim_{x \rightarrow p/8} (\operatorname{tg} 2x)^{\sin(p/8+x)}.$$

$$20. \lim_{x \rightarrow p} (x + \sin x)^{\sin x + x}.$$

$$22. \lim_{x \rightarrow 1} (\sqrt{x} + 1)^{p/\operatorname{arctg} x}.$$

$$24. \lim_{x \rightarrow 1} \frac{e^{\sin px} - 1}{e^{x^2+1} - 1} \cdot \frac{x^2+1}{x-1}.$$

$$26. \lim_{x \rightarrow 1/2} (\arcsin x + \arccos x)^{\frac{1}{x}}.$$

$$28. \lim_{x \rightarrow 1} (\sqrt[3]{x} + x - 1)^{\sin(px/4)}.$$

$$29. \lim_{x \rightarrow 1} \frac{x^2 + 2x - 3}{x^2 + 4x - 5}.$$

$$30. \lim_{x \rightarrow 1} \frac{1 + \cos px}{\operatorname{tg}^2 px}.$$

Javoblar. 19.1 $\frac{1}{e}$; 19.2 1; 19.3 1; 19.4 $\sin 2$; 19.5 1; 19.6 $\frac{1}{2}$; 19.7 1; 19.8 $\frac{4}{9}$; 19.9 $(1+e)^p$;

19.10 $\frac{3}{2}$; 19.11 $-\frac{1}{3p}$; 19.12 1; 19.13 $\frac{p^2}{4}$; 19.14 0; 19.15 $\cos a$; 19.16 $\frac{1}{4}$; 19.17 $\sqrt{2}$;

19.18 1; 19.19 1; 19.20 p^p ; 19.21 1; 19.22 16; 19.23 3; 19.24 p^2 ; 19.25 1; 19.26 $\frac{p^2}{4}$;

19.27 1; 19.28 1; 19.29 $\frac{2}{3}$; 19.30 $\frac{1}{2}$.

20–masala. Funksiyaning yoki sonli ketma-ketlikning limitini hisoblang.

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\sqrt{n^2 + 3n - 1} + \sqrt[3]{2n^2 + 1}}{n + 2 \sin n} &= \\ = \lim_{n \rightarrow \infty} \frac{\frac{1}{n}(\sqrt{n^2 + 3n - 1} + \sqrt[3]{2n^2 + 1})}{\frac{1}{n}(n + 2 \sin n)} &= \\ = \lim_{n \rightarrow \infty} \frac{\sqrt{1 + \frac{3}{n} - \frac{1}{n^2}} + \sqrt[3]{\frac{2}{n} + \frac{1}{n^3}}}{1 + 2 \frac{\sin n}{n}} &= \lim_{n \rightarrow \infty} \frac{\sin n}{n} = 0 \\ = \frac{\sqrt{1+0-0} + \sqrt[3]{0+0}}{1+2 \cdot 0} &= \frac{\sqrt{1} + \sqrt[3]{0}}{1} = 1. \end{aligned}$$

1. $\lim_{x \rightarrow 0} \sqrt{4 \cos 3x + x \operatorname{arctg} \frac{1}{x}}.$

2. $\lim_{x \rightarrow p/2} \sqrt{3 \sin x + (2x - p) \sin \frac{x}{2x - p}}.$

3. $\lim_{n \rightarrow \infty} \frac{2n - \sin n}{\sqrt{n} - \sqrt[3]{n^3 - 7}}.$

4. $\lim_{x \rightarrow 0} \frac{\operatorname{tg} x \cdot \cos \frac{1}{x} + \lg(2+x)}{\lg(4+x)}.$

$$5. \lim_{n \rightarrow \infty} \frac{e^{1/n} + \sin \frac{n}{n^2 + 1} \cos n}{1 + \cos \frac{1}{n}}.$$

$$6. \lim_{n \rightarrow \infty} \frac{\sqrt[4]{2 + n^5} - \sqrt{2n^3 + 3}}{(n + \sin n)\sqrt{7n}}.$$

$$7. \lim_{x \rightarrow p/4} \frac{\sqrt[3]{tgx} + (4x - p) \cos \frac{x}{4x - p}}{\lg(2 + tgx)}.$$

$$8. \lim_{n \rightarrow \infty} (\sin \sqrt{n^2 + 1} \operatorname{arctg} \frac{n}{n^2 + 1}).$$

$$9. \lim_{n \rightarrow \infty} \frac{n^2 - \sqrt{3n^5 - 7}}{(n^2 - n \cos n + 1)\sqrt{n}}.$$

$$10. \lim_{n \rightarrow \infty} \frac{3 \sin n + \sqrt{n - 1}}{n + \sqrt{n + 1}}.$$

$$11. \lim_{n \rightarrow \infty} \frac{(1 - \cos n)\sqrt[3]{n}}{\sqrt{2n + 1} - 1}.$$

$$12. \lim_{x \rightarrow 0} \ln \frac{2 + \sqrt{\arctg x \sin \frac{1}{x}}}{e}.$$

$$13. \lim_{x \rightarrow -2} \sqrt{\frac{1 + \cos px}{4 + (x + 2) \sin \frac{x}{x + 2}}}.$$

$$14. \lim_{n \rightarrow \infty} \frac{n}{\sqrt[3]{n^4 - 3} + \sin n}.$$

$$15. \lim_{n \rightarrow \infty} \frac{\sqrt[3]{n^2 \cos n} + \sqrt{3n^2 + 2}}{\sqrt[5]{n^6 + 1}}.$$

$$16. \lim_{x \rightarrow 0} \frac{\sqrt[3]{tgx} \operatorname{arctg} \frac{1}{x} + 3}{2 - \lg(1 + \sin x)}.$$

$$17. \lim_{x \rightarrow 0} \sqrt{\operatorname{arctg} \sin^2 \frac{1}{x} + 5 \cos x}.$$

$$18. \lim_{x \rightarrow 0} \sqrt{4 \cos x + \sin \frac{1}{x} \ln(1+x)}.$$

$$19. \lim_{x \rightarrow 0} \sqrt{2 \cos^2 x + (e^x - 1) \sin \frac{1}{x}}.$$

$$20. \lim_{x \rightarrow 0} \frac{2 + \ln(e + x \sin \frac{1}{x})}{\cos x + \sin x}.$$

$$21. \lim_{x \rightarrow 0} \ln \left((e^{x^2} - \cos x) \cos(\frac{1}{x}) + \operatorname{tg}(x + \frac{\rho}{3}) \right)$$

$$22. \lim_{x \rightarrow 0} \frac{\cos x + \ln(1+x) \sqrt{2 + \cos(1/x)}}{2 + e^x}.$$

$$23. \lim_{x \rightarrow 1} \frac{\cos 2\rho x}{2 + (e^{\sqrt{x-1}} - 1) \operatorname{arctg} \frac{x+2}{x-1}}.$$

$$24. \lim_{x \rightarrow 0} \sqrt{(e^{\sin x} - 1) \cos \frac{1}{x} + 4 \cos x}$$

$$25. \lim_{x \rightarrow 0} \frac{\cos(1+x)}{(2 + \sin \frac{1}{x}) \ln(1+x) + 2}.$$

26.

$$\lim_{x \rightarrow 2} \sqrt[3]{\lg(x+2) + \sin \sqrt{4-x^2} \cos \frac{x+2}{x-2}}.$$

$$27. \lim_{x \rightarrow \rho/2} \frac{2 + \cos x \sin \frac{2}{2x-\rho}}{3 + 2x \sin x}.$$

$$28. \lim_{x \rightarrow 1} \operatorname{tg}(\cos x + \sin \frac{x-1}{x+1} \cos \frac{x+1}{x-1}).$$

$$29. \lim_{x \rightarrow 0} \sqrt{x(2 + \sin \frac{1}{x}) + 4 \cos x}$$

$$30. \lim_{x \rightarrow 1} \frac{\sin x + \sin \rho x \times \arctg \frac{1+x}{1+\cos x}}{(2 + \sin \frac{1}{x}) \ln(1+x) + 2}.$$

Javoblar. 20.1 2; 20.2 $\sqrt{3}$; 20.3 - 2; 20.4 $\frac{1}{2}$; 20.5 $\frac{1}{2}$; 20.6 $-\frac{2}{7}$; 20.7 $\frac{1}{\lg(3)}$; 20.8 0;

20.9 $-\sqrt{3}$; 20.10 0; 20.11 0; 20.12 $\ln 2$; 20.13 $\frac{\sqrt{2}}{2}$; 20.14 1; 20.15 0; 20.16 $\frac{3}{2}$; 20.17 $\sqrt{5}$; 20.18

2; 20.19 $\sqrt{2}$; 20.20 3; 20.21 $\ln \sqrt{3}$; 20.22 $\frac{1}{3}$; 20.23 $\frac{1}{2}$; 20.24 2; 20.25 $\frac{\cos 1}{2}$;

20.26 $\sqrt[3]{\lg 4}$; 20.27 $\frac{2}{3+\rho}$; 20.28 $\operatorname{tg}(\cos 1)$; 20.29 2; 20.30 $\frac{\sin 1}{1+\cos 1}$.

III BOB. Funktsiyaning hosilasi

Bir argumentli funktsiyalarni differensiallash bo'limida siz funktsiyaning hosilasini topish, differensiallashning asosiy qoidalari, asosiy formulalari, yuqori tartibli hosilalarni hisoblash, shuningdek funktsiyalarni umumiy tekshirish masalalari bilan tanishasiz.

Ta'rif. Agar ushbu

$$\lim_{Dx \rightarrow 0} \frac{Dy}{Dx} = \lim_{Dx \rightarrow 0} \frac{f(x_0 + Dx) - f(x_0)}{Dx}$$

limit mavjud va chekli bo'lsa, u $f(x)$ funktsiyaning x_0 dagi hosilasi deyiladi va

$$f'(x_0) = \lim_{Dx \rightarrow 0} \frac{f(x_0 + Dx) - f(x_0)}{Dx} \quad (1)$$

kabi belgilanadi.

Agar $x_0 + Dx = x$ deyilsa, unda $Dx = x - x_0$ va $Dx \rightarrow 0$ da $x \rightarrow x_0$ bo'lib, (1) munosabat quyidagi

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$$

ko'rinishga keladi.

1-masala. Hosila ta'rifidan foydalanib, $f'(0)$ toping.

$$f(x) = \begin{cases} 1 - \cos\left(x \sin \frac{1}{x}\right), & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$x = 0$ nuqtada hosilasi

$$f'(0) = \lim_{\Delta x \rightarrow 0} \frac{f(0 + \Delta x) - f(0)}{\Delta x}$$

Hosila ta'rifidan foydalanib:

$$\begin{aligned} f'(0) &= \lim_{\Delta x \rightarrow 0} \frac{f(0 + \Delta x) - f(0)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{1 - \cos\left(\Delta x \sin \frac{1}{\Delta x}\right) - 0}{\Delta x} = \\ &= \lim_{\Delta x \rightarrow 0} \frac{1 - \left(1 - 2 \sin^2\left(\frac{1}{2} \Delta x \sin \frac{1}{\Delta x}\right)\right)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{2 \sin^2\left(\frac{1}{2} \Delta x \sin \frac{1}{\Delta x}\right)}{\Delta x} = \\ &= \left[\sin\left(\frac{1}{2} \Delta x \sin \frac{1}{\Delta x}\right) \gg \frac{1}{2} \Delta x \sin \frac{1}{\Delta x} \right] = \\ &= \lim_{\Delta x \rightarrow 0} \frac{2 \left(\frac{1}{2} \Delta x \sin \frac{1}{\Delta x}\right)^2}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{1}{2} \Delta x \sin^2 \frac{1}{\Delta x} = \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{2} \Delta x \sin^2 \frac{1}{\Delta x} - \text{chegaralangan, u holda} \lim_{\Delta x \rightarrow 0} \frac{1}{2} \Delta x \sin^2 \frac{1}{\Delta x} = 0. \end{aligned}$$

$$1. f(x) = \begin{cases} \operatorname{tg}\left(x^3 + x^2 \sin\left(\frac{x}{2}\right)\right), & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$2. f(x) = \begin{cases} \arcsin\left(x^2 \cos\left(\frac{1}{9x}\right)\right) + \frac{2}{3}x, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$3. f(x) = \begin{cases} \operatorname{arctg}\left(x \cos\left(\frac{1}{5x}\right)\right), & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$4. f(x) = \begin{cases} \ln\left(1 - \sin\left(x^3 \sin \frac{1}{x}\right)\right), & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$5. f(x) = \begin{cases} \sin\left(x \sin \frac{3}{x}\right), & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$6. f(x) = \begin{cases} \sqrt{1 + \ln\left(1 + x^2 \sin \frac{1}{x}\right)} - 1, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$7. f(x) = \begin{cases} \sin(e^{x^2 \sin \frac{5}{x}} - 1) + x, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$8. f(x) = \begin{cases} x^2 \cos\left(\frac{4}{3x}\right) + \frac{x^2}{2}, & x \neq 0; \\ 0, & x = 0 \end{cases}.$$

$$9. f(x) = \begin{cases} \operatorname{arctg}\left(x^3 - x^{\frac{3}{2}} \sin \frac{1}{3x}\right), & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$10. f(x) = \begin{cases} \sin x \times \cos \frac{5}{x}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$11. f(x) = \begin{cases} x + \arcsin\left(x^2 \sin \frac{6}{x}\right), & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$12. f(x) = \begin{cases} \operatorname{tg}\left(2^{x^2 \cos\left(\frac{1}{8x}\right)} - 1 + x\right), & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$13. f(x) = \begin{cases} \operatorname{arctg} x \times \sin \frac{7}{x}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$14. f(x) = \begin{cases} 2x^2 + x^2 \cos \frac{1}{9x}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$15. f(x) = \begin{cases} x^2 \cos^2 \frac{11}{x}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$16. f(x) = \begin{cases} 2x^2 + x^2 \cos \frac{1}{x}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$17. f(x) = \begin{cases} \frac{\ln(\cos x)}{x}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$18. f(x) = \begin{cases} 6x + x \sin \frac{1}{x}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$19. f(x) = \begin{cases} \frac{e^{x^2} - \cos x}{x}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$20. f(x) = \begin{cases} e^{x \sin \frac{5}{x}} - 1, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$21. f(x) = \begin{cases} 3^{x^2 \sin \frac{2}{x}} - 1 + 2x, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$22. f(x) = \begin{cases} \sqrt{1 + \ln(1 + 3x^2 \cos \frac{2}{x})} - 1, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$23. f(x) = \begin{cases} e^{x \sin \frac{3}{5x}} - 1, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$24. f(x) = \begin{cases} \frac{e^{\operatorname{tg} x} - e^{\sin x}}{x^2}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$25. f(x) = \begin{cases} \operatorname{arctg}(\frac{3x}{2} - x^2 \sin \frac{1}{x}), & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$26. f(x) = \begin{cases} e^{\sin \frac{3}{x^2} \sin \frac{2}{x}} - 1 + x^2, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$27. f(x) = \begin{cases} \sqrt[3]{1 - 2x^3 \sin \frac{5}{x}} - 1 + x, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$28. f(x) = \begin{cases} x^2 e^{|x|} \sin \frac{1}{x^2}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$29. f(x) = \begin{cases} \frac{\ln(1+2x^2+x^3)}{x}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

$$30. f(x) = \begin{cases} \frac{\cos x - \cos 3x}{x}, & x \neq 0; \\ 0, & x = 0 \end{cases}$$

Javoblar. 1.1 0; 1.2 $\frac{2}{3}$; 1.3 mavjud emas; 1.4 0; 1.5 mavjud emas; 1.6 0; 1.7 1; 1.8 0;

1.9 0; 1.10 mavjud emas; 1.11 1; 1.12 1; 1.13 mavjud emas; 1.14 0; 1.15 0; 1.16 0;

1.17 $-\frac{1}{2}$; 1.18 mavjud emas; 1.19 1,5; 1.20 mavjud emas; 1.21 $-\frac{1}{2}$; 1.22 0; 1.23 mavjud

emas; 1.24 $\ln \sqrt{2}$; 1.25 $\frac{3}{2}$; 1.26 0; 1.27 1; 1.28 0; 1.29 2; 1.30 4.

Differensiallanuvchi $y = f(x)$ funksiya grafigining $M_0(x_0, y_0)$ ($y_0 = f(x_0)$) nuqtasida o'tkazilgan urinma tenglamasi

$$y - y_0 = f'(x_0)(x - x_0)$$

ko'rinishga,

$f'(x_0) \neq 0$ da normal tenglamasi

$$y - y_0 = -\frac{1}{f'(x_0)}(x - x_0)$$

ko'rinishga ega bo'ladi.

2-masala. Funksiya grafigining abssissasi x_0 bo'lgan nuqtasiga o'tkazilgan urinma tenglamasini tuzing.

$$y = \sqrt[3]{x} - \frac{16\sqrt[4]{x}}{3}, \quad x_0 = 1.$$

Echim:

$$\begin{aligned}
 y &= 6\sqrt[3]{x} - \frac{16\sqrt[4]{x}}{3} = 6x^{\frac{1}{3}} - \frac{16x^{\frac{1}{4}}}{3} = \\
 &= 6 \times \frac{1}{3} \times x^{-\frac{2}{3}} - \frac{16}{3} \times \frac{1}{4} \times x^{-\frac{3}{4}} = 2x^{-\frac{2}{3}} - \frac{4}{3}x^{-\frac{3}{4}}. \\
 y &= y(x_0) = 2x^{-\frac{2}{3}} - \frac{4}{3}x^{-\frac{3}{4}} = 2 - \frac{4}{3} = \frac{2}{3}.
 \end{aligned}$$

y funksiya x_0 nuqtada hosilaga ega ekanligidan, urinma tenglamasi quyidagi ko'rinishda bo'ladi:

$$y - y_0 = y'(x - x_0), \text{ bu yerda } y' = \frac{2}{3}$$

$$y_0 = y(x_0) = 6\sqrt[3]{1} - \frac{16\sqrt[4]{1}}{3} = 6 - \frac{16}{3} = \frac{2}{3}$$

U holda:

$$y - \frac{2}{3} = \frac{2}{3}(x - 1)$$

$$y = \frac{2}{3}x - \frac{2}{3} + \frac{2}{3}$$

$$y = \frac{2}{3}x$$

Shunday qilib, urinma tenglamasi:

$$y = \frac{2}{3}x.$$

Funksiya grafigining abssissasi x_0 bo'lgan nuqtasiga o'tkazilgan normal (1–12 variantlarda) yoki urinma (13–30 variantlarda) tenglamasini tuzing.

$$1. y = \frac{4x - x^2}{4}, \quad x_0 = 2.$$

$$2. y = 2x^2 + 3x - 1, \quad x_0 = -2.$$

$$3. y = x^2 + 8\sqrt{x} - 32, \quad x_0 = 4.$$

$$4. y = x + \sqrt{x^3}, \quad x_0 = 1.$$

$$5. y = \sqrt[3]{x^2} - 20, x_0 = -8.$$

$$7. y = 8\sqrt[4]{x} - 70, x_0 = 16.$$

$$9. y = \frac{x^2 - 3x + 6}{x^2}, x_0 = 3.$$

$$11. y = \frac{x^3 + 2}{x^3 - 2}, x_0 = 2.$$

$$13. y = \frac{x^{29} + 6}{x^4 + 1}, x_0 = 1.$$

$$15. y = \frac{-2(x^8 + 2)}{3(x^4 + 1)}, x_0 = 1.$$

$$17. y = \frac{x^{16} + 9}{1 - 5x^2}, x_0 = 1.$$

$$19. y = \frac{1}{3x + 2}, x_0 = 2.$$

$$21. y = \frac{x^2 - 3x + 3}{x}, x_0 = 3.$$

$$23. y = -2(\sqrt[3]{x} + 3\sqrt{x}), x_0 = 1.$$

$$25. y = 14\sqrt{x} - 15\sqrt[3]{x} + 2, x_0 = 1.$$

$$27. y = \frac{3x - 2x^3}{3}, x_0 = 1.$$

$$29. y = \frac{x^2 - 2x - 3}{4}, x_0 = 4.$$

$$6. y = \frac{1 + \sqrt{x}}{1 - \sqrt{x}}, x_0 = 4.$$

$$8. y = 2x^2 - 3x + 1, x_0 = 1.$$

$$10. y = \sqrt{x} - 3\sqrt[3]{x}, x_0 = 64.$$

$$12. y = 2x^2 + 3, x_0 = -1.$$

$$14. y = 2x + \frac{1}{x}, x_0 = 1.$$

$$16. y = \frac{x^5 + 1}{x^4 + 1}, x_0 = 1.$$

$$18. y = 3(\sqrt[3]{x} - 2\sqrt{x}), x_0 = 1.$$

$$20. y = \frac{x}{x^2 + 1}, x_0 = -2.$$

$$22. y = \frac{2x}{x^2 + 1}, x_0 = 1.$$

$$24. y = \frac{1 + 3x^2}{3 + x^2}, x_0 = 1.$$

$$26. y = 3\sqrt[4]{x} - \sqrt{x}, x_0 = 1.$$

$$28. y = \frac{x^2}{10} + 3, x_0 = 2.$$

$$30. y = x - x^3, x_0 = -1.$$

Javoblar. 2.1 $x = 2$; 2.2 $y = \frac{x}{5} + 1\frac{2}{5}$; 2.3 $y = -\frac{x}{10} + \frac{2}{5}$; 2.4 $y = -\frac{2x}{5} + 2\frac{2}{5}$;

2.5 $y = 3x + 8$; 2.6 $y = -2x + 5$; 2.7 $y = -4x + 10$; 2.8 $y = -x + 1$; 2.9 $y = 9x - 26\frac{1}{3}$; 2.10

$$x = 64; \text{2.11 } y = \frac{3}{4}x + \frac{1}{6}; \text{2.12 } y = -4x + 1; \text{2.13 } y = 7,5x - 4; \text{2.14 } y = x + 2;$$

$$\text{2.15 } y = -\frac{2}{3}x - \frac{1}{3}; \text{2.16 } y = \frac{x}{2} + \frac{1}{2}; \text{2.17 } y = \frac{9}{4}x - \frac{19}{4}; \text{2.18 } y = -2x - 1;$$

$$\text{2.19 } y = -\frac{3}{64}x + \frac{7}{32}; \text{2.20 } y = -\frac{3}{25}x - \frac{16}{25}; \text{2.21 } y = x - 2; \text{2.22 } y = 1;$$

$$\text{2.23 } 3y + 11x + 13 = 0; \text{2.24 } y = x; \text{2.25 } y = 2x + 1; \text{2.26 } y = \frac{x}{4} + \frac{7}{4}; \text{2.27 } y = -x + 1\frac{1}{3}; \text{2.28 }$$

$$y = \frac{2}{5}x + \frac{13}{5}; \text{2.29 } y = \frac{3}{2}x - \frac{19}{4}; \text{2.30 } y = \frac{x}{2} + \frac{1}{2}.$$

Funksiyaning differentsiali

Agar $y = f(x)$ funksiyaning Dy orttirmasi $Dy = A \Delta x + o(\Delta x)$ ko'rinishda yozilishi mumkin bo'lsa, ortirmaning Δx ga nisbatan chiziqli qismi $A \Delta x$ funksiyaning differentsiali deyiladi va dy yoki $df(x)$ orqali belgilanadi: $dy = A(x)\Delta x$. Differentsila mavjud bo'lishi uchun chekli hosila $f'(x)$ ning mavjudligi va yetarlidir. Bunda $dy = f'(x)dx = y' dx$ bo'ladi.

Agar $y = f(u)$, $u = j(x)$ murakkab funksiya berilgan bo'lsa, u holda $dy = f'(u)du$ bo'ladi.

3-masala. Differentsial dy ni toping.

$$y = x\sqrt{x^2 - 1} + \ln|x + \sqrt{x^2 - 1}|.$$

$$\begin{aligned} dy &= y' dx = \left(x\sqrt{x^2 - 1} + \ln|x + \sqrt{x^2 - 1}| \right)' dx = \\ &= \left(\frac{x}{\sqrt{x^2 - 1}} + x \times \frac{1}{2\sqrt{x^2 - 1}} \times 2x + \frac{1}{x + \sqrt{x^2 - 1}} \times \frac{x}{\sqrt{x^2 - 1}} + \frac{1}{2\sqrt{x^2 - 1}} \times 2x \right) dx = \\ &= \left(\frac{x}{\sqrt{x^2 - 1}} + \frac{x^2}{\sqrt{x^2 - 1}} + \frac{x - \sqrt{x^2 - 1}}{x^2 - (x^2 - 1)} \times \frac{x}{\sqrt{x^2 - 1}} + \frac{x}{\sqrt{x^2 - 1}} \right) dx = \\ &= \left(\frac{x}{\sqrt{x^2 - 1}} + \frac{x^2}{\sqrt{x^2 - 1}} + x - \sqrt{x^2 - 1} + \frac{x}{\sqrt{x^2 - 1}} \right) dx = \\ &= \left(\frac{x}{\sqrt{x^2 - 1}} + \frac{x^2}{\sqrt{x^2 - 1}} + x - \sqrt{x^2 - 1} + \frac{x^2}{\sqrt{x^2 - 1}} - x \right) dx = \frac{2x^2 dx}{\sqrt{x^2 - 1}} \end{aligned}$$

$$1. y = x \arcsin\left(\frac{1}{x}\right) + \ln|x + \sqrt{x^2 - 1}|, \quad x > 0.$$

$$2. y = \operatorname{tg}\left(2 \arccos \sqrt{1 - 2x^2}\right), \quad x > 0.$$

$$3. y = \sqrt{1 + 2x} - \ln|x + \sqrt{1 + 2x}|.$$

$$4. y = x^2 \operatorname{arctg}\left(\sqrt{x^2 - 1}\right) - \sqrt{x^2 - 1}.$$

$$5. y = \arccos \frac{1}{\sqrt{1 + 2x^2}}, \quad x > 0.$$

$$6. y = x \ln|x + \sqrt{x^2 + 3}| - \sqrt{x^2 + 3}.$$

$$7. y = \operatorname{arctg}(\operatorname{sh} x) + (\operatorname{sh} x) \ln(\operatorname{ch} x).$$

$$8. y = \arccos \frac{x^2 - 1}{x^2 \sqrt{2}}.$$

$$9. y = \ln\left(\cos^2 x + \sqrt{1 + \cos^4 x}\right).$$

$$10. y = \ln\left(x + \sqrt{1 + x^2}\right) - \sqrt{1 + x^2} \operatorname{arctg} x.$$

$$11. y = \frac{\ln|x|}{1 + x^2} - \frac{1}{2} \ln \frac{x^2}{1 + x^2}.$$

$$12. y = \ln\left(e^x + \sqrt{e^{2x} - 1}\right) + \arcsin e^x.$$

$$13. y = x\sqrt{4 - x^2} + a \arcsin \frac{x}{2}$$

$$14. y = \ln\left(\operatorname{tg} \frac{x}{2}\right) - \frac{x}{\sin x}.$$

$$15. y = 2x + \ln|\sin x + 2 \cos x|.$$

$$16. y = \sqrt{\operatorname{ctg} x} - \frac{\sqrt{\operatorname{tg}^3 x}}{3}.$$

$$17. y = \ln \left| \frac{x + \sqrt{x^2 + 1}}{2x} \right|.$$

$$18. y = \sqrt[3]{\frac{x+2}{x-2}}.$$

$$19. y = \operatorname{arctg} \frac{x^2 - 1}{x}.$$

$$20. y = \ln|x^2 - 1| - \frac{1}{x^2 - 1}.$$

$$21. y = \operatorname{arctg}\left(\operatorname{tg} \frac{x}{2} + 1\right).$$

$$22. y = \ln|2x + 2\sqrt{x^2 + x + 1}|.$$

$$23. y = \ln|\cos \sqrt{x}| + \sqrt{x} \operatorname{tg} \sqrt{x}.$$

$$24. y = e^x (\cos 2x + 2 \sin 2x).$$

$$25. y = x(\sin(\ln x) - \cos(\ln x))$$

$$26. y = \frac{1}{2} \sqrt{x-1} - \frac{1}{2} e^{2\sqrt{x-1}}.$$

$$27. y = \cos x \ln(\operatorname{tg} x) - \ln\left(\operatorname{tg} \frac{x}{2}\right).$$

$$28. y = \sqrt{3+x^2} - x \ln|x + \sqrt{3+x^2}|.$$

$$29. y = \sqrt{x} - (1+x) \operatorname{arctg} \sqrt{x}.$$

$$30. y = x \operatorname{arctg} x - \ln \sqrt{1+x^2}.$$

Javoblar. 3.1 $\arcsin \frac{x}{\sqrt{1+x^2}} dx$; 3.2 $\frac{2\sqrt{2}}{(1-4x^2)^2 \sqrt{1-2x^2}} dx$; 3.3 $\frac{x-1}{(x+\sqrt{1+2x})\sqrt{1+2x}} dx$; 3.4

$2x \operatorname{arctg}(\sqrt{x^2-1}) dx$; 3.5 $\frac{\sqrt{2}x}{1+2x^2} dx$; 3.6 $\ln|x + \sqrt{x^2+3}| dx$; 3.7

$\operatorname{ch} x (1 + \ln(\operatorname{ch} x)) dx$; 3.8 $-\frac{2}{x\sqrt{x^4+2x^2-1}} dx$; 3.9 $-\frac{\sin 2x}{\sqrt{1+\cos^4 x}} dx$; 3.10

$-\frac{x}{\sqrt{1+x^2}} \operatorname{arctg} x dx$; 3.11 $-\frac{2x \ln|x|}{(1+x^2)^2} dx$; 3.12 $\frac{e^x}{\sqrt{e^{2x}-1}} + \frac{e^x}{\sqrt{1-e^{2x}}} dx$; 3.13

$-\frac{2x^2+4+2a}{\sqrt{4-x^2}} dx$; 3.14 $\frac{2\operatorname{ctg} x}{\sin x} dx$; 3.15 $\frac{5\cos x}{\sin x + 2\cos x} dx$; 3.16

$-\frac{\sqrt{2}}{\cos x \sqrt{\sin^3 2x}} dx$; 3.17 $-\frac{dx}{x(x+\sqrt{x^2+1})}$; 3.18 $\frac{4dx}{3(x-2)\sqrt{(x+2)^2(x-2)}}$;

3.19 $\frac{x^2+1}{x^4-x^2+1} dx$; 3.20 $\frac{2x^3}{(x^2-1)^2} dx$; 3.21 $\frac{dx}{3+2\sin x + \cos x}$; 3.22 $\frac{dx}{\sqrt{x^2+x}}$;

3.23 $\frac{dx}{2\cos^2 \sqrt{x}}$; 3.24 $5e^x \cos 2x dx$; 3.25 $2\sin(\ln x) dx$; 3.26 $e^{2\sqrt{x-1}} dx$;

3.27 $-\sin x \ln \operatorname{tg} x dx$; 3.28 $-\ln|x + \sqrt{3+x^2}| dx$; 3.29 $-\operatorname{arctg} \sqrt{x} dx$; 3.30 $\operatorname{arctg} x dx$.

Funksiyaning differentsiali

Agar x argumentning ortirmasi $Dx = x - x_0$ absolyut qiymati kichik bo'lsa, u holda

$$f(x) = f(x_0 + Dx) \approx f(x_0) + f'(x_0) \cdot Dx$$

ko'rinishda yoziladi. Bu formuladan funksiyalarning qiymatlarini taqribiy hisoblashlarda foydalaniladi.

4–masala. Differensial yordamida taqribiy hisoblang.

$$y = \frac{1}{\sqrt{2x+1}}, \quad x = 1,58.$$

Echim:

Agar x argumentning ortirmasi $Dx = x - x_0$ absolyut qiymati kichik bo'lsa, u holda

$$f(x) = f(x_0 + Dx) \approx f(x_0) + f'(x_0) \cdot Dx$$

$x_0 = 1,5$ ni deb olamiz

U holda:

$$Dx = 0,08$$

Hisoblaymiz:

$$y(1,5) = \frac{1}{\sqrt{2 \cdot 1,5 + 1}} = \frac{1}{\sqrt{4}} = \frac{1}{2}.$$

$$y'(x) = \frac{d}{dx} \frac{1}{\sqrt{2x+1}} = \left((2x+1)^{-\frac{1}{2}} \right)' = -\frac{1}{2} \cdot (2x+1)^{-\frac{3}{2}} \cdot 2 = -\frac{1}{\sqrt{(2x+1)^3}}.$$

$$y'(1,5) = -\frac{1}{\sqrt{(2 \cdot 1,5 + 1)^3}} = -\frac{1}{\sqrt{4^3}} = -\frac{1}{8}.$$

Natijada:

$$y(1,58) \approx y(1,5) + y'(1,5) \times 0,08 = \frac{1}{2} - \frac{1}{8} \times 0,08 = 0,5 - 0,01 = 0,49.$$

$$1. y = \sqrt[3]{x}, x = 7,76.$$

$$2. y = \sqrt[3]{x^3 + 7x}, x = 1,012.$$

$$3. y = \frac{x + \sqrt{5 - x^2}}{2}, x = 0,98.$$

$$4. y = \sqrt[3]{x}, x = 27,54.$$

$$5. y = \arcsin x, x = 0,08.$$

$$6. y = \sqrt[3]{x^2 + 7x + 5}, x = 0,97.$$

$$7. y = \sqrt[3]{x}, x = 26,46.$$

$$8. y = \sqrt[3]{x^2 + x + 3}, x = 1,97.$$

$$9. y = x^{11}, x = 1,021.$$

$$10. y = \sqrt[3]{x}, x = 1,21.$$

$$11. y = x^{21}, x = 0,998.$$

$$12. y = \sqrt[3]{x^2}, x = 1,03.$$

$$13. y = x^6, x = 2,01.$$

$$14. y = \sqrt[3]{x}, x = 8,24.$$

$$15. y = x^7, x = 1,996.$$

$$16. y = \sqrt[3]{x}, x = 7,64.$$

$$17. y = \sqrt{4x - 1}, x = 2,56.$$

$$18. y = \frac{1}{\sqrt{2x^2 + x + 1}}, x = 1,016.$$

$$19. y = \sqrt[3]{x}, x = 8,36.$$

$$20. y = \frac{1}{\sqrt{x}}, x = 4,16.$$

$$21. y = x^7, x = 2,002.$$

$$22. y = \sqrt{4x - 3}, x = 1,78.$$

$$23. y = \sqrt{x^3}, x = 0,98.$$

$$24. y = x^5, x = 2,997.$$

$$25. y = \sqrt[5]{x^2}, x = 1,03.$$

$$26. y = x^4, x = 3,998.$$

$$27. y = \sqrt{1 + x + \sin x}, x = 0,01.$$

$$28. y = \sqrt[3]{3x + \cos x}, x = 0,01.$$

$$29. y = \sqrt[4]{2x - \sin \frac{\pi x}{2}}, x = 1,02.$$

$$30. y = \sqrt{x^2 + 5}, x = 1,97.$$

Javoblar. 4.1 1,98; 4.2 2,01; 4.3 1,495; 4.4 3,02; 4.5 0,08; 4.6 1,99; 4.7 2,98; 4.8 2,975;
 4.9 1,231; 4.10 1,07; 4.11 0,958; 4.12 1,02; 4.13 65,92; 4.14 2,02; 4.15 126,208;
 4.16 1,97; 4.17 3,04; 4.18 0,495; 4.19 2,03; 4.20 0,49; 4.21 128,896; 4.22 2,03;
 4.23 0,98; 4.24 241,785; 4.25 1,012; 4.26 255,488; 4.27 1,01; 4.28 1,01; 4.29 1,01;

Differensiallashning asosiy qoidalari

Funksiya hosilasini topish amaliga funksiyani differensiallash deyiladi. Differensiallashning asosiy qoidalari:

Agar $u(x)$, $v(x)$ differensiallanuvchi funksiyalar bo'lib, C - *const*, ya'ni o'zgarmas bo'lsin.

$$1. [u(x) \pm v(x)]' = u'(x) \pm v'(x).$$

$$2. [Cu(x)]' = Cu'(x).$$

$$3. [u(x) \cdot v(x)]' = u'(x) \cdot v(x) + u(x) \cdot v'(x).$$

$$4. \frac{u'(x) \cdot v(x) - u(x) \cdot v'(x)}{v^2(x)} = \left(\frac{u(x)}{v(x)} \right)', \text{ bu yerda } v(x) \neq 0.$$

5-masala. Funksiyaning hosilasini toping.

$$\begin{aligned} y' &= \frac{3x^6 + 4x^4 - x^2 - 2}{15\sqrt{1+x^2}} \\ &= \frac{(3x^6 + 4x^4 - x^2 - 2)' \sqrt{1+x^2} - (3x^6 + 4x^4 - x^2 - 2) \left(\sqrt{1+x^2} \right)'}{15(1+x^2)} \\ &= \frac{(18x^5 + 16x^3 - 2x) \sqrt{1+x^2} - (3x^6 + 4x^4 - x^2 - 2) \cdot \frac{1}{\sqrt{1+x^2}} \cdot 2x}{15(1+x^2)} \end{aligned}$$

$$\begin{aligned}
&= \frac{(18x^5 + 16x^3 - 2x)\sqrt{1+x^2} - (3x^6 + 4x^4 - x^2 - 2) \times \frac{1}{\sqrt{1+x^2}} \times 2x}{15(1+x^2)} = \\
&= \frac{x(18x^4 + 16x^2 - 2)(1+x^2) - (3x^6 + 4x^4 - x^2 - 2) \times x}{15\sqrt{1+x^2}(1+x^2)} = \\
&= \frac{x(18x^4 + 16x^2 - 2 + 18x^6 + 16x^4 - 2x^2 - 3x^6 - 4x^4 + x^2 + 2)}{15\sqrt{1+x^2}(1+x^2)} = \\
&= \frac{x(15x^6 + 30x^4 + 15x^2)}{15\sqrt{1+x^2}(1+x^2)} = \frac{15x^3(x^4 + 2x^2 + 1)}{15\sqrt{1+x^2}(1+x^2)} = \frac{x^3(x^4 + 2x^2 + 1)}{\sqrt{1+x^2}(1+x^2)}.
\end{aligned}$$

$$1. y = \frac{2(3x^3 + 4x^2 - x - 2)}{15\sqrt{1+x}}.$$

$$2. y = \frac{(2x^2 - 1)\sqrt{1+x^2}}{3x^3}.$$

$$3. y = \frac{x^4 - 8x^2}{2(x^2 - 4)}.$$

$$4. y = \frac{2x^2 - x - 1}{3\sqrt{2+4x}}.$$

$$5. y = \frac{(1+x^8)\sqrt{1+x^8}}{12x^{12}}.$$

$$6. y = \frac{x^2}{2\sqrt{1-3x^4}}.$$

$$7. y = \frac{(x^2 - 6)\sqrt{(4+x^2)^3}}{120x^5}.$$

$$8. y = \frac{(x^2 - 8)\sqrt{x^2 - 8}}{6x^3}.$$

$$9. y = \frac{4 + 3x^3}{x \times \sqrt[3]{(2+x^3)^2}}.$$

$$10. y = \sqrt[3]{\frac{(1+x^{3/4})^2}{x^{3/2}}}.$$

$$11. y = \frac{x^6 + x^3 - 2}{\sqrt{1-x^3}}.$$

$$12. y = \frac{(x^2 - 2)\sqrt{4+x^2}}{24x^3}.$$

$$13. y = \frac{1+x^2}{2\sqrt{1+2x^2}}.$$

$$14. y = \frac{\sqrt{x-1} \times (3x+2)}{4x^2}.$$

$$15. y = \frac{\sqrt{(1+x^2)^3}}{3x^3}.$$

$$16. y = \frac{x^6 + 8x^3 - 128}{\sqrt{8-x^3}}.$$

$$17. y = \frac{\sqrt{2x+3} \times (x-2)}{x^2}.$$

$$18. y = (1-x^2)^5 \sqrt{x^3 + \frac{1}{x}}.$$

$$19. y = \frac{(2x^2 + 3)\sqrt{x^2 - 3}}{9x^3}.$$

$$20. y = \frac{x - 1}{(x^2 + 5)\sqrt{x^2 + 5}}.$$

$$21. y = \frac{(2x + 1)\sqrt{x^2 - x}}{x^2}.$$

$$22. y = 2 \times \sqrt{\frac{1 - \sqrt{x}}{1 + \sqrt{x}}}.$$

$$23. y = \frac{1}{(x + 2)\sqrt{x^2 + 4x + 5}}.$$

$$24. y = 3 \times \frac{\sqrt[3]{x^2 + x + 1}}{x + 1}.$$

$$25. y = 3 \times \sqrt[3]{\frac{x + 1}{(x - 1)^2}}.$$

$$26. y = \frac{x + 7}{6\sqrt{x^2 + 2x + 7}}.$$

$$27. y = \frac{x\sqrt{x + 1}}{x^2 + x + 1}.$$

$$28. y = \frac{x^2 + 2}{2\sqrt{1 - x^4}}.$$

$$29. y = \frac{(x + 3)\sqrt{2x - 1}}{2x + 7}.$$

$$30. y = \frac{3x + \sqrt{x}}{\sqrt{x^2 + 2}}.$$

Javoblar. 5.1 $x\sqrt{1+x}$; 5.2 $\frac{1}{x^4\sqrt{1+x^2}}$; 5.3 $x + \frac{16}{(x^2 - 4)^2}$; 5.4 $\frac{x}{\sqrt{2+4x}}$; 5.5 $-\frac{\sqrt{1+x^8}}{x^{13}}$;

5.6 $\frac{x}{(1 - 3x^4)\sqrt{1 - 3x^4}}$; 5.7 $\frac{\sqrt{4+x^2}}{x^6}$; 5.8 $\frac{4\sqrt{x^2 - 8}}{x^4}$; 5.9 $\frac{-8}{x^2(2+x^3)\sqrt{(2+x^3)^2}}$;

5.10 $-\frac{1}{2x \times \sqrt{x} \times \sqrt[3]{1+x^{3/4}}}$; 5.11 $\frac{9x^5}{2\sqrt{1-x^3}}$; 5.12 $\frac{-x^4 + x^3 + 2x^2 - 2x + 24}{24x^4\sqrt{4+x^2}}$;

5.13 $\frac{x^3}{(1+2x^2)\sqrt{1+2x^2}}$; 5.14 $\frac{-3x^2+8}{8x^3\sqrt{x-1}}$; 5.15 $-\frac{\sqrt{1+x^2}}{x^2}$; 5.16 $\frac{9x^5}{2\sqrt{8-x^3}}$;

5.17 $\frac{-x^2+3x+12}{x^3\sqrt{2x+3}}$; 5.18 $\frac{1}{5\sqrt{(x^3+\frac{1}{x})^4}} \times \frac{1}{x^2}$; 5.19 $\frac{3}{x^4\sqrt{x^2-3}}$;

5.20 $\frac{-2x^2+3x+5}{\sqrt{(x^2+5)^5}}$; 5.21 $\frac{3}{x^2\sqrt{x^2-3}}$; 5.22 $-\frac{1}{\sqrt{x(1-x)} \times (1+\sqrt{x})}$;

5.23 $-\frac{2x^2+8x+9}{(x+2)^2 \times \sqrt{(x^2+4x+5)^3}}$; 5.24 $-\frac{x^2+2}{\sqrt[3]{(x^2+x+1)^2} \times (x+1)^2}$;

$$\begin{aligned}
 5.25 &= \sqrt[3]{\frac{x-1}{(x+1)^2}} \times \frac{x+3}{(x-1)^2}; \quad 5.26 = \frac{x}{(x^2+2x+7) \times \sqrt{x^2+2x+7}}; \\
 5.27 &= \frac{-x^3 - x^2 + 3x + 2}{2\sqrt{x+1} \times (x^2+x+1)^2}; \quad 5.28 = \frac{2x^3+x}{(1-x^4) \times \sqrt{1-x^4}}; \quad 5.29 = \frac{2x^2+15x+20}{(2x+7)^2 \times \sqrt{2x-1}}; \\
 5.30 &= \frac{12\sqrt{x}+2-x^2}{2\sqrt{x}(x^2+2)\sqrt{x^2+2}}.
 \end{aligned}$$

Murakkab funksiyaning hosilasi

Agar $y = f(x)$ bo'lib, $u = j(x)$ bo'lsa, ya'ni y funksiya x argument bilan oraliq argument orqali bog'langan bo'lsa, y ni x ning murakkab funksiyasi deyiladi.

Murakkab funksiyaning hosilasi, uning oraliq argument bo'yicha hosilasini oraliq argumentning erkli argument bo'yicha hosilasiga ko'paytmasiga teng, ya'ni:

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} \quad \text{yoki} \quad y' = f'(u) \times u'(x).$$

6-masala. Funksiyaning hosilasini toping.

$$y = \frac{e^{x^2}}{1+x^2}.$$

$$\begin{aligned}
 y' &= \frac{e^{x^2}}{1+x^2} = \frac{(e^{x^2})'(1+x^2) - e^{x^2}(1+x^2)'}{(1+x^2)^2} = \\
 &= \frac{e^{x^2} \times 2x \times (1+x^2) - e^{x^2} \times 2x}{(1+x^2)^2} = \frac{2x^3 \times e^{x^2}}{(1+x^2)^2}.
 \end{aligned}$$

$$1. y = x - \ln\left(2 + e^x + 2\sqrt{e^{2x} + e^x + 1}\right)$$

$$2. y = \frac{e^{2x}(2 - \sin 2x - \cos 2x)}{8}.$$

$$3. y = \frac{1}{2} \operatorname{arctg} \frac{e^x - 3}{2}.$$

$$4. y = \frac{1}{\ln 4} \ln \frac{1 + 2^x}{1 - 2^x}.$$

$$5. y = 2\sqrt{e^x + 1} + \ln \frac{\sqrt{e^x + 1} - 1}{\sqrt{e^x + 1} + 1}.$$

$$6. y = \frac{2}{3} \sqrt{(\operatorname{arctg} e^x)^3}.$$

$$7. y = \frac{1}{2} \ln(e^{2x} + 1) - 2 \operatorname{arctg} e^x.$$

$$8. y = \ln(e^x + 1) + \frac{18e^{2x} + 27e^x + 11}{6(e^x + 1)^3}.$$

$$9. y = \frac{2(\sqrt{2^x - 1} - \operatorname{arctg} \sqrt{2^x - 1})}{\ln 2}.$$

$$10. y = 2(x - 2)\sqrt{1 + e^x} - 2 \ln \frac{\sqrt{1 + e^x} - 1}{\sqrt{1 + e^x} + 1}.$$

$$11. y = \frac{e^{ax}(a \sin bx - b \cos bx)}{a^2 + b^2}.$$

$$12. y = \frac{e^{ax}(b \sin bx - a \cos bx)}{a^2 + b^2}.$$

$$13. y = e^{ax} \frac{1}{2a} + \frac{a \cos 2bx + 2b \sin 2bx}{2(a^2 + 4b^2)}.$$

$$14. y = x + \frac{1}{1 + e^x} - \ln(1 + e^x).$$

$$15. y = x - 3 \ln(1 + e^{\frac{x}{6}}) \sqrt{1 + e^{\frac{x}{6}}} - 3 \operatorname{arctg} e^{\frac{x}{6}}.$$

$$16. y = x + \frac{8}{1 + e^{\frac{x}{4}}}.$$

$$17. y = \ln(e^x + \sqrt{e^{2x} - 1}) + \arcsin e^{-x}.$$

$$18. y = x - e^{-x} \arcsin e^x - \ln(1 + \sqrt{1 - e^{2x}}).$$

$$19. y = x - \ln(1 + e^x) - 2e^{-\frac{x}{2}} \arctg e^{\frac{x}{2}} - \left(\arctg e^{-\frac{x}{2}}\right)^2.$$

$$20. y = \frac{e^{x^3}}{1 + x^3}.$$

$$21. y = \frac{1}{m\sqrt{ab}} \arctg \frac{e^{mx}}{e} \times \sqrt{\frac{a}{b}}.$$

$$22. y = 3e^{\sqrt[3]{x}} \left(\sqrt[3]{x^2} - 2\sqrt[3]{x} + 2 \right).$$

$$23. y = \ln \frac{\sqrt{1 + e^x + e^{2x}} - e^x - 1}{\sqrt{1 + e^x + e^{2x}} - e^x + 1}.$$

$$24. y = e^{\sin x} \frac{1}{\cos x}.$$

$$25. y = \frac{e^x}{2} \left((x^2 - 1) \cos x + (x - 1)^2 \sin x \right).$$

$$26. y = \arctg(e^x - e^{-x}).$$

$$27. y = 3e^{\sqrt[3]{x}} \left(\sqrt[3]{x^5} - 5\sqrt[3]{x^4} + 20x - 60\sqrt[3]{x^2} + 120\sqrt[3]{x} - 120 \right).$$

$$28. y = - \frac{e^{3x}}{3 \operatorname{sh}^3 x}.$$

$$29. y = \arcsin e^{-x} - \sqrt{1 - e^{2x}}.$$

$$30. y = - \frac{1}{2} e^{-x^2} (x^4 + 2x^2 + 2).$$

Javoblar. 6.1 $\frac{1}{\sqrt{e^{2x} + e^x + 1}}$; 6.2 $e^{2x} \sin^2 x$; 6.3 $\frac{e^x}{e^{2x} - 6e^x + 13}$; 6.4 $\frac{2^x}{1 - 2^{2x}}$;

6.5 $\sqrt{e^x + 1}$; 6.6 $\frac{e^x \times \sqrt{\arctg e^x}}{1 + e^{2x}}$; 6.7 $\frac{e^{2x} - e^x}{1 + e^{2x}}$; 6.8 $\frac{e^{4x}}{(e^x + 1)^4}$; 6.9 $\sqrt{2^x - 1}$;

6.10 $\frac{2e^x + x \times e^{2x} - 2}{e^x \sqrt{1 + e^x}}$; 6.11 $e^{ax} \times \sin bx$;

6.12 $\frac{e^{ax}(b^2 \times \cos bx + 2ab \sin 3x - a^2 \times \cos bx)}{a^2 + b^2}$; 6.13 $e^{ax} > \cos^2 bx$; 6.14 $\frac{1}{(1 + e^x)^2}$;

6.15 $\frac{1 - e^{x/2} - e^{x/3} - e^{x/6}}{2(1 + e^{x/3})(1 + e^{x/6})}$; 6.16 $\frac{1 + e^{x/2}}{(1 + e^{x/4})^2}$; 6.17 $\sqrt{\frac{e^x - 1}{e^x + 1}}$; 6.18 $e^{-x} \arcsin e^x$;

6.19 $\frac{\arctg e^{x/2}}{e^{x/2} \times (1 + e^x)}$; 6.20 $\frac{3x^5 \times e^{x^3}}{(1 + x^3)^2}$; 6.21 $\frac{e^{mx}}{b + a > e^{2mx}}$; 6.22 $e^{\sqrt[3]{x}}$;

6.23 $\frac{1}{\sqrt{1 + e^x + e^{2x}}}$; 6.24 $e^{\sin x} \times \frac{\sin x}{\cos^2 x}$; 6.25 $x^2 > e^x > \cos x$;

6.26 $\frac{e^{3x} + e^x}{e^{4x} - e^{2x} + 1}$; 6.27 $x \times e^{\sqrt[3]{x}}$; 6.28 $\frac{e^{3x} \times (\operatorname{ch} x - \operatorname{sh} x)}{\operatorname{sh}^4 x}$; 6.29 $\frac{e^x \sqrt{e^{2x} - 1} - \sqrt{e^{-2x} - 1}}{\sqrt{1 - e^{-2x}} \times \sqrt{1 - e^{2x}}}$;

6.30 $x^5 \times e^{-x^2}$.

Funksiyaning hosilasi

7-masala. Funksiyaning hosilasini toping.

$$y = \ln \ln^2 \ln^3 x.$$

$$\begin{aligned} y' &= (\ln \ln^2 \ln^3 x)' = \frac{1}{\ln^2 \ln^3 x} \times 2 \ln \ln^3 x \times \frac{1}{\ln^3 x} \times 3 \ln^2 x \times \frac{1}{x} = \\ &= \frac{2}{\ln \ln^3 x} \times \frac{3}{\ln x} \times \frac{1}{x} = \frac{6}{x \ln x \ln^3 x}. \end{aligned}$$

1. $y = \sqrt{x} \ln(\sqrt{x} + \sqrt{x+a}) - \sqrt{x+a}$. 2. $y = \ln(x + \sqrt{a^2 + x^2})$

3. $y = 2\sqrt{x} - 4 \ln(2 + \sqrt{x})$. 4. $y = \ln \frac{x^2}{\sqrt{1 - ax^4}}$.

5. $y = \ln(\sqrt{x} + \sqrt{x+1})$. 6. $y = \ln \frac{a^2 + x^2}{a^2 - x^2}$.

$$7. y = \ln^2(x + \cos x).$$

$$8. y = \ln^3(x + \cos x).$$

$$9. y = \ln \frac{x^2}{1 - x^2}.$$

$$10. y = \ln \operatorname{tg}\left(\frac{\rho}{4} + \frac{x}{2}\right).$$

$$11. y = \ln \sqrt[4]{\frac{1+2x}{1-2x}}.$$

$$12. y = x + \frac{1}{\sqrt{2}} \ln \frac{x - \sqrt{2}}{x + \sqrt{2}} + a^{\rho^{\sqrt{2}}}.$$

$$13. y = \ln_{\frac{e}{\sin}} \frac{2x+4}{x+1} \frac{\ddot{\circ}}{\varnothing}$$

$$14. y = \log_{16} \log_5 \operatorname{tg} x.$$

$$15. y = \log_4 \log_2 \operatorname{tg} x.$$

$$16. y = \frac{x(\cos(\ln x) + \sin(\ln x))}{2}.$$

$$17. y = \ln \cos \frac{2x+3}{x+1}.$$

$$18. y = \lg \ln(\operatorname{ctg} x).$$

$$19. y = \log_a \frac{1}{\sqrt[6]{1-x^4}} \frac{\ddot{\circ}}{\varnothing}$$

$$20. y = \frac{1}{\sqrt{2}} \ln \left(\sqrt{2} \operatorname{tg} x + \sqrt{1+2\operatorname{tg}^2 x} \right)$$

$$21. y = \ln \left(\arcsin \sqrt{1 - e^{2x}} \right)$$

$$22. y = \ln \arccos \sqrt{1 - e^{4x}}.$$

$$23. y = \ln \left(bx + \sqrt{a^2 + b^2 x^2} \right)$$

$$24. y = \ln \frac{\sqrt{x^2+1} + x\sqrt{2}}{\sqrt{x^2+1} - x\sqrt{2}}.$$

$$25. y = \ln_{\frac{e}{\arccos}} \frac{1}{\sqrt{x}} \frac{\ddot{\circ}}{\varnothing}$$

$$26. y = \ln \left(e^x + \sqrt{1 + e^{2x}} \right)$$

$$27. y = \ln \frac{\sqrt{5} + \operatorname{tg} \frac{x}{2}}{\sqrt{5} - \operatorname{tg} \frac{x}{2}}.$$

$$28. y = \ln_{\frac{e}{\sin}} \frac{\ln x}{\sin \frac{1}{x}} \frac{\ddot{\circ}}{\varnothing}$$

$$29. y = \ln \ln \sin \left(1 + \frac{1}{x} \right).$$

$$30. y = \ln \ln^3 \ln^2 x.$$

Javoblar. 7.1 $\frac{1}{2\sqrt{x}} \times \ln(\sqrt{x} + \sqrt{x+a})$; 7.2 $\frac{1}{\sqrt{a^2+x^2}}$; 7.3 $\frac{\sqrt{x}}{\sqrt{x} \times (2+\sqrt{x})}$; 7.4 $\frac{2}{x - ax^5}$;

7.5 $\frac{1}{2\sqrt{x^2+x}}$; 7.6 $\frac{4a^2x}{a^4 - x^4}$; 7.7 $\frac{1 - \sin x}{x + \cos x} \times 2 \ln(x + \cos x)$;

7.8 $-\frac{3 \times \sin x \times \ln^2(1 + \cos x)}{1 + \cos x}$; 7.9 $\frac{2}{x \times (1 - x^2)}$; 7.10 $\frac{1}{\cos x}$; 7.11 $\frac{1}{1 - 4x^2}$; 7.12 $\frac{x^2}{x^2 - 2}$;

$$\begin{aligned}
& 7.13 - \frac{2}{(x+1)^2} \times \operatorname{ctg} \frac{2x+4}{x+1}; 7.14 \frac{1}{\sin 2x \times \ln 4 \times \ln \operatorname{tg} x}; 7.15 \frac{1}{\sin 2x \times \ln 2 \times \ln \operatorname{tg} x}; \\
& 7.16 \cos(\ln x); 7.17 \frac{\operatorname{tg} \frac{2x+3}{x+1}}{(x+1)^2}; 7.18 - \frac{2}{\ln(\operatorname{ctg} x) \times \ln 10 \times \sin 2x}; 7.19 \frac{2x^3}{\ln a \times (1-x^4)}; \\
& 7.20 \frac{1}{\cos^2 x \sqrt{1+2\operatorname{tg}^2 x}}; 7.21 - \frac{e^x}{\sqrt{1-e^{2x}} \times \arcsin \sqrt{1-e^{2x}}}; \\
& 7.22 \frac{2e^{2x}}{\sqrt{1-e^{4x}} \times \arccos \sqrt{1-e^{4x}}}; 7.23 \frac{b}{\sqrt{a^2+b^2x^2}}; 7.24 \frac{2\sqrt{2}}{(1-x^2)\sqrt{x^2+x}}; \\
& 7.25 \frac{1}{2x \times \sqrt{x-1} \times \arccos \frac{1}{\sqrt{x}}}; 7.26 \frac{e^x}{\sqrt{1+e^{2x}}}; 7.27 \frac{\sqrt{5}}{6\cos^2 \frac{x}{2} - 1}; \\
& 7.28 \frac{\sin \frac{1}{x} - x \times \ln x \times \cos \frac{1}{x}}{x \times \ln x \times \sin \frac{1}{x}}; 7.29 - \frac{\operatorname{ctg}(1+\frac{1}{x})}{x^2 \times \ln \sin(1+\frac{1}{x})}; 7.30 \frac{6}{x \times \ln x \times \ln \ln^2 x}.
\end{aligned}$$

Funksiyaning hosilasi

8-masala. Funksiyaning hosilasini toping.

$$y = \operatorname{tg} \sqrt{\cos \frac{1}{3} + \frac{\sin^2 31x}{31 \cos 62x}}.$$

$$\begin{aligned}
y' &= \operatorname{ctg} \sqrt{\cos \frac{1}{3} + \frac{\sin^2 31x}{31 \cos 62x}} = \frac{\sin^2 31x}{31 \cos 62x} = \\
&= \frac{2 \sin^2 31x}{62 \cos 62x} = \frac{1 - \cos 62x}{62 \cos 62x} = \frac{1}{62 \cos 62x} - \frac{1}{62} = \\
&= \frac{1}{62 \cos 62x} - \frac{1}{62 \cos^2 62x} \times (-\sin 62x) \times 62 = \frac{\operatorname{tg} 62x}{\cos 62x}.
\end{aligned}$$

$$1. y = \sin \sqrt{3} + \frac{1}{3} \times \frac{\sin^2 3x}{\cos 6x}.$$

$$2. y = \cos \ln 2 - \frac{1}{3} \times \frac{\cos^2 3x}{\sin 6x}.$$

$$3. y = \operatorname{tg} \lg \frac{1}{3} + \frac{1}{4} \times \frac{\sin^2 4x}{\cos 8x}.$$

$$4. y = \frac{\cos(\sin 5) \times \sin^2 2x}{2 \cos 4x}.$$

$$5. y = \operatorname{ctg} \sqrt[3]{5} - \frac{1}{8} \times \frac{\cos^2 4x}{\sin 8x}.$$

$$7. y = \frac{\cos \ln 7 \times \sin^2 7x}{7 \cos 14x}.$$

$$9. y = \operatorname{ctg}(\cos 2) + \frac{1}{6} \times \frac{\sin^2 6x}{\cos 12x}.$$

$$11. y = \frac{1}{3} \times \cos \operatorname{ctg} \frac{1}{2} + \frac{1}{10} \times \frac{\sin^2 10x}{\cos 20x}.$$

$$13. y = 8 \sin(\operatorname{ctg} 3) + \frac{1}{5} \times \frac{\sin^2 5x}{\cos 10x}.$$

$$15. y = \frac{\cos\left(\operatorname{tg} \frac{1}{3}\right) \times \sin^2 15x}{15 \cos 30x}.$$

$$17. y = \frac{\operatorname{ctg}\left(\sin \frac{1}{3}\right) \times \sin^2 17x}{17 \cos 34x}.$$

$$19. y = \frac{\operatorname{tg}(\ln 2) \times \sin^2 19x}{19 \cos 38x}.$$

$$21. y = \sqrt{\operatorname{tg} 4} + \frac{\sin^2 21x}{21 \cos 42x}.$$

$$23. y = \ln \cos \frac{1}{3} + \frac{\sin^2 23x}{23 \cos 46x}.$$

$$25. y = \sin(\ln 2) + \frac{\sin^2 25x}{25 \cos 50x}.$$

$$27. y = \sqrt[7]{\operatorname{tg}(\cos 2)} + \frac{\sin^2 27x}{27 \cos 54x}.$$

$$29. y = \cos^2(\sin 3) + \frac{\sin^2 29x}{29 \cos 58x}.$$

$$6. y = \frac{\sin(\cos 3) \times \cos^2 2x}{4 \sin 4x}.$$

$$8. y = \cos(\operatorname{ctg} 2) - \frac{1}{16} \times \frac{\cos^2 8x}{\sin 16x}.$$

$$10. y = \sqrt[3]{\operatorname{ctg} 2} - \frac{1}{20} \times \frac{\cos^2 10x}{\sin 20x}.$$

$$12. y = \ln \sin \frac{1}{2} - \frac{1}{24} \times \frac{\cos^2 12x}{\sin 24x}.$$

$$14. y = \frac{\sin(\operatorname{ctg} 3) \times \cos^2 14x}{28 \sin 28x}.$$

$$16. y = \frac{\sin\left(\operatorname{tg} \frac{1}{7}\right) \times \cos^2 16x}{32 \sin 32x}.$$

$$18. y = \frac{\sqrt[5]{\operatorname{ctg} 2} \times \cos^2 18x}{36 \sin 36x}.$$

$$20. y = \operatorname{ctg}(\cos 5) - \frac{1}{40} \times \frac{\cos^2 20x}{\sin 40x}.$$

$$22. y = \cos(\ln 13) - \frac{1}{44} \times \frac{\cos^2 22x}{\sin 44x}.$$

$$24. y = \operatorname{ctg} \sin \frac{1}{13} - \frac{1}{48} \times \frac{\cos^2 24x}{\sin 48x}.$$

$$26. y = \sqrt[3]{\cos \sqrt{2}} - \frac{1}{52} \times \frac{\cos^2 26x}{\sin 52x}.$$

$$28. y = \sin \sqrt[3]{\operatorname{tg} 2} - \frac{\cos^2 28x}{56 \sin 56x}.$$

$$30. y = \sin^3(\cos 2) - \frac{\cos^2 30x}{60 \sin 60x}.$$

Javoblar. 8.1 $\frac{\operatorname{tg} 6x}{\cos 6x}$; 8.2 $\frac{1}{2 \sin^2 3x}$; 8.3 $\frac{\operatorname{tg} 8x}{\cos 8x}$; 8.4 $\frac{\cos(\sin 5) \times \operatorname{tg} 4x}{\cos 4x}$; 8.5 $\frac{1}{4 \sin^2 4x}$;

8.6 $-\frac{\sin(\cos 3)}{4 \sin^2 2x}$; 8.7 $\frac{\cos \ln 7 \times \operatorname{tg} 14x}{\cos 14x}$; 8.8 $\frac{1}{4 \sin^2 8x}$; 8.9 $\frac{\operatorname{tg} 12x}{\cos 12x}$; 8.10 $\frac{1}{4 \sin^2 10x}$;

$$\begin{aligned}
& \text{8.11 } \frac{\operatorname{tg} 20x}{\cos 20x}; \text{8.12 } \frac{1}{4\sin^2 12x}; \text{8.13 } \frac{\operatorname{tg} 10x}{\cos 10x}; \text{8.14 } -\frac{\cos(\operatorname{ctg} 3)}{4\sin^2 14x}; \text{8.15 } \frac{\cos(\operatorname{tg} \frac{1}{3}) \operatorname{tg} 30x}{\cos 30x}; \\
& \text{8.16 } -\frac{\sin(\operatorname{tg} \frac{1}{7})}{4\sin^2 16x}; \text{8.17 } \frac{\operatorname{ctg}(\sin \frac{1}{3}) \operatorname{tg} 34x}{\cos 34x}; \text{8.18 } -\frac{\sqrt[5]{\operatorname{ctg} 2}}{4\sin^2 18x}; \text{8.19 } \frac{\operatorname{tg}(\ln 2) \operatorname{tg} 38x}{\cos 38x}; \\
& \text{8.20 } \frac{1}{4\sin^2 20x}; \text{8.21 } \frac{\operatorname{tg} 42x}{\cos 42x}; \text{8.22 } \frac{1}{4\sin^2 22x}; \text{8.23 } \frac{\operatorname{tg} 46x}{\cos 46x}; \text{8.24 } \frac{1}{4\sin^2 24x}; \\
& \text{8.25 } \frac{\sin 50x}{\cos^2 50x}; \text{8.26 } \frac{1}{2\sin^2 26x}; \text{8.27 } \operatorname{tg} 54x \operatorname{tg} 54x; \text{8.28 } \frac{1}{4\sin^2 28x}; \text{8.29 } \frac{\operatorname{tg} 58x}{\cos 58x}; \quad \text{8.30} \\
& \frac{1}{4\sin^2 30x}.
\end{aligned}$$

Funksiyaning hosilasi

9-masala. Funksiyaning hosilasini toping.

$$\begin{aligned}
y' &= \frac{1}{1 + \operatorname{tg}^2 \frac{x}{2} + 1} \times \frac{1}{2} \times \frac{1}{\cos^2 \frac{x}{2}} \times \frac{1}{2} = \\
&= \frac{1}{\operatorname{tg}^2 \frac{x}{2} + 2\operatorname{tg} \frac{x}{2} + 5} \times \frac{1}{2} \times \frac{1}{\cos^2 \frac{x}{2}} \times \frac{1}{2} = \\
&= \frac{1}{\sin^2 \frac{x}{2} + 2\sin \frac{x}{2} \cos \frac{x}{2} + 5\cos^2 \frac{x}{2}} = \frac{1}{1 + \sin x + 4\cos^2 \frac{x}{2}} = \\
&= \frac{1}{\sin x + 2\cos x + 2} = \frac{1}{\sin x + 2\cos x + 3}.
\end{aligned}$$

$$1. y = \arctg \frac{\operatorname{tg} x - \operatorname{ctg} x}{\sqrt{2}}.$$

$$2. y = \arcsin \frac{\sqrt{x} - 2}{\sqrt{5x}}.$$

$$3. y = \frac{2x-1}{4} \times \sqrt{2+x-x^2} + \frac{9}{8} \times \arcsin \frac{2x-1}{3}.$$

$$4. y = \operatorname{arctg} \frac{\sqrt{1+x^2} - 1}{x}.$$

$$5. y = \arccos \frac{x^2 - 4}{\sqrt{x^2 + 16}}.$$

$$6. y = \sqrt{\frac{2}{3}} \times \operatorname{arctg} \frac{3x - 1}{\sqrt{6x}}.$$

$$7. y = \frac{1}{4} \times \ln \frac{x - 1}{x + 1} - \frac{1}{2} \operatorname{arctg} x.$$

$$8. y = \frac{1}{2} \times (x - 4) \sqrt{8x - x^2} - 7 - 9 \arccos \sqrt{\frac{x - 1}{6}}.$$

$$9. y = \frac{(1+x) \operatorname{arctg} \sqrt{x}}{x^2} + \frac{1}{3x\sqrt{x}}.$$

$$10. y = \frac{x^3}{3} \times \arccos x - \frac{2+x^2}{9} \times \sqrt{1-x^2}.$$

$$11. y = \frac{1}{2\sqrt{x}} + \frac{1+x}{2x} \times \operatorname{arctg} \sqrt{x}.$$

$$12. y = \frac{3+x}{2} \times \sqrt{x(2-x)} + 3 \arccos \sqrt{\frac{x}{2}}.$$

$$13. y = \frac{4+x^4}{x^3} \times \operatorname{arctg} \frac{x^2}{2} + \frac{4}{x}.$$

$$14. y = \arcsin \sqrt{\frac{x}{x+1}} + \operatorname{arctg} \sqrt{x}.$$

$$15. y = \frac{1}{2} \times \sqrt{\frac{1}{x^2} - 1} - \frac{\arccos x}{2x^2}.$$

$$16. y = 6 \arcsin \frac{\sqrt{x}}{2} - \frac{6+x}{2} \times \sqrt{x(4-x)}.$$

$$17. y = \frac{x-3}{2} \sqrt{6x-x^2} - 8 + \arcsin \sqrt{\frac{x}{2} - 1}.$$

$$18. y = \frac{(1+x)\operatorname{arctg}\sqrt{x} - \sqrt{x}}{x}.$$

$$19. y = \frac{2\sqrt{1-x} \times \arcsin \sqrt{x}}{x} + \frac{2}{\sqrt{x}}.$$

$$20. y = \frac{2x-5}{4} \times \sqrt{5x-4-x^2} + \frac{9}{4} \times \arcsin \sqrt{\frac{x-1}{3}}.$$

$$21. y = \operatorname{arctg} x + \frac{5}{6} \times \ln \frac{x^2+1}{x^2+4}.$$

$$22. y = \arcsin \frac{x-2}{(x-1)\sqrt{2}}.$$

$$23. y = \sqrt{1-x^2} - x \times \arcsin \sqrt{1-x^2}.$$

$$24. y = \sqrt{x} + \frac{1}{3} \times \operatorname{arctg} \sqrt{x} + \frac{8}{3} \times \operatorname{arctg} \frac{\sqrt{x}}{2}.$$

$$25. y = \operatorname{arctg} \frac{\sqrt{1-x}}{1-\sqrt{x}}.$$

$$26. y = (2x^2 + 6x + 5) \operatorname{arctg} \frac{x+1}{x+2} - x.$$

$$27. y = \frac{x}{2\sqrt{1-4x^2}} \arcsin 2x + \frac{1}{8} \times \ln(1-4x^2).$$

$$28. y = \frac{2}{3} 2x^2 - x + \frac{1}{2} \times \operatorname{arctg} \frac{x^2-1}{x\sqrt{3}} - \frac{x^2}{2\sqrt{3}} - \frac{\sqrt{3}}{2} \times x.$$

$$29. y = (x - 2\sqrt{x} + 2) \operatorname{arctg} \frac{\sqrt{x}}{\sqrt{x}+2} - \sqrt{x}.$$

$$30. y = \sqrt{1+2x-x^2} \arcsin \frac{x\sqrt{2}}{1+x} - \sqrt{2} \ln(1+x).$$

Javoblar. 9.1 $\frac{\sqrt{2}}{\sin^4 x + \cos^4 x}$; 9.2 $\frac{1}{2x\sqrt{x+\sqrt{x}-1}}$; 9.3 $\sqrt{2+x-x^2}$;

9.4 $\frac{-1+\sqrt{1+x^2}}{(x^2+(\sqrt{1+x^2}-1)^2)\sqrt{1+x^2}}$; 9.5 $-\frac{2\sqrt{2}(4+x^2)}{x^4+16}$; 9.6 $\frac{3x+1}{\sqrt{x}(9x^2+1)}$; 9.7 $\frac{1}{x^4-1}$;

$$\begin{aligned}
& \text{9.8 } \sqrt{8x - x^2 - 7}; \text{9.9 } - \frac{(2+x) \operatorname{arctg} \sqrt{x}}{x^3}; \text{9.10 } x^2 > \arccos x; \\
& \text{9.11 } - \frac{1}{2x^2} \operatorname{arctg} \sqrt{x}; \text{9.12 } - \frac{x^2}{\sqrt{x(2-x)}}; \text{9.13 } \frac{x^4 - 12}{x^4} \operatorname{arctg} \frac{x^2}{2}; \text{9.14 } \frac{1}{\sqrt{x(x+1)}}; \\
& \text{9.15 } \frac{x + \sqrt{1-x^2} \arccos x}{x^3 \sqrt{1-x^2}}; \text{9.16 } \frac{x^2 - 3}{\sqrt{x(4-x)}}; \text{9.17 } \sqrt{6x - x^2 - 8}; \\
& \text{9.18 } - \frac{1}{x^2} \operatorname{arctg} \sqrt{x} + \frac{1}{x\sqrt{x}}; \text{9.19 } \frac{x-2}{x^2 \sqrt{1-x}} \arcsin \sqrt{x}; \text{9.20 } \sqrt{5x - 4 - x^2}; \\
& \text{9.21 } \frac{x^2 + 9}{(1+x^2) \sqrt{x^2 + 4}}; \text{9.22 } \frac{1}{(x-1) \sqrt{x^2 - 2}}; \text{9.23 } - \arcsin \sqrt{1-x^2}; \\
& \text{9.24 } \frac{3x^2 + 16x + 32}{6\sqrt{x}(x+1)(x+4)}; \text{9.25 } \frac{1}{4\sqrt{x(1-x)}}; \text{9.26 } (4x+6) \operatorname{arctg} \frac{x+1}{x+2}; \\
& \text{9.27 } \frac{\arcsin 2x}{2(1-4x^2) \sqrt{1-4x^2}}; \text{9.28 } (4x-1) \operatorname{arctg} \frac{x^2-1}{x\sqrt{3}} + \frac{\sqrt{3}(x^2+1)(3x^2-2x-x^4)}{2(x^4+x^2+1)}; \\
& \text{9.29 } \frac{1}{\sqrt{x}} \operatorname{arctg} \frac{\sqrt{x}}{\sqrt{x+2}}; \text{9.30 } \frac{1-x}{\sqrt{1+2x-x^2}} \arcsin \frac{x\sqrt{2}}{1+x}.
\end{aligned}$$

Funksiyaning hosilasi

10-masala. Funksiyaning hosilasini toping.

$$y = \frac{2}{3} \operatorname{th} x - \frac{\operatorname{ch} x}{3 \operatorname{sh}^3 x}.$$

$$\begin{aligned}
y' &= \frac{2}{3} \operatorname{th} x - \frac{\operatorname{ch} x}{3 \operatorname{sh}^3 x} = -\frac{2}{3} \frac{1}{\operatorname{sh}^2 x} - \frac{(\operatorname{ch} x) \operatorname{sh}^3 x - \operatorname{ch} x (\operatorname{sh}^3 x)'}{3 \operatorname{sh}^6 x} = \\
&= -\frac{2}{3} \frac{1}{\operatorname{sh}^2 x} - \frac{\operatorname{sh} x \operatorname{sh}^3 x - \operatorname{ch} x \times 3 \operatorname{sh}^2 x \times \operatorname{ch} x}{3 \operatorname{sh}^6 x} = -\frac{2 \operatorname{sh}^2 x}{3 \operatorname{sh}^4 x} - \frac{\operatorname{sh}^2 x - 3 \operatorname{ch}^2 x}{3 \operatorname{sh}^4 x} = \\
&= \frac{-2 \operatorname{sh}^2 x - \operatorname{sh}^2 x + 3 \operatorname{ch}^2 x}{3 \operatorname{sh}^4 x} = \frac{3 \operatorname{ch}^2 x - 3 \operatorname{sh}^2 x}{3 \operatorname{sh}^4 x} = \frac{3}{3 \operatorname{sh}^4 x} = \frac{1}{\operatorname{sh}^4 x}.
\end{aligned}$$

$$1. y = \frac{1}{4\sqrt{5}} \ln \frac{2 + \sqrt{5} \operatorname{th} x}{2 - \sqrt{5} \operatorname{th} x}.$$

$$2. y = \frac{\operatorname{sh} x}{4 \operatorname{ch}^4 x} + \frac{3 \operatorname{sh} x}{8 \operatorname{ch}^2 x} + \frac{3}{8} \operatorname{arctg}(\operatorname{sh} x).$$

$$3. y = \frac{1}{2} \ln \frac{1 + \sqrt{\operatorname{th} x}}{1 - \sqrt{\operatorname{th} x}} + \operatorname{arctg} \sqrt{\operatorname{th} x}.$$

$$4. y = \frac{3}{8\sqrt{2}} \ln \frac{\sqrt{2} - \operatorname{th} x}{\sqrt{2} + \operatorname{th} x} - \frac{\operatorname{th} x}{4(2 - \operatorname{th}^2 x)}.$$

$$5. y = \frac{1}{2} \operatorname{th} x + \frac{1}{4\sqrt{2}} \ln \frac{1 + \sqrt{2} \operatorname{th} x}{1 - \sqrt{2} \operatorname{th} x}.$$

$$6. y = -\frac{1}{2} \ln \frac{\operatorname{ch} x}{\operatorname{sh} x} - \frac{x}{2 \operatorname{sh}^2 x} - \frac{\operatorname{ch} x}{2 \operatorname{sh}^2 x}.$$

$$7. y = \frac{1}{2a\sqrt{1+a^2}} \ln \frac{a + \sqrt{1+a^2} \operatorname{th} x}{a - \sqrt{1+a^2} \operatorname{th} x}.$$

$$8. y = \frac{1}{18\sqrt{2}} \ln \frac{1 + \sqrt{2} \operatorname{cth} x}{1 - \sqrt{2} \operatorname{cth} x}.$$

$$9. y = \operatorname{arctg} \frac{\sqrt{\operatorname{sh} 2x}}{\operatorname{ch} x - \operatorname{sh} x}.$$

$$10. y = \frac{1}{6} \ln \frac{1 - \operatorname{sh} 2x}{1 + \operatorname{sh} 2x}.$$

$$11. y = \sqrt[4]{\frac{1 + \operatorname{th} x}{1 - \operatorname{th} x}}.$$

$$12. y = \frac{\operatorname{sh} x}{1 + \operatorname{ch} x}.$$

$$13. y = \frac{\operatorname{ch} x}{\sqrt{\operatorname{sh} 2x}}.$$

$$14. y = \frac{\operatorname{sh} 3x}{\sqrt{\operatorname{ch} 6x}}.$$

$$15. y = \frac{1 + 8\operatorname{ch}^2 x \times \ln(\operatorname{ch} x)}{2\operatorname{ch}^2 x}.$$

$$16. y = - \frac{12\operatorname{sh}^2 x + 1}{3\operatorname{sh}^2 x}.$$

$$17. y = - \frac{\operatorname{sh} x + 1}{2\operatorname{ch}^2 x} + \frac{3}{2} \times \arcsin(\operatorname{th} x).$$

$$18. y = \frac{1}{\sqrt{8}} \arcsin \frac{3 + \operatorname{ch} x}{1 + 3\operatorname{ch} x}.$$

$$19. y = \frac{1}{\sqrt{8}} \arcsin \frac{4 + \sqrt{8} \operatorname{th} \frac{x}{2}}{4 - \sqrt{8} \operatorname{th} \frac{x}{2}}.$$

$$20. y = \frac{1}{4} \ln \left| \operatorname{th} \frac{x}{2} \right| - \frac{1}{4} \times \ln \frac{3 + \operatorname{ch} x}{\operatorname{sh} x}.$$

$$21. y = - \frac{1}{4} \arcsin \frac{5 + 3\operatorname{ch} x}{3 + 5\operatorname{ch} x}.$$

$$22. y = \frac{1 - 8\text{ch}^2 x}{4\text{ch}^4 x}.$$

$$23. y = \frac{2}{\text{sh} x} - \frac{1}{3\text{sh}^3 x} + \frac{\text{sh} x}{2\text{ch}^2 x} + \frac{5}{2} \text{arctg}(\text{sh} x).$$

$$24. y = \frac{8}{3} \times \text{cth} x - \frac{1}{3\text{ch} x \times \text{sh}^3 x}.$$

$$25. y = \frac{1}{2} \times \text{arctg}(\text{sh} x) - \frac{\text{sh} x}{2\text{ch}^2 x}.$$

$$26. y = \frac{3}{2} \times \ln\left(\text{th} \frac{x}{2}\right) + \text{ch} x - \frac{\text{ch} x}{2\text{sh}^2 x}.$$

$$27. y = -\frac{\text{sh} x}{2\text{ch}^2 x} - \frac{1}{\text{sh} x} - \frac{3}{2} \text{arctg}(\text{sh} x).$$

$$28. y = \frac{\text{sh} x}{2\text{ch}^2 x} + \frac{1}{2} \times \text{arctg}(\text{sh} x).$$

$$29. y = \frac{1}{2} \frac{\text{sh} x}{\text{ch}^2 x} + \text{arctg}(\text{sh} x) \frac{\ddot{\circ}}{\emptyset}$$

$$30. y = -\frac{\text{ch} x}{2\text{sh}^2 x} - \frac{1}{2} \ln \frac{\text{sh} x}{\text{ch} x} \frac{\ddot{\circ}}{\emptyset}$$

Javoblar. 10.1 $\frac{1}{4 - \text{sh}^2 x}$; 10.2 $\frac{1 - 3\text{ch}^2 x}{4\text{ch}^5 x}$; 10.3 $\sqrt{\text{th} x}$; 10.4 $\frac{1}{(1 + \text{ch}^2 x)^2}$;

10.5 $\frac{1}{\text{ch}^2 x (1 - \text{sh}^2 x)}$; 10.6 $\frac{1}{\text{sh}^3 x}$; 10.7 $\frac{1}{a^2 \times \text{ch}^2 x + (1 + a^2) \times \text{sh}^2 x}$; 10.8 $\frac{1}{9 \times (1 + \text{ch}^2 x)}$;

10.9 $\frac{\text{ch} x + \text{sh} x}{\sqrt{\text{sh} 2x} \times \text{ch} 2x}$; 10.10 $\frac{\text{ch} 2x}{\text{sh}^2 2x + \text{sh} 2x - 2}$; 10.11 $\frac{1}{2\sqrt{\text{ch} x - \text{sh} x}}$; 10.12 $\frac{1}{1 + \text{ch} x}$;

$$\begin{aligned}
& \text{10.13} - \frac{1}{2\operatorname{sh}x\sqrt{\operatorname{sh}2x}}; \text{10.14} \frac{3\operatorname{ch}3x}{\operatorname{ch}6x\sqrt{\operatorname{ch}6x}}; \text{10.15} \frac{\operatorname{sh}x(4\operatorname{ch}^2x-1)}{\operatorname{ch}^3x}; \text{10.16} - \frac{2\operatorname{ch}x}{3\operatorname{sh}^3x}; \\
& \text{10.17} \frac{\operatorname{ch}2x}{\operatorname{ch}^3x}; \quad \text{10.18} - \frac{9\operatorname{sh}x}{8(1+3\operatorname{ch}x)}; \text{10.19} \frac{1}{2(\operatorname{ch}^2\frac{x}{2}+1)}; \text{10.20} \frac{1}{2\operatorname{sh}x}; \\
& \text{10.21} \frac{1}{3+5\operatorname{ch}x}; \text{10.22} \frac{4\operatorname{th}^3x}{\operatorname{ch}^2x}; \text{10.23} - \frac{2\operatorname{ch}x}{\operatorname{sh}^2x} + \frac{\operatorname{ch}x}{\operatorname{sh}^4x} + \frac{1-\operatorname{sh}^2x}{2\operatorname{ch}^3x} + \frac{5}{2\operatorname{ch}x}; \\
& \text{10.24} \frac{1-4\operatorname{sh}^3x}{\operatorname{ch}^2x\sqrt{\operatorname{sh}^4x}}; \text{10.25} \frac{\operatorname{sh}^2x}{\operatorname{ch}^3x}; \text{10.26} \frac{\operatorname{ch}^4x}{\operatorname{sh}^3x}; \text{10.27} \frac{\operatorname{ch}x}{\operatorname{sh}^2x} - \frac{2-\operatorname{sh}^2x}{\operatorname{ch}^3x}; \text{10.28} \frac{1}{\operatorname{ch}^3x}; \\
& \text{10.29} \frac{1}{\operatorname{ch}^3x}; \text{10.30} \frac{1}{\operatorname{sh}^3x}.
\end{aligned}$$

Funksiyaning hosilasi

11-masala. Funksiyaning hosilasini toping.

$$y = x^{e^x} \times x^9.$$

$$\ln y = \ln(x^{e^x} \times x^9) = e^x \times \ln x + 9 \times \ln x = \ln x \times (e^x + 9).$$

$$\frac{y'}{y} = (\ln x \times (e^x + 9))' = \frac{1}{x} \times (e^x + 9) + \ln x \times e^x = e^x \ln x + \frac{1}{x} + \frac{9}{x}.$$

$$y' = y \times e^x \ln x + \frac{1}{x} + \frac{9}{x} = x^{e^x} \times x^9 \times e^x \ln x + \frac{1}{x} + \frac{9}{x}.$$

$$1. y = (\operatorname{arctg} x)^{\frac{1}{2} \ln(\operatorname{arctg} x)}.$$

$$2. y = (\sin \sqrt{x})^{\ln(\sin \sqrt{x})}.$$

$$3. y = (\sin x)^{5e^x}.$$

$$4. y = (\arcsin x)^{e^x}.$$

$$5. y = (\ln x)^{3^x}.$$

$$6. y = x^{\arcsin x}.$$

$$7. y = (\operatorname{ctg} 3x)^{2e^x}.$$

$$8. y = x^{e^{\operatorname{tg} x}}.$$

$$9. y = (\operatorname{tg} x)^{4e^x}.$$

$$10. y = (\cos 5x)^{e^x}.$$

$$11. y = (x \sin x)^{8 \ln(x \sin x)}.$$

$$12. y = (x - 5)^{\operatorname{ch} x}.$$

$$13. y = (x^3 + 4)^{\operatorname{tg} x}.$$

$$14. y = x^{\sin x^3}.$$

15. $y = (x^2 - 1)^{\operatorname{sh} x}$.

16. $y = (x^4 + 5)^{\operatorname{ctg} x}$.

17. $y = (\sin x)^{\frac{5x}{2}}$.

18. $y = (x^2 + 1)^{\cos x}$.

19. $y = 19^{x^{19}} \times x^{19}$.

20. $y = x^{3^x} \times 2^x$.

21. $y = (\sin \sqrt{x})^{e^{1/x}}$.

22. $y = x^{e^{\operatorname{ctg} x}}$.

23. $y = x^{e^{\cos x}}$.

24. $y = x^{2^x} \times 5^x$.

25. $y = x^{e^{\sin x}}$.

26. $y = (\operatorname{tg} x)^{\ln \frac{\operatorname{tg} x}{4}}$.

27. $y = x^{e^{\operatorname{arctg} x}}$.

28. $y = (x^8 + 4)^{\operatorname{th} x}$.

29. $y = 29^{x^{29}} \times x^{29}$.

30. $y = \cos 2x^{\frac{\ln \cos 2x}{4}}$

Javoblar. 11.1 $(\operatorname{arctg} x)^{\frac{1}{2} \ln(\operatorname{arctg} x)} \times \frac{\ln(\operatorname{arctg} x)}{\operatorname{arctg} x \times (1 + x^2)};$

11.2 $(\sin \sqrt{x})^{\ln(\sin \sqrt{x})} \times \frac{\ln(\sin \sqrt{x}) \operatorname{ctg} \sqrt{x}}{\sqrt{x}};$

11.3 $5e^x \times (\sin x)^{5e^x} \times (\ln(\sin x) + \operatorname{ctg} x);$

11.4 $(\arcsin x)^{e^x} \times e^x \times \frac{\ln(\arcsin x) + \frac{1}{\sqrt{1-x^2} \arcsin x}}{e};$

11.5 $(\ln x)^{3^x} \times 3^x \times \frac{\ln(\ln 3 \times \ln(\ln x)) + \frac{1}{x \ln x}}{e};$ **11.6** $x^{\arcsin x} \times \frac{\ln x}{e \sqrt{1-x^2}} + \frac{\arcsin x}{x};$

11.7 $2e^x \times (\operatorname{ctg} 3x)^{2e^x} \times \frac{\ln(\operatorname{ctg} 3x) - \frac{6}{\sin 6x}}{e};$ **11.8** $x^{e^{\operatorname{tg} x}} \times e^{\operatorname{tg} x} \times \frac{\ln x}{e \cos^2 x} + \frac{1}{x};$

11.9 $(\operatorname{tg} x)^{4e^x} \times 4e^x \times \frac{\ln(\operatorname{tg} x) + \frac{2}{\sin 2x}}{e};$ **11.10** $(\cos 5x)^{e^x} \times e^x \times (\ln(\cos 5x) - \operatorname{tg} 5x);$

11.11 $\frac{16(x \sin x)^{8 \ln(x \sin x)} \times \ln(x \sin x) \times (1 + x \times \operatorname{ctg} x)}{x};$

11.12 $(x - 5)^{\operatorname{ch} x} \times \frac{\operatorname{sh} x \times \ln(x - 5) + \frac{\operatorname{ch} x}{x - 5}}{e};$ **11.13** $(x^3 + 4)^{\operatorname{tg} x} \times \frac{\ln(x^3 + 4)}{e \sqrt{1 + x^2}} + \frac{3x^2 \times \operatorname{tg} x}{x^3 + 4};$

$$11.14 \quad x^{\sin x^3} \times 3x^2 \times \ln x \times \cos x^3 + \frac{\sin x^3}{x};$$

$$11.15 \quad (x^2 - 1)^{\operatorname{sh} x} \times \operatorname{ch} x \times \ln(x^2 - 1) + \frac{2x \times \operatorname{sh} x}{x^2 - 1};$$

$$11.16 \quad (x^4 + 5)^{\operatorname{ctg} x} \times \frac{x^3 \times \operatorname{ctg} x}{x^4 + 5} - \frac{\ln(x^4 + 5)}{\sin^2 x}; \quad 11.17 \quad \frac{5}{2} \times (\sin x)^{\frac{5x}{2}} \times (\ln(\sin x) + x \times \operatorname{ctg} x);$$

$$11.18 \quad (x^2 + 1)^{\cos x} \times \frac{x \times \cos x}{x^2 + 1} - \sin x \times \ln(x^2 + 1);$$

$$11.19 \quad 19^{x^{19}} \times x^{19} \times \ln 19 + \frac{1}{x}; \quad 11.20 \quad x^{3^x} \times 2^x \times 3^x \times \ln 3 \times \ln(x) + \frac{3^x}{x} + \ln 2;$$

$$11.21 \quad (\sin \sqrt{x})^{e^{1/x}} \times e^{\frac{1}{x}} \times \frac{\ln(\sin \sqrt{x})}{x} + \frac{\operatorname{tg} \sqrt{x}}{2\sqrt{x}}; \quad 11.22 \quad x^{e^{\operatorname{ctg} x}} \times e^{\operatorname{ctg} x} \times \frac{1}{x} - \frac{\ln x}{\sin^2 x};$$

$$11.23 \quad x^{e^{\cos x}} \times e^{\cos x} \times \frac{1}{x} - \sin x \times \ln x; \quad 11.24 \quad x^{2^x} \times 5^x \times 2^x \times \ln 2 \times \ln(x) + \frac{2^x}{x} + \ln 5;$$

$$11.25 \quad x^{e^{\sin x}} \times e^{\sin x} \times \cos x \times \ln x + \frac{1}{x}; \quad 11.26 \quad \operatorname{tg} x^{\frac{\ln \operatorname{tg} x}{4}} \times \frac{\ln(\operatorname{tg} x)}{\sin 2x};$$

$$11.27 \quad x^{e^{\operatorname{arctg} x}} \times e^{\operatorname{arctg} x} \times \frac{\ln x}{1 + x^2} + \frac{1}{x}; \quad 11.28 \quad (x^8 + 1)^{\operatorname{th} x} \times \frac{\ln(x^8 + 1)}{\operatorname{ch}^2 x} + \frac{8x^7 \times \operatorname{th} x}{x^8 + 1};$$

$$11.29 \quad x^{29^x} \times 29^x \times 29^x \times \ln 29 \times \ln(x) + \frac{29^x}{x} + \ln 29;$$

$$11.30 \quad -\cos 2x^{\frac{\ln \cos 2x}{4}} \times \operatorname{tg} 2x \times \ln(\cos 2x).$$

Funksiyaning hosilasi

12–masala. Funksiyaning hosilasini toping.

$$y = \arcsin(e^{-2x}) + \ln(e^{2x} + \sqrt{e^{4x} - 1})$$

$$\begin{aligned}
y' &= \left(\arcsin(e^{-2x}) + \ln(e^{2x} + \sqrt{e^{4x} - 1}) \right)' = \\
&= \frac{1}{\sqrt{1 - (e^{-2x})^2}} + \frac{1}{e^{2x} + \sqrt{e^{4x} - 1}} \times 2e^{2x} + \frac{1}{2\sqrt{e^{4x} - 1}} \times e^{4x} = \\
&= \frac{1}{\sqrt{1 - e^{-4x}}} + \frac{1}{e^{2x} + \sqrt{e^{4x} - 1}} \times 2e^{2x} \times \frac{\sqrt{e^{4x} - 1}}{\sqrt{e^{4x} - 1}} + \frac{e^{2x}}{\sqrt{e^{4x} - 1}} = \\
&= \frac{e^{2x}}{\sqrt{e^{4x} - 1}} + \frac{2e^{2x}}{e^{2x} + \sqrt{e^{4x} - 1}} \times \frac{\sqrt{e^{4x} - 1} + e^{2x}}{\sqrt{e^{4x} - 1}} = \\
&= \frac{e^{2x}}{\sqrt{e^{4x} - 1}} + \frac{e^{2x}}{\sqrt{e^{4x} - 1}} \frac{3e^{2x}}{\sqrt{e^{4x} - 1}}.
\end{aligned}$$

$$1. y = \frac{1}{24} (x^2 + 8) \sqrt{x^2 - 4} + \frac{x^2}{16} \arcsin \frac{2}{x}, \quad x > 0.$$

$$2. y = \frac{4x+1}{16x^2+8x+3} + \frac{1}{\sqrt{2}} \operatorname{arctg} x \frac{4x+1}{\sqrt{2}}.$$

$$3. y = 2x - \ln(1 + \sqrt{1 - e^{4x}}) - e^{-2x} \arcsin(e^{2x}).$$

$$4. y = \sqrt{9x^2 - 12x + 5} \operatorname{arctg}(3x - 2) - \ln(3x - 2 + \sqrt{9x^2 - 12x + 5}).$$

$$5. y = \frac{2}{x-1} \sqrt{2x-x^2} + \ln \frac{1 + \sqrt{2x-x^2}}{x-1}.$$

$$6. y = \frac{x^4}{81} \arcsin \frac{3}{x} + \frac{1}{81} (x^2 + 18) \sqrt{x^2 - 9}, \quad x > 0.$$

$$7. y = \frac{1}{\sqrt{2}} \operatorname{arctg} \frac{3x-1}{\sqrt{2}} + \frac{1}{3} \times \frac{3x-1}{3x^2-2x+1}.$$

$$8. y = 3x - \ln(1 + \sqrt{1 - e^{6x}}) - e^{-3x} \arcsin(e^{3x}).$$

$$9. y = \ln(4x - 1 + \sqrt{16x^2 - 8x + 2}) - \sqrt{16x^2 - 8x + 2} \operatorname{arctg}(4x - 1).$$

$$10. y = \ln \frac{1 + 2\sqrt{-x-x^2}}{2x+1} + \frac{4}{2x+1} \sqrt{-x-x^2}.$$

$$11. y = (2x+3)^4 \times \arcsin \frac{1}{2x+3} + \frac{2}{3} \times (4x^2 + 12x + 11) \times \sqrt{x^2 + 3x + 2}, \quad 2x+3 > 0.$$

$$12. y = \frac{x+2}{x^2 + 4x + 6} + \frac{1}{\sqrt{2}} \times \operatorname{arctg} \frac{x+2}{\sqrt{2}}.$$

$$13. y = 5x - \ln(1 + \sqrt{1 - e^{10x}}) - e^{-5x} \times \arcsin(e^{5x}).$$

$$14. y = \sqrt{x^2 - 8x + 17} \times \operatorname{arctg}(x - 4) - \ln(x - 4 + \sqrt{x^2 - 8x + 17}).$$

$$15. y = \ln \frac{1 + \sqrt{-3 + 4x - x^2}}{2 - x} + \frac{2}{2 - x} \times \sqrt{-3 + 4x - x^2}.$$

$$16. y = (3x^2 - 4x + 2) \times \sqrt{9x^2 - 12x + 3} + (3x - 2)^4 \times \arcsin \frac{1}{3x - 2}, \quad 3x - 2 > 0.$$

$$17. y = \frac{1}{\sqrt{2}} \times \operatorname{arctg} \frac{x-1}{\sqrt{2}} + \frac{x-1}{x^2 - 2x + 3}.$$

$$18. y = \ln(e^{5x} + \sqrt{e^{10x} - 1}) + \arcsin(e^{-5x}).$$

$$19. y = \ln(2x - 3 + \sqrt{4x^2 - 12x + 10}) + \sqrt{4x^2 - 12x + 10} \times \operatorname{arctg}(2x - 3).$$

$$20. y = \ln \frac{1 + \sqrt{-3 - 4x - x^2}}{-x - 2} - \frac{2}{x + 2} \times \sqrt{-3 - 4x - x^2}.$$

$$21. y = \frac{2}{3} \times (4x^2 - 4x + 3) \times \sqrt{x^2 - x} + (2x - 1)^4 \times \arcsin \frac{1}{2x - 1}, \quad 2x - 1 > 0.$$

$$22. y = \frac{2x - 1}{4x^2 - 4x + 3} + \frac{1}{\sqrt{2}} \times \operatorname{arctg} \frac{2x - 1}{\sqrt{2}}.$$

$$23. y = \arcsin(e^{-4x}) + \ln(e^{4x} + \sqrt{e^{8x} - 1})$$

$$24. y = \ln(5x + \sqrt{25x^2 + 1}) - \sqrt{25x^2 + 1} \times \operatorname{arctg} 5x.$$

$$25. y = \frac{2}{3x - 2} \times \sqrt{-3 + 12x - 9x^2} + \ln \frac{1 + \sqrt{-3 + 12x - 9x^2}}{3x - 2}.$$

$$26. y = (3x + 1)^4 \times \arcsin \frac{1}{3x + 1} + (3x^2 + 2x + 1) \times \sqrt{9x^2 + 6x}, \quad 3x + 1 > 0.$$

$$27. y = \frac{1}{\sqrt{2}} \times \operatorname{arctg} \frac{2x+1}{\sqrt{2}} + \frac{2x+1}{4x^2 + 4x + 3}.$$

$$28. y = \ln(e^{3x} + \sqrt{e^{6x} - 1}) + \arcsin(e^{-3x}).$$

$$29. y = \sqrt{49x^2 + 1} \times \arctg 7x - \ln(7x + \sqrt{49x^2 + 1}).$$

$$30. y = \frac{1}{x} \times \sqrt{1 - 4x^2} + \ln \frac{1 + \sqrt{1 + 4x^2}}{2x}.$$

$$\text{Javoblar. 12.1 } \frac{x^3 - x}{8\sqrt{x^2 - 4}} + \frac{8}{x} \times \arcsin \frac{2}{x}; \quad 12.2 \frac{16}{(16x^2 + 8x + 3)^2};$$

$$12.3 \ 2e^{-2x} \times \arcsin(e^{2x}); 12.4 \ \frac{(9x - 6) \times \arctg(3x - 2)}{\sqrt{9x^2 - 12x + 5}}; \quad 12.5 \ \frac{2x^2 - 7x + 3}{(x - 1)^2 \sqrt{2x - x^2}};$$

$$12.6 \ \frac{4x^3}{81} \times \arcsin \frac{3}{x} + \frac{x \times (x^2 - 1)}{27\sqrt{x^2 - 9}}; \quad 12.7 \ \frac{4}{3(3x^2 - 2x + 1)^2}; \quad 12.8 \ 3e^{-3x} \times \arcsin(e^{3x});$$

$$12.9 \ \frac{4(1 - 4x)}{\sqrt{16x^2 - 8x + 2}} \times \arctg(4x - 1); \quad 12.10 \ - \frac{2x + 3}{\sqrt{-x - x^2} \times (2x + 1)^2};$$

$$12.11 \ 8(2x + 3)^3 \times \arcsin \frac{1}{2x + 3}; \quad 12.12 \ \frac{4}{(x^2 + 4x + 6)^2}; \quad 12.13 \ 5e^{-5x} \times \arcsin(e^{5x});$$

$$12.14 \ \frac{x - 4}{\sqrt{x^2 - 8x + 17}} \times \arctg(x - 4); \quad 12.15 \ \frac{4 - x}{(2 - x)^2 \times \sqrt{-3 + 4x - x^2}};$$

$$12.16 \ 12(3x - 2)^3 \times \arcsin \frac{1}{3x - 2}; \quad 12.17 \ \frac{4}{(x^2 - 2x + 3)^2}; \quad 12.18 \ 5e^{5x} \times \sqrt{1 - e^{-10x}};$$

$$12.19 \ \frac{4x - 6}{\sqrt{4x^2 - 12x + 10}} \times \arctg(2x - 3);$$

$$12.20 \ \frac{2x^2 + 8x + 9 + \sqrt{-3 - 4x - x^2}}{(x + 2) \times \sqrt{-3 - 4x - x^2} \times (1 + \sqrt{-3 - 4x - x^2})}; \quad 12.21 \ 8(2x - 1) \times \arcsin \frac{1}{2x - 1};$$

$$12.22 \ \frac{8}{(4x^2 - 4x + 3)^2}; \quad 12.23 \ 4\sqrt{\frac{e^{4x} - 1}{e^{4x} + 1}}; \quad 12.24 \ \frac{25x \times \arctg 5x}{\sqrt{25x^2 + 1}};$$

$$12.25 \ \frac{3 - 9x}{\sqrt{-3 + 12x - 9x^2} \times (3x - 2)};$$

$$12.26 \ 12(3x+1)^3 \times \arcsin \frac{1}{3x+1} + (3x+1) \times \frac{18x^2}{\sqrt{9x^2+6x}}; \quad 12.27 \ \frac{8}{(4x^2+4x+3)^2};$$

$$12.28 \ 3\sqrt{\frac{e^{3x}-1}{e^{3x}+1}}; \quad 12.29 \ \frac{7 \times \operatorname{arctg} 7x}{2\sqrt{49x^2+1}}; \quad 12.30 \ - \frac{1}{x^2\sqrt{1-4x^2}} - \frac{1}{x\sqrt{1+4x^2}}.$$

Funksiyaning hosilasi

13–masala. Funksiyaning hosilasini toping.

$$y = \frac{\arcsin x}{\sqrt{1-x^2}} + \frac{1}{2} \ln \frac{1-x}{1+x}.$$

$$\begin{aligned} y' &= \frac{x \times \arcsin x}{\sqrt{1-x^2}} + \frac{1}{2} \ln \frac{1-x}{1+x} = \frac{(x \times \arcsin x) \times \sqrt{1-x^2} - \arcsin x (\sqrt{1-x^2})}{1-x^2} + \\ &+ \frac{1}{2} \times \frac{1+x}{1-x} \times \frac{-1 \times (1+x) - (1-x) \times 1}{(1+x)^2} = \\ &= \frac{\frac{1}{\sqrt{1-x^2}} \times \sqrt{1-x^2} - \arcsin x \times \frac{1}{2\sqrt{1-x^2}} \times 2x}{1-x^2} + \\ &+ \frac{1}{2} \times \frac{1}{1-x} \times \frac{-2}{1+x} = \frac{1 - \frac{x \times \arcsin x}{\sqrt{1-x^2}}}{1-x^2} - \frac{1}{1-x^2} = \frac{x \times \arcsin x}{(1-x^2)\sqrt{1-x^2}}. \end{aligned}$$

$$1. \ y = \frac{x \times \arcsin x}{\sqrt{1-x^2}} + \ln \sqrt{1-x^2}.$$

$$2. \ y = 4 \ln \frac{x}{1+\sqrt{1-4x^2}} - \frac{\sqrt{1-4x^2}}{x^2}.$$

$$3. \ y = x(2x^2+5)\sqrt{x^2+1} + 3 \ln(x+\sqrt{x^2+1})$$

$$4. \ y = x^3 \arcsin x + \frac{x^2+2}{3} \sqrt{1-x^2}.$$

$$5. y = 3 \arcsin \frac{3}{4x+1} + 2\sqrt{4x^2 + 2x - 2}, \quad 4x+1 > 0.$$

$$6. y = \sqrt{1+x^2} \operatorname{arctg} x - \ln(x + \sqrt{1+x^2}).$$

$$7. y = 2 \arcsin \frac{2}{3x+4} + \sqrt{9x^2 + 24x + 12}, \quad 3x+4 > 0.$$

$$8. y = x(2x^2 + 1)\sqrt{x^2 + 1} - \ln(x + \sqrt{x^2 + 1})$$

$$9. y = \ln(x + \sqrt{x^2 + 1}) - \frac{\sqrt{x^2 + 1}}{x}.$$

$$10. y = \sqrt{1 - 3x - 2x^2} + \frac{3}{2\sqrt{2}} \arcsin \frac{4x+3}{\sqrt{17}}.$$

$$11. y = \sqrt{(4+x)(1+x)} + 3 \ln(\sqrt{4+x} + \sqrt{1+x}).$$

$$12. y = \ln \frac{\sqrt{x^2 - x + 1}}{x} + \sqrt{3} \operatorname{arctg} \frac{2x-1}{3}.$$

$$13. y = \frac{1}{12} \ln \frac{x^4 - x^2 + 1}{(x^2 + 1)^2} - \frac{1}{2\sqrt{3}} \operatorname{arctg} \frac{\sqrt{3}}{2x^2 - 1}.$$

$$14. y = 4 \arcsin \frac{4}{2x+3} + \sqrt{4x^2 + 12x - 7}, \quad 2x+3 > 0.$$

$$15. y = 2 \arcsin \frac{2}{3x+1} + \sqrt{9x^2 + 6x - 3}, \quad 3x+1 > 0.$$

$$16. y = (2+3x)\sqrt{x-1} - \frac{3}{2} \operatorname{arctg} \sqrt{x-1}.$$

$$17. y = \frac{1}{3}(x-2)\sqrt{x+1} + \ln(\sqrt{x+1} + 1).$$

$$18. y = \sqrt{x^2 + 1} - \frac{1}{2} \ln \frac{\sqrt{x^2 + 1} - x}{\sqrt{x^2 + 1} + 1}.$$

$$19. y = \sqrt[3]{\frac{x-1}{x+1}} - \frac{1}{2} \operatorname{arctg} \frac{x-1}{x^2-1} + \frac{1}{x^2-1} \operatorname{arctg} x.$$

$$20. y = x \ln(\sqrt{1-x} + \sqrt{1+x}) + \frac{1}{2}(\arcsin x - x).$$

$$21. y = \operatorname{arctg} \sqrt{x^2 - 1} - \frac{\ln x}{\sqrt{x^2 - 1}}.$$

$$22. y = 3 \arcsin \frac{3}{x+2} + \sqrt{x^2 + 4x - 5}, \quad x+2 > 0.$$

$$23. y = \sqrt{(3-x)(2+x)} + 5 \arcsin \sqrt{\frac{x+2}{5}}.$$

$$24. y = x(\arcsin x)^2 + 2\sqrt{1-x^2} \arcsin x - 2x.$$

$$25. y = \frac{\sqrt{1-x^2}}{x} + \arcsin x.$$

$$26. y = x^2 \arccos x - \frac{x^2 + 2}{3} \sqrt{1-x^2}.$$

$$27. y = \frac{\sqrt{x^2 + 2}}{x^2} - \frac{1}{\sqrt{2}} \ln \frac{\sqrt{2} + \sqrt{x^2 + 2}}{x}.$$

$$28. y = \frac{x}{4}(10 - x^2)\sqrt{4-x^2} + 6 \arcsin \frac{x}{2}.$$

$$29. y = \arcsin \frac{1}{2x+3} + 2\sqrt{x^2 + 3x + 2}, \quad 2x+3 > 0.$$

$$30. y = x \times \arcsin \sqrt{\frac{x}{x+1}} - \sqrt{x} + \operatorname{arctg} \sqrt{x}.$$

Javoblar. 13.1 $\frac{\arcsin x}{\sqrt{(1-x^2)^3}}$; 13.2 $\frac{2}{x^3 \sqrt{1-4x^2}}$; 13.3 $8\sqrt{(x^2+1)^3}$; 13.4 $3x^2 \arcsin x$;

13.5 $\frac{7 \times (4x+1)}{2\sqrt{4x^2+2x-2}}$; 13.6 $\frac{x \times \operatorname{arctg} x}{\sqrt{1+x^2}}$; 13.7 $\frac{8(3x+4)}{\sqrt{9x^2+24x+12}}$; 13.8 $8x^2 \sqrt{x^2+1}$;

13.9 $\frac{\sqrt{x^2+1}}{x^2}$; 13.10 $-\frac{2x}{\sqrt{1-3x-2x^2}}$; 13.11 $\sqrt{\frac{4+x}{1+x}}$; 13.12 $\frac{2x-1}{x \times (x^2-x+1)}$;

13.13 $\frac{x^3}{(x^4-x^2+1) \times (x^2+1)}$; 13.14 $\frac{2\sqrt{4x^2+12x-7}}{2x+3}$; 13.15 $\frac{3\sqrt{9x^2+6x-3}}{3x+1}$;

$$\begin{aligned}
& \mathbf{13.16} \frac{18x^2 - 8x - 3}{4x\sqrt{x-1}}; \mathbf{13.17} \frac{3x\sqrt{x+1} + 3x - \sqrt{x+1} + 2}{6\sqrt{x+1}(\sqrt{x+1} + 1)}; \mathbf{13.18} \frac{2\sqrt{x^2+1} + x + 2}{2(\sqrt{x^2+1} + 1)\sqrt{x^2+1}}; \\
& \mathbf{13.19} \frac{5x^2 + 17}{12(x^4 - 1)} + \frac{x \operatorname{arctg} x}{(x^2 - 1)^2}; \mathbf{13.20} \ln(\sqrt{1-x} + \sqrt{1+x}); \mathbf{13.21} \frac{x \ln x}{\sqrt{(x^2 - 1)^3}}; \mathbf{13.22} \\
& \frac{\sqrt{x^2 + 4x - 5}}{x + 2}; \mathbf{13.23} \sqrt{\frac{3-x}{2+x}}; \mathbf{13.24} (\arcsin x)^2; \mathbf{13.25} - \frac{\sqrt{1-x^2}}{x^2}; \mathbf{13.26} \\
& 2x \operatorname{arccos} x - x^2 \sqrt{\frac{1-x}{1+x}}; \mathbf{13.27} - \frac{4}{x^3 \sqrt{x^2 + 2}}; \mathbf{13.28} \sqrt{(4 - x^2)^3}; \mathbf{13.29} \\
& \frac{4\sqrt{x^2 + 3x + 2}}{2x + 3}; \mathbf{13.30} \arcsin \sqrt{\frac{x}{x+1}}.
\end{aligned}$$

Funksiyaning hosilasi

14-masala. Funksiyaning hosilasini toping.

$$\begin{aligned}
 y &= \sqrt{\frac{\operatorname{tg} x + \sqrt{2\operatorname{tg} x + 1}}{\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1}}}. \\
 y' &= \frac{\frac{\frac{1}{\cos^2 x} + \frac{1}{2\sqrt{2\operatorname{tg} x}} \times \frac{2}{\cos^2 x} \cdot \frac{\sqrt{2\operatorname{tg} x + 1}}{\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1}} - (\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1}) \cdot \frac{\frac{1}{\cos^2 x} + \frac{1}{2\sqrt{2\operatorname{tg} x}} \times \frac{2}{\cos^2 x} \cdot \frac{\sqrt{2\operatorname{tg} x + 1}}{\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1}}}{(\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1})^2}}{\frac{\sqrt{2\operatorname{tg} x + 1}}{\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1}}} \times \frac{(1 + \frac{1}{\sqrt{2\operatorname{tg} x}})(\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1}) - (\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1})(1 - \frac{1}{\sqrt{2\operatorname{tg} x}})}{\cos^2 x \times (\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1})^2}} = \\
 &= \frac{1}{2} \sqrt{\frac{\operatorname{tg} x + \sqrt{2\operatorname{tg} x + 1}}{\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1}}} \times \frac{\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1} + \frac{\sqrt{\operatorname{tg} x}}{\sqrt{2}} - 1 + \frac{1}{\sqrt{2\operatorname{tg} x}} - \operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1} + \frac{\sqrt{\operatorname{tg} x}}{\sqrt{2}} + 1 + \frac{1}{\sqrt{2\operatorname{tg} x}}}{\cos^2 x \times (\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1})^2} = \\
 &= \frac{1}{2} \sqrt{\frac{\operatorname{tg} x + \sqrt{2\operatorname{tg} x + 1}}{\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1}}} \times \frac{\sqrt{2\operatorname{tg} x} - 2\sqrt{2\operatorname{tg} x} + \frac{\sqrt{2}}{\sqrt{\operatorname{tg} x}}}{\cos^2 x \times (\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1})^2} = \\
 &= \frac{1}{2} \sqrt{\frac{\operatorname{tg} x + \sqrt{2\operatorname{tg} x + 1}}{\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1}}} \times \frac{\frac{\sqrt{2}}{\sqrt{\operatorname{tg} x}} - \sqrt{2\operatorname{tg} x}}{\cos^2 x \times (\operatorname{tg} x - \sqrt{2\operatorname{tg} x + 1})^2} =
 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{\sqrt{2}} \sqrt{\frac{\operatorname{tg} x - \sqrt{2\operatorname{tg} x} + 1}{(\operatorname{tg} x + \sqrt{2\operatorname{tg} x} + 1)(\operatorname{tg} x - \sqrt{2\operatorname{tg} x} + 1)^2}} \times \frac{\frac{1}{\sqrt{\operatorname{tg} x}} - \sqrt{\operatorname{tg} x}}{\cos^2 x \times (\operatorname{tg} x - \sqrt{2\operatorname{tg} x} + 1)} = \\
&= \frac{1}{\sqrt{2}} \sqrt{\frac{1}{\operatorname{tg}^2 x + 1}} \times \frac{\frac{1}{\sqrt{\operatorname{tg} x}} - \sqrt{\operatorname{tg} x}}{\cos^2 x \times (\operatorname{tg} x - \sqrt{2\operatorname{tg} x} + 1)} = \\
&= \frac{1}{\sqrt{2}} \cos x \times \frac{\frac{1}{\sqrt{\operatorname{tg} x}} - \sqrt{\operatorname{tg} x}}{\cos^2 x \times (\operatorname{tg} x - \sqrt{2\operatorname{tg} x} + 1)} = \\
&= \frac{1 - \operatorname{tg} x}{\sqrt{2\operatorname{tg} x} \times (\sin x - \sqrt{\sin 2x} + \cos x)}.
\end{aligned}$$

$$1. y = \frac{1}{\sin a} \ln(\operatorname{tg} x + \operatorname{ctg} a).$$

$$2. y = x > \cos a + \sin a > \ln \sin(x - a).$$

$$3. y = \frac{1}{2\sqrt{2}} (\sin(\ln x) - (\sqrt{2} - 1) \times \cos(\ln x)) x^{\sqrt{2}+1}.$$

$$4. y = \operatorname{arctg} \frac{\cos x}{\sqrt[4]{\cos 2x}}.$$

$$5. y = 3 \frac{\sin x}{\cos^2 x} + 2 \frac{\sin x}{\cos^4 x}.$$

$$6. y = (a^2 + b^2)^{-\frac{1}{2}} \times \arcsin \frac{\sqrt{a^2 + b^2} \times \sin x}{b}.$$

$$7. y = \frac{7^x (3 \sin 3x + \cos 3x \times \ln 7)}{9 + \ln^2 7}.$$

$$8. y = \ln \frac{\sin x}{\cos x + \sqrt{\cos 2x}}.$$

$$9. y = \frac{1}{a(1 + a^2)} \left(\operatorname{arctg}(a \cos x) + a \ln \left(\operatorname{tg} \frac{x}{2} \right) \right).$$

$$10. y = -\frac{1}{3 \sin^3 x} - \frac{1}{\sin x} + \frac{1}{2} \ln \frac{1 + \sin x}{1 - \sin x}.$$

$$11. y = (1 + x^2)e^{\operatorname{arctg} x}.$$

$$12. y = \frac{\operatorname{ctg} x + x}{1 - x \times \operatorname{ctg} x}.$$

$$13. y = \frac{1}{2 \sin \frac{a}{2}} \times \operatorname{arctg} \frac{2x \sin \frac{a}{2}}{1 - x^2}.$$

$$14. y = \operatorname{arctg} \frac{\sqrt{\sqrt{x^4 + 1} - x^2}}{x}, \quad x > 0.$$

$$15. y = \frac{6^x (\sin 4x \times \ln 6 - 4 \cos 4x)}{16 + \ln^2 6}.$$

$$16. y = \operatorname{arctg} \frac{\sqrt{2 \operatorname{tg} x}}{1 - \operatorname{tg} x}.$$

$$17. y = \operatorname{arctg} \frac{2 \sin x}{\sqrt{9 \cos^2 x - 4}}.$$

$$18. y = \frac{5^x (2 \sin 2x + \cos 2x \times \ln 5)}{4 + \ln^2 5}.$$

$$19. y = \ln \frac{\sqrt{2} + \operatorname{th} x}{\sqrt{2} - \operatorname{th} x}.$$

$$20. y = \frac{3^x (4 \sin 4x + \ln 3 \times \cos 4x)}{16 + \ln^2 3}.$$

$$21. y = \frac{4^x (\ln 4 \times \sin 4x - 4 \cos 4x)}{16 + \ln^2 4}.$$

$$22. y = \frac{\cos x}{\sin^2 x} - 2 \cos x - 3 \ln \left(\operatorname{tg} \frac{x}{2} \right).$$

$$23. y = \frac{5^x (\sin 3x \times \ln 5 - 3 \cos 3x)}{9 + \ln^2 5}.$$

$$24. y = x - \ln(1 + e^x) - 2e^{-\frac{x}{2}} \operatorname{arctg} e^{\frac{x}{2}}.$$

$$25. y = \frac{2^x (\sin x + \cos x \times \ln 2)}{1 + \ln^2 2}.$$

$$26. y = \frac{\ln(\operatorname{ctg} x + \operatorname{ctg} a)}{\sin a}.$$

$$27. y = 2 \frac{\cos x}{\sin^4 x} + 3 \frac{\cos x}{\sin^2 x}.$$

$$28. y = \frac{\cos x}{3(2 + \sin x)} + \frac{4}{3\sqrt{3}} \operatorname{arctg} \frac{2\operatorname{tg}\left(\frac{x}{2}\right) + 1}{\sqrt{3}}.$$

$$29. y = \frac{3^x (\ln 3 \times \sin 2x - 2 \cos 2x)}{\ln^2 3 + 4}.$$

$$30. y = \frac{1}{2} \ln \frac{1 + \cos x}{1 - \cos x} - \frac{1}{\cos x} - \frac{1}{3 \cos^3 x}.$$

Javoblar. 14.1 $\frac{1}{\cos x \times \cos(a - x)}$; 14.2 $\frac{\sin x}{\sin(x - a)}$;

14.4 $-\frac{\sin^3 x}{(\sqrt{\cos 2x + \cos^2 x}) \times \sqrt[4]{(\cos 2x)^3}};$

14.5 $\frac{3 + 3 \sin^2 x}{\cos^3 x} + \frac{2 - 6 \sin^2 x}{\cos^5 x}$; 14.6 $\frac{\cos x}{\sqrt{b^2 \times \cos^2 x - a^2 \times \sin^2 x}}$; 14.7 $7^x > \cos 3x$;

14.8 $\frac{1}{\sin x \times \sqrt{\cos 2x}}$; 14.9 $\frac{\cos x > \operatorname{ctg} x}{1 + a^2 > \cos^2 x}$; 14.10 $\frac{1}{\cos x > \sin^4 x}$; 14.11 $(2x + 1) \times e^{\operatorname{arctg} x}$;

14.12 $-\frac{x^2}{(\sin x - x \times \cos x)^2}$; 14.13 $\frac{1 + x^2}{(1 - x^2)^2 + 4x^2 \times \sin^2 \frac{a}{2}}$;

14.14 $-\frac{1}{(x^4 + 1) \times \sqrt{\sqrt{x^4 + 1} - x^2}}$; 14.15 $6^x \sin 4x$; 14.16 $\frac{1 - \operatorname{tg} x + \sqrt{2 \operatorname{tg} x}}{\sqrt{2 \operatorname{tg} x}}$;

14.17 $\frac{2}{\cos x \sqrt{9 \cos^2 x - 4}}$; 14.18 $5^x > \cos 2x$; 14.19 $\frac{2\sqrt{2}}{\operatorname{ch}^2 x + 1}$; 14.20 $3^x > \cos 4x$;

14.21 $4^x \sin 4x$; 14.22 $-\frac{2 + 3 \sin^2 x}{\sin^3 x}$; 14.23 $5^x \sin 3x$; 14.24 $x \times e^{-\frac{x}{2}} \times \operatorname{arctg} e^{\frac{x}{2}}$;

14.25 $2^x > \cos x$; 14.26 $-\frac{1}{\sin(a + x)}$; 14.27 $3 \operatorname{cosec} x - 8 \operatorname{cosec}^5 x$; 14.28 $\frac{2 \sin x + 7}{3(2 + \sin x)^2}$;

14.29 $3^x \sin 2x$; 14.30 $-\frac{1}{\sin x > \cos^4 x}$.

Parametrik berilgan funksiyalarning hosilalari

Agar x ning funksiyasi y ushbu

$$\begin{cases} x = x(t), \\ y = y(t) \end{cases}$$

parametrik tenglamalar bilan berilgan bo'lsa, u holda y ning x bo'yicha hosilasi y'_x

$$y'_x = \frac{y'_t}{x'_t}$$

tenglik bilan aniqlanadi.

15-masala. y'_x funksiyaning hosilasini toping.

$$\begin{cases} x = \ln(t + \sqrt{1+t^2}) \\ y = \sqrt{1+t^2} - \ln \frac{1+\sqrt{1+t^2}}{t} \end{cases}$$

$$\begin{aligned} y'_x &= \left(\ln(t + \sqrt{1+t^2}) \right)' = \frac{1}{t + \sqrt{1+t^2}} \times 1 + \frac{1}{2\sqrt{1+t^2}} \times 2t = \\ &= \frac{1}{t + \sqrt{1+t^2}} \times \frac{\sqrt{1+t^2} + t}{\sqrt{1+t^2}} = \frac{1}{\sqrt{1+t^2}}. \end{aligned}$$

$$\begin{aligned}
 y' &= \left(\sqrt{1+t^2} - \ln \frac{1+\sqrt{1+t^2}}{t} \right)' = \frac{1}{2\sqrt{1+t^2}} \times 2t - \frac{1}{1+\sqrt{1+t^2}} \times \frac{1+\sqrt{1+t^2}}{t} = \\
 &= \frac{t}{\sqrt{1+t^2}} - \frac{t}{1+\sqrt{1+t^2}} \times \frac{\frac{1}{2\sqrt{1+t^2}} \times 2t \times (1+\sqrt{1+t^2})}{t^2} = \\
 &= \frac{t}{\sqrt{1+t^2}} - \frac{t}{1+\sqrt{1+t^2}} \times \frac{t^2 - \sqrt{1+t^2} - 1 - t^2}{t^2 \sqrt{1+t^2}} = \\
 &= \frac{t}{\sqrt{1+t^2}} + \frac{1}{1+\sqrt{1+t^2}} \times \frac{\sqrt{1+t^2} + 1}{t\sqrt{1+t^2}} = \frac{t^2 + 1}{t\sqrt{1+t^2}}.
 \end{aligned}$$

Natijada:

$$y' = \frac{y'}{x'} = \frac{1}{\sqrt{1+t^2}} \cdot \frac{t^2+1}{t\sqrt{1+t^2}} = \frac{t}{t^2+1}.$$

$$1. \begin{cases} x = \frac{3t^2+1}{t^3} \\ y = \sin\left(\frac{t^3}{3} + t\right) \end{cases}$$

$$2. \begin{cases} x = \sqrt{1-t^2} \\ y = \operatorname{tg} \sqrt{1+t} \end{cases}$$

$$3. \begin{cases} x = \sqrt{2t-t^2} \\ y = \frac{1}{\sqrt[3]{(1-t)^2}} \end{cases}$$

$$4. \begin{cases} x = \arcsin(\sin t) \\ y = \arccos(\cos t) \end{cases}$$

$$5. \begin{cases} x = \ln(t + \sqrt{t^2+1}) \\ y = t\sqrt{t^2+1} \end{cases}$$

$$6. \begin{cases} x = \sqrt{2t-t^2} \\ y = \arcsin(t-1) \end{cases}$$

$$7. \begin{cases} x = \operatorname{ctg}(2e^t) \\ y = \ln(\operatorname{tg} e^t) \end{cases}$$

$$8. \begin{cases} x = \ln(\operatorname{ctg} t) \\ y = \frac{1}{\cos^2 t} \end{cases}$$

$$9. \begin{cases} x = \operatorname{arctg} e^{\frac{t}{2}} \\ y = \sqrt{e^t+1} \end{cases}$$

$$10. \begin{cases} x = \ln \sqrt{\frac{1-t}{1+t}} \\ y = \sqrt{1-t^2} \end{cases}$$

$$11. \begin{cases} \dot{x} = \ln \frac{1}{\sqrt{1-t^4}} \\ \ddot{y} = \arcsin \frac{1-t^2}{1+t^2} \end{cases}$$

$$13. \begin{cases} \dot{x} = \arcsin(\sqrt{1-t^2}) \\ \dot{y} = (\arccos t)^2 \end{cases}$$

$$15. \begin{cases} \dot{x} = (1 + \cos^2 t)^2 \\ \ddot{y} = \frac{\cos t}{\sin^2 t} \end{cases}$$

$$17. \begin{cases} \dot{x} = \arccos \frac{1}{t} \\ \ddot{y} = \sqrt{1-t^2} + \arcsin \frac{1}{t} \end{cases}$$

$$19. \begin{cases} \dot{x} = \arcsin \sqrt{t} \\ \dot{y} = \sqrt{1+\sqrt{t}} \end{cases}$$

$$21. \begin{cases} \dot{x} = t\sqrt{t^2+1} \\ \ddot{y} = \ln \frac{1+\sqrt{1+t^2}}{t} \end{cases}$$

$$23. \begin{cases} \dot{x} = \ln(1-t^2) \\ \dot{y} = \arcsin \sqrt{1-t^2} \end{cases}$$

$$25. \begin{cases} \dot{x} = \ln \sqrt{\frac{1-\sin t}{1+\sin t}} \\ \ddot{y} = \frac{1}{2} \operatorname{tg}^2 t + \ln \cos t \end{cases}$$

$$27. \begin{cases} \dot{x} = \ln(\operatorname{tg} t) \\ \ddot{y} = \frac{1}{\sin^2 t} \end{cases}$$

$$29. \begin{cases} \dot{x} = e^{\sec^2 t} \\ \dot{y} = \operatorname{tg} t \times \ln \cos t + \operatorname{tg} t - t \end{cases}$$

$$12. \begin{cases} \dot{x} = \sqrt{1-t^2} \\ \ddot{y} = \frac{t}{\sqrt{1-t^2}} \end{cases}$$

$$14. \begin{cases} \dot{x} = \frac{t}{\sqrt{1-t^2}} \\ \ddot{y} = \ln \frac{1+\sqrt{1-t^2}}{t} \end{cases}$$

$$16. \begin{cases} \dot{x} = \ln \frac{1-t}{1+t} \\ \ddot{y} = \sqrt{1-t^2} \end{cases}$$

$$18. \begin{cases} \dot{x} = \frac{1}{\ln t} \\ \ddot{\ln} \frac{1+\sqrt{1-t^2}}{t} \end{cases}$$

$$20. \begin{cases} \dot{x} = (\arcsin t)^2 \\ \dot{y} = \frac{t}{\sqrt{1-t^2}} \end{cases}$$

$$22. \begin{cases} \dot{x} = \operatorname{arctg} t \\ \dot{y} = \ln \frac{\sqrt{1+t^2}}{t+1} \end{cases}$$

$$24. \begin{cases} \dot{x} = \operatorname{arctg} \frac{t+1}{t-1} \\ \dot{y} = \arcsin \sqrt{1-t^2} \end{cases}$$

$$26. \begin{cases} \dot{x} = \sqrt{t-t^2} - \operatorname{arctg} \sqrt{\frac{1-t}{t}} \\ \dot{y} = \sqrt{t} - \sqrt{1-t} \times \arcsin \sqrt{t} \end{cases}$$

$$28. \begin{cases} \dot{x} = \frac{t^2 \ln t}{1-t^2} + \ln \sqrt{1-t^2} \\ \dot{y} = \frac{t}{\sqrt{1-t^2}} \arcsin t + \ln \sqrt{1-t^2} \end{cases}$$

$$30. \begin{cases} \dot{x} = \frac{t}{\sqrt{1-t^2}} \times \arcsin t + \ln \sqrt{1-t^2} \\ \dot{y} = \frac{t}{\sqrt{1-t^2}} \end{cases}$$

Javoblar. 15.1 $-t^4 \times \cos \frac{\pi}{3} + t^{\frac{3}{2}}$; 15.2 $\frac{\sqrt{1-t}}{2t \times \cos^2(\sqrt{1+t})}$; 15.3 $\frac{2\sqrt{2t-t^2}}{3(1-t^2) \times \sqrt{(1-t)^2}}$;
 15.4 1; 15.5 $2t^2 + 1$; 15.6 $\frac{1}{(1-t)}$; 15.7 $-\sin(2e^t)$; 15.8 $-2tg^2t$; 15.9 $\sqrt{e^{2t} + e^t}$;
 15.10 $t \times \sqrt{1-t^2}$; 15.11 $\frac{t^2-1}{t^3}$; 15.12 $\frac{1}{t \times (t^2-1)}$; 15.13 $\frac{2 \arccos t \times \sqrt{1-t^2}}{\sqrt{1-t^2}}$; 15.14 $\frac{t^2-1}{t}$;
 15.15 $\frac{1}{4 \sin^4 t \times \cos t}$; 15.16 $\frac{t \times \sqrt{1-t^2}}{2}$; 15.17 $t^2 - 1$; 15.18 $\frac{\ln^2 t}{\sqrt{1-t^2}}$; 15.19 $2\sqrt{\frac{1-t}{1+\sqrt{t}}}$;
 15.20 $\frac{1}{(1-t^2) \times 2 \arcsin t}$; 15.21 $-\frac{1}{t \times (2t^2+1)}$; 15.22 $\frac{t-1}{t+1}$; 15.23 $\frac{\sqrt{1-t^2}}{2t}$;
 15.24 $\frac{t^2+1}{\sqrt{1-t^2}}$; 15.25 $\frac{\sin t \times \cos t - 1}{\cos t}$; 15.26 $\frac{\sqrt{t} \times \arcsin \sqrt{t}}{2 \times (1-t)}$; 15.27 $-2ctg^2t$;
 15.28 $\frac{\arcsin t \times \sqrt{1-t^2}}{2t \times \ln t}$; 15.29 $\frac{1}{2} \times t g t \times \ln \cos t \times e^{-\sec^2 t}$; 15.30 $\frac{1}{\arcsin t}$.

Egri chiziqqa nuqtadan o'tkazilgan urinma va normal tenglamasi

$y = f(x)$ egri chiziqning $M_0(x_0, f(x_0))$ nuqtasidan o'tkazilgan urinma tenglamasi:

$$y - f(x_0) = f'(x_0)(x - x_0)$$

$y = f(x)$ egri chiziqning $M_0(x_0, f(x_0))$ nuqtasidan o'tkazilgan normal(perpendikulyar) tenglamasi:

$$y - y_0 = -\frac{1}{f'(x_0)}(x - x_0) \quad (f'(x_0) \neq 0).$$

16-masala. Funksiya grafigining $t = t_0$ parametrning qiymatiga mos kelgan egri chiziqning nuqtasiga o'tkazilgan urinma va normal tenglamasini tuzing.

$$\begin{cases} x = 2e^t \\ y = e^{-t}, \quad t_0 = 0 \end{cases}$$

Echim:

$t_0 = 0$ ekanligidan, u holda

$$x_0 = 2e^0 = 2$$

$$y_0 = e^{-0} = 1$$

Hosilalarni topamiz:

$$x_t' = (2e^t)' = 2e^t$$

$$y_t' = (e^{-t})' = -e^{-t}$$

$$y_x' = \frac{y_t'}{x_t'} = \frac{-e^{-t}}{2e^t} = -\frac{1}{2e^{2t}}$$

u holda

$$y_x' = -\frac{1}{2e^{2x_0}} = -\frac{1}{2}.$$

urinma tenglamasi:

$$y - y_0 = y_x'(x - x_0)$$

$$y - 1 = -\frac{1}{2}(x - 2)$$

$$y = -\frac{1}{2}x + 2$$

normal tenglamasi:

$$y - y_0 = -\frac{1}{y_x'}(x - x_0)$$

$$y - 1 = -\frac{1}{(-\frac{1}{2})}(x - 2)$$

$$y = 2x - 3.$$

$$1. \begin{cases} \dot{x} = a \sin^3 t \\ \dot{y} = a \cos^3 t, \quad t_0 = \frac{\rho}{3} \end{cases}$$

$$2. \begin{cases} \dot{x} = \sqrt{3} \times \cos t \\ \dot{y} = \sin t, \quad t_0 = \frac{\rho}{3} \end{cases}$$

$$3. \begin{cases} \dot{x} = a(t - \sin t) \\ \dot{y} = a(1 - \cos t), \quad t_0 = \frac{\rho}{3} \end{cases}$$

$$4. \begin{cases} \dot{x} = 2t - t^2 \\ \dot{y} = 3t - t^3, \quad t_0 = 1 \end{cases}$$

$$5. \begin{cases} \dot{x} = \frac{2t+t^2}{1+t^3} \\ \dot{y} = \frac{2t-t^2}{1+t^3}, \quad t_0 = 1 \end{cases}$$

$$6. \begin{cases} \dot{x} = \arcsin \frac{t}{\sqrt{1+t^2}} \\ \dot{y} = \arccos \frac{t}{\sqrt{1+t^2}}, \quad t_0 = -1 \end{cases}$$

$$7. \begin{cases} \dot{x} = t(t \times \cos t - 2 \sin t) \\ \dot{y} = t(t \times \sin t + 2 \cos t), \quad t_0 = \frac{\rho}{4} \end{cases}$$

$$8. \begin{cases} \dot{x} = \frac{3at}{1+t^2} \\ \dot{y} = \frac{3at^2}{1+t^2}, \quad t_0 = 2 \end{cases}$$

$$9. \begin{cases} \dot{x} = 2 \ln(\operatorname{ctgt}) + \operatorname{ctgt} \\ \dot{y} = \operatorname{tgt} + \operatorname{ctgt}, \quad t_0 = \frac{\rho}{4} \end{cases}$$

$$10. \begin{cases} \dot{x} = \frac{1}{2} \times^2 - \frac{1}{4} \times^4 \\ \dot{y} = \frac{1}{2} \times^2 + \frac{1}{3} \times^3, \quad t_0 = 0 \end{cases}$$

$$11. \begin{cases} \dot{x} = a \times t \times \cos t \\ \dot{y} = a \times \sin t, \quad t_0 = \frac{\rho}{2} \end{cases}$$

$$12. \begin{cases} \dot{x} = \sin t \\ \dot{y} = \cos t, \quad t_0 = \frac{\rho}{6} \end{cases}$$

$$13. \begin{cases} \dot{x} = \arcsin \frac{t}{\sqrt{1+t^2}} \\ \dot{y} = \arccos \frac{1}{\sqrt{1+t^2}}, \quad t_0 = 1 \end{cases}$$

$$14. \begin{cases} \dot{x} = \frac{1+\ln t}{t^2} \\ \dot{y} = \frac{3+2\ln t}{t}, \quad t_0 = 1 \end{cases}$$

$$15. \begin{cases} \dot{x} = \frac{1+t}{t^2} \\ \dot{y} = \frac{3}{2t^2} + \frac{2}{t}, \quad t_0 = 2 \end{cases}$$

$$16. \begin{cases} \dot{x} = a \times \sin^3 t \\ \dot{y} = a \times \cos^3 t, \quad t_0 = \frac{\rho}{6} \end{cases}$$

$$17. \begin{cases} \dot{x} = a(t \times \sin t + \cos t) \\ \dot{y} = a(\sin t - t \times \cos t), \quad t_0 = \frac{\rho}{4} \end{cases}$$

$$18. \begin{cases} \dot{x} = \frac{t+1}{t} \\ \dot{y} = \frac{t-1}{t}, \quad t_0 = -1 \end{cases}$$

$$19. \begin{cases} \dot{x} = 1 - t^2 \\ \dot{y} = 1 - t^3, \quad t_0 = 2 \end{cases}$$

$$20. \begin{cases} \dot{x} = \ln(1+t^2) \\ \dot{y} = t - \operatorname{arctgt}, \quad t_0 = 1 \end{cases}$$

$$21. \begin{cases} \dot{x} = t(1 - \sin t) \\ \dot{y} = t \times \cos t, \quad t_0 = 0 \end{cases}$$

$$22. \begin{cases} \dot{x} = \frac{1+t^3}{t^2-1} \\ \dot{y} = \frac{t}{t^2-1}, \quad t_0 = 2 \end{cases}$$

$$23. \begin{cases} \dot{x} = 3 \cos t \\ \dot{y} = 4 \sin t, \quad t_0 = \frac{\rho}{4} \end{cases}$$

$$24. \begin{cases} \dot{x} = t - t^4 \\ \dot{y} = t^2 - t^3, \quad t_0 = 1 \end{cases}$$

$$25. \begin{cases} \dot{x} = t^3 + 1 \\ \dot{y} = t^2 + t + 1, \quad t_0 = 1 \end{cases}$$

$$26. \begin{cases} \dot{x} = 2 \cos t \\ \dot{y} = \sin t, \quad t_0 = -\frac{\rho}{3} \end{cases}$$

$$27. \begin{cases} \dot{x} = 2 \operatorname{tg} t \\ \dot{y} = 2 \sin^2 t + \sin 2t, \quad t_0 = \frac{\rho}{4} \end{cases}$$

$$28. \begin{cases} \dot{x} = t^3 + 1 \\ \dot{y} = t^2, \quad t_0 = -2 \end{cases}$$

$$29. \begin{cases} \dot{x} = \sin t \\ \dot{y} = a^t, \quad t_0 = 0 \end{cases}$$

$$30. \begin{cases} \dot{x} = \sin t \\ \dot{y} = \cos 2t, \quad t_0 = \frac{\rho}{6} \end{cases}$$

Javoblar. 16.1 $y = -\frac{x}{\sqrt{3}} + \frac{a}{2}$; $y = \sqrt{3}x - a$; **16.2** $y = 3x - \sqrt{3}$; **16.3** $y = -\frac{x}{\sqrt{3}} + \frac{a\rho}{3\sqrt{3}}$;

16.4 $y = 3x - 1$; $y = -\frac{x}{3} + 2\frac{1}{3}$; **16.5** $y = 3x - 4$; $y = -\frac{x}{3} + 1$; **16.6** $y = 2x + \frac{3\rho}{4}$;

$y = -\frac{x}{2} + \frac{\rho}{8}$; **16.7** $y = -x + \frac{\rho^2 \times \sqrt{2}}{16}$; $y = x + \frac{\rho \times \sqrt{2}}{2}$; **16.8** $y = -\frac{4}{3}x - 4a$;

$y = \frac{3}{4}x + \frac{3a}{2}$; **16.9** $y = 2$; $x = 1$; **16.10** $y = x$; $y = -x$; **16.11** $y = -\frac{2x}{\rho} + \frac{a\rho}{2}$;

$y = -\frac{\rho \times x}{2} + \frac{a\rho}{2}$; **16.12** $y = -\frac{1}{\sqrt{3}}x + \frac{2}{\sqrt{3}}$; $y = \sqrt{3}x$; **16.13** $y = 2x - \frac{\rho}{4}$; $y = -\frac{x}{2} + \frac{3\rho}{8}$;

16.14 $y = x + 2$; $y = -x + 4$; **16.15** $y = \frac{7x}{4} + 2\frac{17}{48}$; $y = -\frac{4x}{7} + 4\frac{2}{21}$; **16.16**

$y = -\sqrt{3}x + \frac{4\sqrt{3}a}{8}$; $y = \frac{x}{\sqrt{3}} + \frac{a}{\sqrt{3}}$; **16.17** $y = x + \frac{\sqrt{2} \times a \times \rho}{4}$; $y = -x + \sqrt{2} \times a$;

16.18 $y = -x + 2$; $y = x + 2$; **16.19** $y = \frac{11x}{4} + \frac{9}{4}$; $y = -\frac{4x}{11} - \frac{78}{11}$;

$$16.20 \quad y = \frac{x}{2} + \frac{4 - p - \ln 4}{4}; \quad y = -2x + \frac{4 - p + \ln 16}{2}; \quad 16.21 \quad y = 0; \quad x = 0;$$

$$16.22 \quad x = 3; \quad y = \frac{2}{3}; \quad 16.23 \quad y = -\frac{4}{3}x + 4\sqrt{2}; \quad y = \frac{3}{4}x + \frac{7\sqrt{2}}{8}; \quad 16.24 \quad y = \frac{x}{3}; \quad y = -3x;$$

$$16.25 \quad y = x + 1; \quad y = -x + 5; \quad 16.26 \quad y = \frac{\sqrt{3}}{6}x - \frac{2\sqrt{3}}{3}; \quad y = -2\sqrt{3}x + \frac{3\sqrt{3}}{2};$$

$$16.27 \quad y = -2x + 6; \quad y = \frac{1}{2}x + 1; \quad 16.28 \quad y = -\frac{1}{3}x + 7; \quad y = 3x - 23;$$

$$16.29 \quad y = x \ln a + 1; \quad y = -\frac{x}{\ln a} + 1; \quad 16.30 \quad y = -2x + 1,5; \quad y = \frac{1}{2}x + 0,25.$$

Yuqori tartibli hosilalar

Birinchi tartibli hosiladan olingan hosila, ya'ni

$$(y)' = (f'(x))' \quad \text{yoki} \quad y'' = f''(x)$$

$y = f(x)$ funksiyaning ikkinchi tartibli hosilasi deyiladi.

Ikkinchi tartibli hosilaning hosilasiga uchinchi tartibli hosila deyiladi

va $y''' = f'''(x)$, $\frac{d^3 y}{dx^3}$ belgilarning biri bilan belgilanadi.

Umuman, $y = f(x)$ funksiyaning n - tartibli hosilasi deb, uning

$(n - 1)$ - tartibli hosilasining hosilasiga aytiladi va $y^{(n)}$, $f^{(n)}(x)$, $\frac{d^n y}{dx^n}$

belgilarning biri bilan belgilanadi.

17-masala. Funksiyaning n -tartibli hosilasini toping.

$$y = 3^{2x+5}.$$

Echim:

$$y' = (3^{2x+5})' = 3^{2x+5} \ln 3 \cdot 2.$$

$$y'' = (3^{2x+5} \ln 3 \cdot 2)' = 3^{2x+5} \ln^2 3 \cdot 2^2.$$

Shunday qilib,

$$y^{(n)} = 3^{2x+5} \ln^n 3 \cdot 2^n = 2^n \ln^n 3 \cdot 3^{2x+5}.$$

$$1. y = x e^{ax}.$$

$$2. y = \sin 2x + \cos(x+1).$$

$$3. y = \sqrt[5]{e^{7x-1}}.$$

$$4. y = \frac{4x+7}{2x+3}.$$

$$5. y = \lg(5x+2).$$

$$6. y = a^{3x}.$$

$$7. y = \frac{x}{2(3x+2)}.$$

$$8. y = \lg(x+4).$$

$$9. y = \sqrt{x}.$$

$$10. y = \frac{2x+5}{13(3x+1)}.$$

$$11. y = 2^{3x+5}.$$

$$12. y = \sin(x+1) + \cos 2x.$$

$$13. y = \sqrt[3]{e^{2x+1}}.$$

$$14. y = \frac{4+15x}{5x+1}.$$

$$15. y = \lg(3x+1).$$

$$16. y = 7^{5x}.$$

$$17. y = \frac{x}{9(4x+9)}.$$

$$18. y = \lg(1+x).$$

$$19. y = \frac{4}{x}.$$

$$20. y = \frac{5x+1}{13(2x+3)}.$$

$$21. y = a^{2x+3}.$$

$$22. y = \sin(3x+1) + \cos 5x.$$

$$23. y = \sqrt{e^{3x+1}}.$$

$$24. y = \frac{11+12x}{6x+5}.$$

$$25. y = \lg(2x+7).$$

$$26. y = 2^{kx}.$$

$$27. y = \frac{x}{x+1}.$$

$$28. y = \log_3(x+5).$$

$$29. y = \frac{1+x}{1-x}.$$

$$30. y = \frac{7x+1}{17(4x+3)}.$$

Javoblar. 17.1 $(n+ax)e^{ax} \cdot a^{n-1}$; 17.2 $2^n \sin \frac{\pi}{2} n + 2x \frac{\pi}{2} + \cos \frac{\pi}{2} n + x + 1 \frac{\pi}{2}$; 17.3

$$17.7 \frac{(-1)^{n-1} n! 3^{n-1}}{(3x+2)^{n+1}}; 17.8 \frac{(-1)^{n-1} (n-1)!}{\ln 10 (x+4)^n}; 17.9 \frac{(-1)^{n-1} \prod_{k=1}^{n-1} (2k-1)}{2^{(n+1)} x^{n-1} \sqrt{x}};$$

$$17.10 \frac{(-1)^n n! 3^{n-1}}{(3x+1)^{n+1}}; 17.11 2^{3x+5} > 3^n > \ln^n 2;$$

$$17.12 \sin \frac{\arcsin p}{2} \times n + x + 1 \frac{\ddot{0}}{\emptyset} + 2^n \times \cos \frac{\arcsin p}{2} \times n + 2x \frac{\ddot{0}}{\emptyset};$$

$$17.13 \frac{\arcsin \ddot{0}}{\emptyset} \times \sqrt[3]{e^{2x+1}}; 17.14 \frac{(-1)^n n! 5^n}{(5x+1)^{n+1}}; 17.15 \frac{(-1)^{n-1} (n-1)! 3^n}{\ln 10 (3x+1)^n}; 17.16 5^n \ln(7)^n \times 7^{5x};$$

$$17.17 \frac{(-1)^{n-1} n! 4^{n-1}}{(4x+9)^{n+1}}; 17.18 \frac{(-1)^{n-1} (n-1)!}{\ln 10 (1+x)^n}; 17.19 \frac{4 \times (-1)^n n!}{x^n}; 17.20 \frac{(-1)^{n-1} n! 2^{n-1}}{(2x+3)^{n+1}};$$

$$17.21 a^{2x+3} > 2^n > \ln^n a; 17.22 3^n \times \sin \frac{\arcsin p}{2} \times n + 3x + 1 \frac{\ddot{0}}{\emptyset} + 5^n \times \cos \frac{\arcsin p}{2} \times n + 5x \frac{\ddot{0}}{\emptyset};$$

$$17.23 \frac{\arcsin \ddot{0}}{\emptyset} \times \sqrt{e^{3x+1}}; 17.24 (-1)^n n! 6^n \times (6x+5)^{-n-1};$$

$$17.25 (-1)^{n-1} \times \frac{2^n (n-1)!}{\ln 10} \times (2x+7)^{-n}; 17.26 2^{kx} > k^n \ln^n 2; 17.27 (-1)^{k+2} \times \frac{(k+1)!}{(x+1)^{k+2}}; 17.28$$

$$\frac{(-1)^{n-1} (n-1)!}{\ln 3 (x+5)^n}; 17.29 \frac{2 > (k+1)!}{(1-x)^{k+2}}; 17.30 \frac{(-1)^{n-1} n! 4^{n-1}}{(4x+3)^{n+1}}.$$

$$17.31 \frac{(-1)^{n-1} (n-1)!}{\ln 3 (x+5)^n}; 17.32 \frac{2 > (k+1)!}{(1-x)^{k+2}}; 17.33 \frac{(-1)^{n-1} n! 4^{n-1}}{(4x+3)^{n+1}}.$$

Yuqori tartibli hosilalar

Birinchi tartibli hosiladan olingan hosila, ya'ni

$$(y' = (f'(x)))' \text{ yoki } y'' = f''(x)$$

$y = f(x)$ funksiyaning ikkinchi tartibli hosilasi deyiladi.

Ikkinchi tartibli hosilaning hosilasiga uchinchi tartibli hosila deyiladi

va $y''' = f'''(x)$, $\frac{d^3 y}{dx^3}$ belgilarning biri bilan belgilanadi.

Umuman, $y = f(x)$ funksiyaning n - tartibli hosilasi deb, uning

$(n - 1)$ - tartibli hosilasining hosilasiga aytiladi va $y^{(n)}, f^{(n)}(x), \frac{d^n y}{dx^n}$

belgilarning biri bilan belgilanadi.

Ikkita funksiya ko'paytmasining n - tartibli hosilasi ushbu

$$\begin{aligned} (u \cdot v)^{(n)} &= u^{(n)}v + nu^{(n-1)}v' + \frac{n(n-1)}{2!}u^{(n-2)}v'' + \dots + \\ &+ \frac{n(n-1)\dots(n-k+1)}{k!}u^{(n-k)}v^{(k)} + \dots + nu^{(n-1)}v' + uv^{(n)} \end{aligned}$$

formuladan foydalanib topiladi. Bu formula Leybnits formulasi deyiladi.

Xususan,

$$(u \cdot v)' = u'v + u \cdot v'$$

18-masala. Funktsiyalarning talab qilingan tartibli hosilasini toping.

$$y = (x^3 + 3)e^{4x+3}, \quad y^{IV} = ?$$

Echim:

$$\begin{aligned} y' &= ((x^3 + 3)e^{4x+3})' = 3x^2e^{4x+3} + (x^3 + 3)e^{4x+3} \cdot 4 = \\ &= (4x^3 + 3x^2 + 12)e^{4x+3}. \end{aligned}$$

$$\begin{aligned} y'' &= ((4x^3 + 3x^2 + 12)e^{4x+3})' = \\ &= (12x^2 + 6x)e^{4x+3} + (4x^3 + 3x^2 + 12)e^{4x+3} \cdot 4 = \\ &= (16x^3 + 24x^2 + 6x + 48)e^{4x+3}. \end{aligned}$$

$$\begin{aligned} y''' &= ((16x^3 + 24x^2 + 6x + 48)e^{4x+3})' = \\ &= (48x^2 + 48x + 6)e^{4x+3} + (16x^3 + 24x^2 + 6x + 48)e^{4x+3} \cdot 4 = \\ &= (64x^3 + 144x^2 + 72x + 198)e^{4x+3}. \end{aligned}$$

$$\begin{aligned} y^{IV} &= ((64x^3 + 144x^2 + 72x + 198)e^{4x+3})' = \\ &= (192x^2 + 288x + 72)e^{4x+3} + (64x^3 + 144x^2 + 72x + 198)e^{4x+3} \cdot 4 = \\ &= (256x^3 + 768x^2 + 576x + 864)e^{4x+3}. \end{aligned}$$

$$1. y = (2x^2 - 7)\ln(x - 1), \quad y^V = ?$$

$$2. y = (3 - x^2)\ln^2 x, \quad y^{\text{III}} = ?$$

$$3. y = x \times \cos x^2, \quad y^{\text{III}} = ?$$

$$4. y = \frac{\ln(x - 1)}{\sqrt{x - 1}}, \quad y^{\text{III}} = ?$$

$$5. y = \frac{\log_2 x}{x^3}, \quad y^{\text{III}} = ?$$

$$6. y = (4x^3 + 5)e^{2x+1}, \quad y^V = ?$$

$$7. y = x^2 \times \sin(5x - 3), \quad y^{\text{III}} = ?$$

$$8. y = \frac{\ln x}{x^2}, \quad y^{\text{IV}} = ?$$

$$9. y = (2x + 3)\ln^2 x, \quad y^{\text{III}} = ?$$

$$10. y = (1 + x^2)\operatorname{arctg} x, \quad y^{\text{III}} = ?$$

$$11. y = \frac{\ln x}{x^3}, \quad y^{\text{IV}} = ?$$

$$12. y = (4x + 3) \times 2^{-x}, \quad y^V = ?$$

$$13. y = e^{1-2x} \times \sin(2 + 3x), \quad y^{\text{IV}} = ?$$

$$14. y = \frac{\ln(3 + x)}{3 + x}, \quad y^{\text{III}} = ?.$$

$$15. y = (2x^3 + 1)\cos x, \quad y^V = ?.$$

$$16. y = (x^2 + 3)\ln(x - 3), \quad y^{\text{IV}} = ?.$$

$$17. y = (1 - x - x^2)e^{\frac{x-1}{2}}, \quad y^{\text{IV}} = ?.$$

$$18. y = \frac{1}{x} \times \sin 2x, \quad y^{\text{III}} = ?.$$

$$19. y = (x + 7)\ln(x + 4), \quad y^V = ?.$$

$$20. y = (3x - 7) \times 3^{-x}, \quad y^{\text{IV}} = ?.$$

$$21. y = \frac{\ln(2x+5)}{2x+5}, \quad y''' = ?.$$

$$22. y = e^{\frac{x}{2}} \times \sin 2x, \quad y^{IV} = ?.$$

$$23. y = \frac{\ln x}{x^5}, \quad y''' = ?.$$

$$24. y = x \ln(1-3x), \quad y^{IV} = ?.$$

$$25. y = (x^2 + 3x + 1)e^{3x+2}, \quad y^V = ?.$$

$$26. y = (5x - 8)e^{-x}, \quad y^{IV} = ?.$$

$$27. y = \frac{\ln(x-2)}{x-2}, \quad y^V = ?.$$

$$28. y = e^{-x} \times (\cos 2x - 3 \sin 2x),$$

$$y^{IV} = ?.$$

$$29. y = (5x - 1) \ln^2 x, \quad y''' = ?.$$

$$30. y = \frac{\log_3 x}{x^2}, \quad y^{IV} = ?.$$

$$\text{Javoblar. 18.1 } \frac{8(x^2 - 5x - 11)}{(x-1)^5}; \text{ 18.2 } \frac{-4 \ln x}{x} + \frac{6 \ln x - 7x^2 - 15}{x^3};$$

$$18.3 - 24x^2 \times \cos x^2 + (8x^4 - 6) \times \sin x^2; \text{ 18.4 } \frac{46 - 15 \ln(x-1)}{8\sqrt{(x-1)^7}}; \text{ 18.5 } \frac{47 - 60 \ln x}{\ln 2 \times x^6};$$

$$18.6 32(4x^3 + 30x^2 + 60x + 35)e^{2x+1};$$

$$18.7 - 150x \times \sin(5x - 3) + (30 - 125x^2) \times \cos(5x - 3); \text{ 18.8 } \frac{-154 + 120 \ln x}{x^6};$$

$$18.9 \frac{4 \ln x \times (3 - x) - 18}{x^3}; \text{ 18.10 } \frac{4}{(1+x^2)^2}; \text{ 18.11 } \frac{-342 + 360 \ln x}{x^7};$$

$$18.12 \left(-\ln^5 2 \times (4x + 3) + 20 \ln^4 2 \right) \times 2^{-x};$$

$$18.13 - 122e^{1-2x} \times \sin(2+3x) - 597e^{1-2x} \times \cos(2+3x); \text{ 18.14 } \frac{11 - 6 \ln(3+x)}{(3+x)^4};$$

$$18.15 (30x^2 - 120) \cos x - (2x^3 - 120x + 1) \sin x; \text{ 18.16 } \frac{-2x^2 + 24x - 126}{(x-3)^4};$$

$$\begin{aligned}
& 18.17 - \frac{1}{16} \times (55 + 17x + x^2) e^{\frac{x-1}{2}}; 18.18 \frac{12x^2 - 6}{x^4} \times \sin 2x + \frac{12 - 8x^2}{x^3} \times \cos 2x; \\
& 18.19 \frac{-120x + 1680}{(x+4)^7}; 18.20 (7 \ln 3 - 12 - 3 \ln 3 \times x) \times \ln^3 3 \times 3^{-x}; \\
& 18.21 \frac{88 - 48 \ln(2x+5)}{(2x+5)^4}; 18.22 \frac{161}{16} \times e^{\frac{x}{2}} \times \sin 2x - 15 \times e^{\frac{x}{2}} \times \cos 2x; 18.23 \frac{107 - 210 \ln x}{x^8}; \\
& 18.24 - \frac{54(4 - 3x)}{(1 - 3x)^4}; 18.25 3^3 \times (9x^2 + 57x + 35) e^{3x+2}; \\
& 18.26 2^{-3} \times \ln^3 2 \times (5 \ln 2 \times x - 8 \ln 2 - 20); 18.27 \frac{274 - 120 \ln(x-2)}{(x-2)^6}; \\
& 18.28 - e^{-x} \times (79 \cos 2x + 3 \sin 2x); 18.29 \frac{6 - 2(5x+2) \ln x}{x^3}; 18.30 \frac{-154 + 120 \ln x}{x^6} > \ln 3.
\end{aligned}$$

Parametrik ko'rinishda berilgan funksiyaning hosilasi.

Agar $y = f(x)$ funksiya parametrik ko'rinishda berilgan bo'lsa, ya'ni

$$\begin{cases} x = j(t), \\ y = y(t), \end{cases} \quad a \leq t \leq b,$$

bo'lsa, u holda

$$f'(x) = \frac{y'(t)}{j'(t)}$$

va

$$f''(x) = \frac{y''(t)j'(t) - j''(t)y'(t)}{j'(t)^3}$$

bo'ladi.

19–masala. Parametrik ko'rinishda berilgan funksiyaning y'' hosilasini toping.

$$\begin{cases} \dot{x} = \ln t \\ \dot{y} = \operatorname{arctg} t \end{cases}$$

Echim:

$$x' = (\ln t)' = \frac{1}{t}$$

$$y' = (\operatorname{arctg} t)' = \frac{1}{t^2 + 1}$$

Natijada:

$$y'' = \frac{y'}{x'} = \frac{\frac{1}{t^2 + 1}}{\frac{1}{t}} = \frac{t}{t^2 + 1}$$

$$(y'')' = \frac{d}{dt} \frac{t}{t^2 + 1} = \frac{1 \cdot (1 + t^2) - t \cdot 2t}{(1 + t^2)^2} = \frac{1 - t^2}{(1 + t^2)^2}$$

U holda:

$$y''' = \frac{(y'')'}{x'} = \frac{\frac{1 - t^2}{(1 + t^2)^2}}{\frac{1}{t}} = \frac{t(1 - t^2)}{(1 + t^2)^2}.$$

$$1. \begin{cases} \dot{x} = \cos 2t \\ \dot{y} = 2 \sec^2 t \end{cases}$$

$$2. \begin{cases} \dot{x} = \sqrt{1 - t^2} \\ \dot{y} = \frac{1}{t} \end{cases}$$

$$3. \begin{cases} \dot{x} = e^t \cos t \\ \dot{y} = e^t \sin t \end{cases}$$

$$4. \begin{cases} \dot{x} = \operatorname{sh}^2 t \\ \dot{y} = \frac{1}{\operatorname{ch}^2 t} \end{cases}$$

$$5. \begin{cases} \dot{x} = t + \sin t \\ \dot{y} = 2 - \cos t \end{cases}$$

$$6. \begin{cases} \dot{x} = \frac{1}{t} \\ \dot{y} = \frac{1}{1 + t^2} \end{cases}$$

$$7. \begin{cases} \dot{x} = \sqrt{t} \\ \dot{y} = \frac{1}{\sqrt{1 - t}} \end{cases}$$

$$8. \begin{cases} \dot{x} = \sin t \\ \dot{y} = \sec t \end{cases}$$

$$9. \begin{cases} \dot{x} = \operatorname{tg} t \\ \dot{y} = \frac{1}{\sin 2t} \end{cases}$$

$$10. \begin{cases} \dot{x} = \sqrt{t-1} \\ \dot{y} = \frac{t}{\sqrt{1-t}} \end{cases}$$

$$11. \begin{cases} \dot{x} = \sqrt{t} \\ \dot{y} = \sqrt[3]{t-1} \end{cases}$$

$$12. \begin{cases} \dot{x} = \frac{\cos t}{1+2\cos t} \\ \dot{y} = \frac{\sin t}{1+2\cos t} \end{cases}$$

$$13. \begin{cases} \dot{x} = \sqrt{t^3-1} \\ \dot{y} = \ln t \end{cases}$$

$$14. \begin{cases} \dot{x} = \operatorname{sh}^2 t \\ \dot{y} = \operatorname{th}^2 t \end{cases}$$

$$15. \begin{cases} \dot{x} = \sqrt{t-1} \\ \dot{y} = \frac{1}{\sqrt{t}} \end{cases}$$

$$16. \begin{cases} \dot{x} = \cos^2 t \\ \dot{y} = \operatorname{tg}^2 t \end{cases}$$

$$17. \begin{cases} \dot{x} = \sqrt{t-3} \\ \dot{y} = \ln(t-2) \end{cases}$$

$$18. \begin{cases} \dot{x} = \sin t \\ \dot{y} = \ln(\cos t) \end{cases}$$

$$19. \begin{cases} \dot{x} = t + \sin t \\ \dot{y} = 2 + \cos t \end{cases}$$

$$20. \begin{cases} \dot{x} = t - \sin t \\ \dot{y} = 2 - \cos t \end{cases}$$

$$21. \begin{cases} \dot{x} = \cos t \\ \dot{y} = \ln(\sin t) \end{cases}$$

$$22. \begin{cases} \dot{x} = \cos t + t \times \sin t \\ \dot{y} = \sin t - t \times \cos t \end{cases}$$

$$23. \begin{cases} \dot{x} = e^t \\ \dot{y} = \arcsin t \end{cases}$$

$$24. \begin{cases} \dot{x} = \cos t \\ \dot{y} = \sin^4\left(\frac{t}{2}\right) \end{cases}$$

$$25. \begin{cases} \dot{x} = \operatorname{ch} t \\ \dot{y} = \sqrt[3]{\operatorname{sh}^2 t} \end{cases}$$

$$26. \begin{cases} \dot{x} = \operatorname{arctg} t \\ \dot{y} = \frac{t^2}{2} \end{cases}$$

$$27. \begin{cases} \dot{x} = 2(t - \sin t) \\ \dot{y} = 4(2 + \cos t) \end{cases}$$

$$28. \begin{cases} \dot{x} = \sin t - t \times \cos t \\ \dot{y} = \cos t + t \times \sin t \end{cases}$$

$$29. \begin{cases} \dot{x} = \frac{1}{t^2} \\ \dot{y} = \frac{1}{t^2 + 1} \end{cases}$$

$$30. \begin{cases} \dot{x} = \cos t + \sin t \\ \dot{y} = \sin 2t \end{cases}$$

Javoblar. 19.1 $\frac{1}{\cos^6 t}$; 19.2 $\frac{3-2t^2}{t^5}$; 19.3 $\frac{2}{e^t \times (\cos t - \sin t)^3}$; 19.4 $\frac{2}{\operatorname{ch}^6 t}$;

19.5 $\frac{1}{(1+\cos t)^2}$; 19.6 $\frac{2(t^2-3) \times^4}{(1+t^2)^3}$; 19.7 $(1+2t)\sqrt{1-t}$; 19.8 $\frac{1+2\sin^2 t}{\cos^5 t}$;

19.9 $-\frac{2 \times \cos^3 t}{\sin t \times \cos 2t}$; 19.10 $\frac{2}{\sqrt{(1-t)^3}}$; 19.11 $-\frac{2(t+3)}{9\sqrt[3]{(t-1)^5}}$; 19.12 $-\frac{(1+2\cos t)^3}{\sin^3 t}$;

19.13 $\frac{2(2-t^3)}{3t^6}$; 19.14 $\frac{2-6\operatorname{sh}^2 t}{\operatorname{ch}^6 t}$; 19.15 $\frac{(2t-3)\sqrt{t}}{t^3}$; 19.16 $\frac{2}{\cos^6 t}$;

$$\begin{aligned}
& \mathbf{19.17} - \frac{2t}{(t-2)^2}; \mathbf{19.18} - \frac{1+\sin^2 t}{\cos^4 t}; \mathbf{19.19} - \frac{1}{(1+\cos t)^2}; \mathbf{19.20} - \frac{1}{(1-\cos t)^2}; \\
& \mathbf{19.21} - \frac{1+\cos^2 t}{\sin^4 t}; \mathbf{19.22} \frac{1}{t > \cos^3 t}; \quad \mathbf{19.23} \frac{t^2+t-1}{e^{2t} \times \sqrt{(1-t^2)^3}}; \\
& \mathbf{19.24} \frac{\cos^2(t/2)+1}{4\cos^3(t/2)}; \mathbf{19.25} - \frac{2(3+\operatorname{ch}^2 t)}{9\operatorname{sh}^4 t}; \mathbf{19.26} \frac{1+3t^2}{1+t}; \mathbf{19.27} \frac{1}{(1-\cos(t))^2} 2; \\
& \mathbf{19.28} - \frac{1}{t > \sin^3 t}; \mathbf{19.29} - \frac{2t^6}{(1+t^2)^3}; \mathbf{19.30} 2.
\end{aligned}$$

20–masala. y funksiya berilgan tenglamaning yechimi bo'lishini ko'rsating.

$$\begin{aligned}
y &= -\sqrt{x^4 - x^2} \\
x \times y \times y &= x^4 \quad (1)
\end{aligned}$$

Echim:

$$y = \left(-\sqrt{x^4 - x^2} \right)' = -\frac{1}{2\sqrt{x^4 - x^2}} \times (4x^3 - 2x) = \frac{x - 2x^3}{\sqrt{x^4 - x^2}}$$

(1) tenglamaga qo'yamiz:

$$x \times \left(-\sqrt{x^4 - x^2} \right) \times \frac{x - 2x^3}{\sqrt{x^4 - x^2}} - \left(-\sqrt{x^4 - x^2} \right)^2 = x^4.$$

Soddalashtiramiz:

$$\begin{aligned}
x \times (2x^3 - x) - (x^4 - x^2) &= x^4 \\
2x^4 - x^2 - x^4 + x^2 &= x^4. \\
x^4 &= x^4.
\end{aligned}$$

Tenglik o'rinli. y funksiya berilgan (1) tenglamaning yechimi bo'ladi.

$$1. \quad y = x \times e^{-\frac{x^2}{2}} \\ x \times y \oslash = (1 - x^2)y \quad (1)$$

$$3. \quad y = 5 \times e^{-2x} + \frac{e^x}{3} \\ y \oslash + 2y = e^x \quad (1)$$

$$5. \quad y = x \times \sqrt{1 - x^2} \\ y \times y \oslash = x - 2x^3 \quad (1)$$

$$7. \quad y = -\frac{1}{3x + c} \\ y \oslash = 3y^2 \quad (1)$$

$$9. \quad y = \sqrt{x^2 - c \times x} \\ (x^2 + y^2)dx - 2 \times x \times y \times dy = 0 \quad (1) \quad 10. \quad y = x \times (c - \ln x) \\ (x - y)dx + x \times dy = 0 \quad (1)$$

$$11. \quad y = e^{\operatorname{tg} \frac{x}{2}} \\ y \oslash \sin x = y \ln y \quad (1)$$

$$13. \quad y = \frac{b + x}{1 + bx} \\ y - x \times y \oslash = b(1 + x^2 \times y \oslash) \quad (1)$$

$$15. \quad y = \sqrt{\ln \frac{e^x + e^x}{e} \frac{2}{2} + 1} \\ (1 + e^x) \times y \times y \oslash = e^x \quad (1)$$

$$17. \quad y = -\sqrt{\frac{2}{x^2} - 1} \\ 1 + y^2 + x \times y \times y \oslash = 0 \quad (1)$$

$$2. \quad y = \frac{\sin x}{x} \\ x \times y \oslash + y = \cos x \quad (1)$$

$$4. \quad y = 2 + c \times \sqrt{1 - x^2} \\ (1 - x^2) \times y \oslash + xy = 2x \quad (1)$$

$$6. \quad y = \frac{c}{\cos x} \\ y \oslash - \operatorname{tg} x \times y = 0 \quad (1)$$

$$8. \quad y = \ln(c + e^x) \\ y \oslash = e^{x-y} \quad (1)$$

$$12. \quad y = \frac{1 + x}{1 - x} \\ y \oslash = \frac{1 + y^2}{1 + x^2} \quad (1)$$

$$14. \quad y = \sqrt{2 + 3x - 3x^2} \\ y \times y \oslash = \frac{1 - 2x}{y} \quad (1)$$

$$16. \quad y = \operatorname{tg} x(\ln 3x) \\ (1 + y^2)dx = x \times dy \quad (1)$$

$$18. \quad y = \sqrt[3]{x - \ln x - 1} \\ \ln x + y^3 - 3 \times x \times y^2 \times y \oslash = 0 \quad (1)$$

19. $y = a + \frac{7x}{ax+1}$
 $y - x \times y \cancel{=} a(1 + x^2 \times y \cancel{=}) \quad (1)$
20. $y = a \times \operatorname{tg} \sqrt{\frac{a}{x} - 1}$
 $a^2 + y^2 + 2x\sqrt{ax - x^2} \times y \cancel{=} 0 \quad (1)$
21. $y = \sqrt[4]{\sqrt{x} + \sqrt{x+1}}$
 $8 \times x \times y \cancel{=} y = \frac{-1}{y^3 \sqrt{x+1}} \quad (1)$
22. $y = (x+1) \times e^{x^2}$
 $y \cancel{=} 2xy = 2 \times x \times e^{x^2} \quad (1)$
23. $y = \frac{2}{x^3+1} + \frac{1}{x}$
 $x \times (x^3+1) \times y \cancel{=} (2x^3 - 1)y = \frac{x^3 - 2}{x} \quad (1)$
24. $y = e^{x+x^2} + 2e^x$
 $y \cancel{=} y = 2xe^{x+x^2} \quad (1)$
25. $y = -x \times \cos x + 3x$
 $x \times y \cancel{=} y + x^2 \sin x \quad (1)$
26. $y = \frac{1}{\sqrt{\sin x + x}}$
 $2 \sin x \times y \cancel{=} y \cos x == y^3 (x \times \cos x - \sin x) \quad (1)$
27. $y = \frac{x}{x-1} + x^2$
 $x \times (x-1) \times y \cancel{=} y = x^2(2x-1) \quad (1)$
28. $y = \frac{x}{\cos x}$
 $y \cancel{=} y \times \operatorname{tg} x = \sec x \quad (1)$
29. $y = (x+1)^n \times (e^x - 1)$
 $y \cancel{=} \frac{n \times y}{x+1} = e^x (x+1)^n \quad (1)$
30. $y = 2 \frac{\sin x}{x} + \cos x$
 $x \times \sin x \times y \cancel{=} (\sin x - x \times \cos x)y = \sin x \times \cos x - x \quad (1)$

IV BOB. Grafiklar

Funksiyaning grafigi

1-masala. Birinchi tartibli hosila yordamida funksiyaning grafigini yasang.


$$y = x^2(x - 2)^2.$$

1) $D(y) = (-\infty; +\infty)$.

2) Funksiya juft ham, toq ham emas.

3) $y' = 2x(x - 2)^2 + 2x^2(x - 2) = 4x(x - 2)(x - 1)$.

$$y' = 0 \text{ da, } \begin{cases} x = 2, \\ x = 1, \\ x = 0 \end{cases}$$




(0;0)- minimum nuqta,

(2;0)- minimum nuqta ,

(1;1)- maksimum nuqta .

Функциянинг графиги



1. $y = 2x^3 - 9x^2 + 12x - 9$.

2. $y = 3x - x^3$.

3. $y = (x^3 - 9x^2)/4 + 6x - 9$.

4. $y = 2 - 3x^2 - x^3$.

5. $y = (x+1)^2(x-1)^2$.

6. $y = 2x^3 - 3x^2 - 4$.

7. $y = 3x^2 - 2 - x^3$.

8. $y = (x-1)^2(x-3)^2$. 9.

$$y = (x^3 + 3x^2)/4 - 5.$$

$$y = 16x^2(x - 1)^2.$$

$$y = 2 - 12x^2 - 8x^3.$$

$$y = 2x^3 + 9x^2 + 12x.$$

$$y = (2x - 1)^2(2x - 3)^2.$$

$$y = x(12 - x^2)/8.$$

$$y = 27(x^3 + x^2)/4 - 5.$$

$$y = -(x^2 - 4)^2/16.$$

$$y = (6x^2 - x^3 - 16)/8.$$

$$y = 16x^3 - 12x^2 - 4.$$

$$y = -(x+1)^2(x-3)^2/16.$$

$$10. y = 6x - 8x^3. \quad 11.$$

$$12. y = 2x^3 + 3x^2 - 5. \quad 13.$$

$$14. y = (2x+1)^2(2x-1)^2. \quad 15.$$

$$16. y = 12x^2 - 8x^3 - 2. \quad 17.$$

$$18. y = 27(x^3 - x^2)/4 - 4. \quad 19.$$

$$20. y = x^2(x-4)^2/16. \quad 21.$$

$$22. y = (16 - 6x^2 - x^3)/8. \quad 23.$$

$$24. y = 16x^3 - 36x^2 + 24x - 9. \quad 25.$$

$$26. y = -(x-2)^2(x-6)^2/16. \quad 27.$$

$$28. y = (11 + 9x - 3x^2 - x^3)/8. \quad 29.$$

$$30. y = 16x^3 + 12x^2 - 5.$$

2-masala. Birinchi tartibli hosila yordamida funksiyaning grafigini yasang.


$$y = 1 - \sqrt[3]{x^2 + 2x}.$$

1) $D(y) = (-\infty; +\infty).$


2) Funksiya juft ham, toq ham emas.

3) $y' = -\frac{2x+2}{3\sqrt[3]{(x^2+2x)^2}}.$

$y' = 0$ da, $x = 1$; $x = 0$ va $x = -2$ nuqtalarda y' mavjud emas.



$(-1; 2)$ – maksimum nuqta .



$$1. y = 2x - 3\sqrt[3]{x^2}.$$

$$3. y = -\frac{12\sqrt[3]{6(x-1)^2}}{x^2 + 2x + 9}.$$

$$5. y = 2x + 6 - 3\sqrt[3]{(x+3)^2}.$$

$$7. y = 1 - \sqrt[3]{x^2 + 4x + 3}.$$

$$9. y = -\frac{6\sqrt[3]{6x^2}}{x^2 + 4x + 12}.$$

$$11. y = \frac{3\sqrt[3]{6(x-4)^2}}{x^2 - 4x + 12}.$$

$$13. y = \sqrt[3]{x^2 + 4x + 3}.$$

$$15. y = 6\sqrt[3]{(x-2)^2} - 4x + 8.$$

$$17. y = 2 + \sqrt[3]{8x(x+2)}.$$

$$19. y = \sqrt[3]{x^2 + 6x + 8}.$$

$$21. y = -\frac{3\sqrt[3]{6(x+2)^2}}{x^2 + 8x + 24}.$$

$$23. y = 1 - \sqrt[3]{x^2 - 4x + 3}.$$

$$25. y = \frac{6\sqrt[3]{6(x+3)^2}}{x^2 + 10x + 33}.$$

$$27. y = -\frac{6\sqrt[3]{6(x-6)^2}}{x^2 - 8x + 24}.$$

$$29. y = \frac{3\sqrt[3]{6(x-1)^2}}{2(x^2 + 2x + 9)}.$$

$$2. y = \frac{12\sqrt[3]{6(x-2)^2}}{x^2 + 8}.$$

$$4. y = 1 - \sqrt[3]{x^2 + 2x}.$$

$$6. y = \frac{6\sqrt[3]{6(x-3)^2}}{x^2 - 2x + 9}.$$

$$8. y = 3\sqrt[3]{(x-3)^2} - 2x + 6.$$

$$10. y = 4x + 8 - 6\sqrt[3]{(x+2)^2}.$$

$$12. y = \sqrt[3]{x(x+2)}.$$

$$14. y = -\frac{3\sqrt[3]{6(x+1)^2}}{x^2 + 6x + 17}.$$

$$16. y = \frac{3\sqrt[3]{6(x-5)^2}}{x^2 - 6x + 17}.$$

$$18. y = 6x - 6 - 9\sqrt[3]{(x-1)^2}.$$

$$20. y = \sqrt[3]{4x(x-1)}.$$

$$22. y = \sqrt[3]{x(x-2)}.$$

$$24. y = 9\sqrt[3]{(x+1)^2} - 6x - 6.$$

$$26. y = 8x - 16 - 12\sqrt[3]{(x-2)^2}.$$

$$28. y = 12\sqrt[3]{(x+2)^2} - 8x - 16.$$

$$30. y = 3\sqrt[3]{(x+4)^2} - 2x - 8.$$

Funksiyaning eng katta va eng kichik qiymatlari

3–masala. Berilgan kesmada funksiyaning eng katta va eng kichik

qiymatlarini toping.

$$y = 2x^2 + \frac{108}{x} - 59, \quad [2, 4].$$

$$x \neq 0.$$

$$y' = 4x - \frac{108}{x^2} = \frac{4x^3 - 108}{x^2}.$$

$$y' = 0 \text{ da, } x = 3 \in [2; 4];$$

$$x = 0 \notin [2; 4] \text{ da } y' \text{ mavjud emas.}$$

$$y(2) = 3,$$

$$y(3) = -5,$$

$$y(4) = 0.$$

$$\max_{[2;4]} y = y(2) = 3;$$

$$\min_{[2;4]} y = y(3) = -5;$$

$$1. y = x^2 + \frac{16}{x} - 16, [1, 4].$$

$$2. y = 4 - x - \frac{4}{x^2}, [1, 4].$$

$$3. y = \sqrt[3]{2(x-2)^2(8-x)} - 1, [0, 6]. \quad 4. y = \frac{2(x^2+3)}{x^2-2x+5}, [-3, 3].$$

$$5. y = 2\sqrt{x} - x, [0, 4].$$

$$6. y = 1 + \sqrt[3]{2(x-1)^2(x-7)}, [-1, 5].$$

$$7. y = x - 4\sqrt{x} + 5, [1, 9].$$

$$8. y = \frac{10x}{1+x^2}, [0, 3].$$

$$9. y = \sqrt[3]{2(x+1)^2(5-x)} - 2, [-3, 3]. \quad 10. y = 3 - x - \frac{4}{(x+2)^2}, [-1, 2].$$

$$11. y = \sqrt[3]{2x^2(x-3)}, [-1, 6].$$

$$12. y = \frac{2(-x^2+7x-7)}{x^2-2x+2}, [1, 4].$$

$$13. y = x - 4\sqrt{x+2} + 8, [-1, 7].$$

$$14. y = \sqrt[3]{2(x-2)^2(5-x)}, [1, 5].$$

$$15. y = \frac{4x}{4+x^2}, [-4, 2].$$

$$16. y = -\frac{x^2}{2} + \frac{8}{x} + 8, [-4, -1].$$

$$17. y = \sqrt[3]{2x^2(x-6)}, [-2, 4].$$

$$18. y = \frac{-2x(2x+3)}{x^2+4x+5}, [1, 4].$$

19. $y = -\frac{2(x^2 + 3)}{x^2 + 2x + 5}, [-5, 1]$ 20. $y = \sqrt[3]{2(x-1)^2(x-4)}, [0, 4]$

21. $y = x^2 - 2x + \frac{16}{x-1} - 13, [2, 5]$ 22. $y = 2\sqrt{x-1} - x + 2, [1, 5]$

23. $y = \sqrt[3]{2(x+2)^2(1-x)}, [-3, 4]$ 24. $y = -\frac{x^2}{2} + 2x + \frac{8}{x-2} + 5, [-2, 1]$

25. $y = 8x + \frac{4}{x^2} - 15, \left[-\frac{1}{2}, \frac{1}{2}\right]$ 26. $y = \sqrt[3]{2(x+2)^2(x-4)} + 3, [-4, 2]$

27. $y = x^2 + 4x + \frac{16}{x+2} - 9, [-1, 2]$ 28. $y = \frac{4}{x^2} - 8x - 15, \left[-\frac{1}{2}, \frac{1}{2}\right]$

29. $y = \sqrt[3]{2(x+1)^2(x-2)}, [-2, 5]$ 30. $y = \frac{10x+10}{x^2+2x+2}, [-1, 2]$

Javoblar. 3.1 $f_{\max} = y(1) = y(4) = 4; f_{\min} = y(-2) = -4$; 3.2 $f_{\max} = y(2) = 1; f_{\min} = y(1) = -1$; 3.3 $f_{\max} = y(0) = y(6) = 3; f_{\min} = y(2) = -1$;

3.4 $f_{\max} = y(-1) = 1; f_{\min} = y(3) = 3$; 3.5 $f_{\max} = y(1) = 1; f_{\min} = y(4) = 0$;

3.6 $f_{\max} = y(1) = 1; f_{\min} = y(5) = -3$; 3.7 $f_{\max} = y(1) = y(9) = 2; f_{\min} = y(4) = 1$;

3.8 $f_{\max} = y(1) = 5; f_{\min} = y(0) = 0$; 3.9 $f_{\max} = y(-3) = y(3) = 2; f_{\min} = y(-1) = -2$;

3.10 $f_{\max} = y(0) = 2; f_{\min} = y(-1) = 0$; 3.11 $f_{\max} = y(6) = 6; f_{\min} = y(-1) = y(2) = -2$;

3.12 $f_{\max} = y(2) = 3; f_{\min} = y(1) = -2$;

3.13 $f_{\max} = y(-1) = y(7) = 3; f_{\min} = y(2) = 2$; 3.14 $f_{\max} = y(1) = y(4) = 2; f_{\min} = y(2) = y(5) = 0$;

3.15 $f_{\max} = y(2) = 1; f_{\min} = y(-2) = -1$;

3.16 $f_{\max} = y(-2) = 2; f_{\min} = y(-4) = -2$; 3.17 $f_{\max} = y(0) = 0; f_{\min} = y(-2) = y(4) = -4$;

3.18 $f_{\max} = y(1) = -1; f_{\min} = y(4) = -\frac{88}{37}$;

3.19 $f_{\max} = y(1) = -1; f_{\min} = y(-3) = -3$;

3.20 $f_{\max} = y(1) = y(4) = 0; f_{\min} = y(0) = -2$; 3.21 $f_{\max} = y(5) = 6; f_{\min} = y(3) = -2$;

3.22 $f_{\max} = y(2) = 2; f_{\min} = y(1) = y(5) = 1$;

3.23 $f_{\max} = y(-3) = y(0) = 2; f_{\min} = y(4) = -6$; 3.24 $f_{\max} = y(0) = 1; f_{\min} = y(-2) = -3$;

3.25 $f_{\max} = y(0) = 5; f_{\min} = y(1) = -3$; 3.26 $f_{\max} = y(-2) = 3; f_{\min} = y(2) = y(-4) = -1$;

3.27 $f_{\max} = y(2) = 7; f_{\min} = y(0) = -1$;

3.28 $f_{\max} = y(0) = 5; f_{\min} = y(-1) = -3$; 3.29 $f_{\max} = y(5) = 6; f_{\min} = y(-2) = y(1) = -2$;

3.30 $f_{\max} = y(-1) = 0; f_{\min} = y(0) = -5$.

4–masala. Yuqori tartibli hosilalar yordamida berilgan nuqta atrofida funksiyani tekshiring.

$$y = 4x - x^2 - 2\cos(x - 2),$$

$$x_0 = 2.$$

$$y' = 4 - 2x + 2\sin(x - 2), \quad y'(2) = 0;$$

$$y'' = -2 + 2\cos(x - 2), \quad y''(2) = 0;$$

$$y''' = -2\sin(x - 2), \quad y'''(2) = 0;$$

$$y^{IV} = -2\cos(x - 2), \quad y^{IV}(2) = -2.$$

$y^{IV} \neq 0$ ekanligidan $x_0 = 2$ nuqtada maksimumga erishadi.

1. $y = x^2 - 4x - (x - 2)\ln(x - 1),$
 $x_0 = 2.$

2. $y = 6e^{x-2} - x^3 + 3x^2 - 6x,$
 $x_0 = 2.$

3. $y = 2\ln(x+1) - 2x + x^2 + 1,$
 $x_0 = 0.$

4. $y = 2x - x^2 - 2\cos(x - 1),$
 $x_0 = 1.$

5. $y = \cos^2(x+1) + x^2 + 2x,$
 $x_0 = -1.$

6. $y = 2\ln x + x^2 - 4x + 3,$
 $x_0 = 1.$

7. $y = 1 - 2x - x^2 - 2\cos(x+1),$
 $x_0 = -1.$

8. $y = x^2 + 6x + 8 - 2e^{x+2},$
 $x_0 = -2.$

9. $y = 4x + x^2 - 2e^{x+1},$
 $x_0 = -1.$

10. $y = (x+1)\sin(x+1) - 2x - x^2,$
 $x_0 = -1.$

11. $y = 6e^{x-1} - 3x - x^3,$
 $x_0 = 1.$

12. $y = 2x + x^2 - (x+1)\ln(2+x),$
 $x_0 = -1.$

13. $y = \sin^2(x+1) - 2x - x^2,$
 $x_0 = -1.$

14. $y = x^2 + 4x + \cos^2(x+2),$
 $x_0 = -2.$

15. $y = x^2 + 2\ln(x+2),$
 $x_0 = -1.$

16. $y = 4x - x^2 + (x-2)\sin(x-2),$
 $x_0 = 2.$

17. $y = 6e^x - x^3 - 3x^2 - 6x - 5,$
 $x_0 = 0.$

18. $y = x^2 - 2x - 2e^{x-2},$
 $x_0 = 2.$

19. $y = \sin^2(x+2) - x^2 - 4x - 4,$
 $x_0 = -2.$

20. $y = \cos^2(x-1) + x^2 - 2x,$
 $x_0 = 1.$

21. $y = x^2 - 2x - (x-1)\ln x,$
 $x_0 = 1.$

22. $y = (x-1)\sin(x-1) + 2x - x^2,$
 $x_0 = 1.$

23. $y = x^2 - 4x + \cos^2(x-2),$
 $x_0 = 2.$

24. $y = x^4 + 4x^3 + 12x^2 + 24(x+1 - e^x),$
 $x_0 = 0.$

25. $y = \sin^2(x-2) - x^2 + 4x - 4,$
 $x_0 = 2.$

26. $y = 6e^{x+1} - x^3 - 6x^2 - 15x - 16,$
 $x_0 = -1.$

27. $y = \sin x + \operatorname{sh} x - 2x,$
 $x_0 = 0.$

28. $y = \sin^2(x-1) - x^2 + 2x,$
 $x_0 = 1.$

29. $y = \cos x + \operatorname{ch} x,$
 $x_0 = 0.$

30. $y = x^2 - 2e^{x-1},$
 $x_0 = 1.$

Funksiyaning aimptotalari

5–masala. Quyidagi funksiyaning aimptotalarini toping va grafigini yasang.

$$y = \frac{17 - x^2}{4x - 5}$$

1) $D(y) = \left(-\frac{5}{4} \right)^{-1} = -\frac{4}{5}; +\infty \rightarrow -\frac{4}{5}.$

2) Funksiya juft ham, toq ham emas.

$$3. a) \lim_{x \rightarrow \frac{5}{4} - 0} \frac{17 - x^2}{4x - 5} = -\infty,$$

$$\lim_{x \rightarrow \frac{5}{4} + 0} \frac{17 - x^2}{4x - 5} = +\infty,$$

$x = \frac{5}{4}$ —vertikal asimptota.


$$6) k = \lim_{x \rightarrow \infty} \frac{f(x)}{x} = \lim_{x \rightarrow \infty} \frac{17 - x^2}{x(4x - 5)} = -\frac{1}{4}.$$

$$b = \lim_{x \rightarrow \infty} (f(x) - kx) = \lim_{x \rightarrow \infty} \left(\frac{17 - x^2}{4x - 5} + \frac{x}{4} \right) = \lim_{x \rightarrow \infty} \frac{68 - 5x}{16x - 20} = -\frac{5}{16}.$$

Demak, $y = -\frac{1}{4}x - \frac{5}{16}$ —og'ma asimptota.

$$4) y'' = \frac{-2x(4x - 5) - 4(17 - x^2)}{(4x - 5)^2} = -\frac{4x^2 + 10x + 68}{(4x - 5)^2}.$$


$x = \frac{5}{4}$ da y'' mavjud emas.



5) O'qlar bilan kesishish nuqtalarini topamiz:

$$x = 0 \text{ da } y = -\frac{17}{5}.$$

$$y = 0 \text{ da } x = \pm 4,12.$$



$$1. y = \frac{x^2 + 1}{\sqrt{4x^2 - 3}}.$$

$$2. y = \frac{x^3 - 4x}{3x^2 - 4}.$$

$$3. y = \frac{4x^2 + 9}{4x + 8}.$$

$$4. y = \frac{4x^3 + 3x^2 - 8x - 2}{2 - 3x^2}.$$

$$5. y = \frac{x^2 - 3}{\sqrt{3x^2 - 2}}.$$

$$6. y = \frac{2x^2 - 6}{x - 2}.$$

$$7. y = \frac{2x^3 + 2x^2 - 3x - 1}{2 - 4x^2}.$$

$$8. y = \frac{x^3 - 5x}{5 - 3x^2}.$$

$$9. y = \frac{x^2 - 6x + 4}{3x - 2}.$$

$$10. y = \frac{2 - x^2}{\sqrt{9x^2 - 4}}.$$

$$11. y = \frac{4x^3 - 3x}{4x^2 - 1}.$$

$$12. y = \frac{3x^2 - 7}{2x + 1}.$$

$$13. y = \frac{x^2 + 16}{\sqrt{9x^2 - 8}}.$$

$$14. y = \frac{x^3 + 3x^2 - 2x - 2}{2 - 3x^2}.$$

$$15. y = \frac{21 - x^2}{7x + 9}.$$

$$16. y = \frac{2x^2 - 1}{\sqrt{x^2 - 2}}.$$

$$17. y = \frac{2x^3 - 3x^2 - 2x + 1}{1 - 3x^2}.$$

$$18. y = \frac{x^2 - 11}{4x - 3}.$$

$$19. y = \frac{2x^2 - 9}{\sqrt{x^2 - 1}}.$$

$$21. y = \frac{x^2 + 2x - 1}{2x + 1}.$$

$$23. y = \frac{x^2 + 6x + 9}{x + 4}.$$

$$25. y = \frac{x^2 - 2x + 2}{x + 3}.$$

$$27. y = \frac{3x^2 - 10}{3 - 2x}.$$

$$29. y = \frac{-8 - x^2}{\sqrt{x^2 - 4}}.$$

$$20. y = \frac{x^3 - 2x^2 - 3x + 2}{1 - x^2}.$$

$$22. y = \frac{x^3 + x^2 - 3x - 1}{2x^2 - 2}.$$

$$24. y = \frac{3x^2 - 10}{\sqrt{4x^2 - 1}}.$$

$$26. y = \frac{2x^3 + 2x^2 - 9x - 3}{2x^2 - 3}.$$

$$28. y = \frac{-x^2 - 4x + 13}{4x + 3}.$$

$$30. y = \frac{9 - 10x^2}{\sqrt{4x^2 - 1}}.$$

Funksiyalarni tekshirish va grafiklarini chizish

6-masala. Quyidagi funktsiyani tekshiring va ularni grafigini yasang.

$$y = \frac{x^2 - 3x + 3}{x - 1}.$$

1) $D(y) = (-\infty; 1) \cup (1; +\infty)$.

2) Funktsiya juft ham, toq ham emas.

3. a) $\lim_{x \rightarrow 1-0} \frac{x^2 - 3x + 3}{x - 1} = -\infty$,

$\lim_{x \rightarrow 1+0} \frac{x^2 - 3x + 3}{x - 1} = +\infty$,

$x = 1$ – vertikal asimptota.

b) $k = \lim_{x \rightarrow \infty} \frac{f(x)}{x} = \lim_{x \rightarrow \infty} \frac{x^2 - 3x + 3}{x(x - 1)} = 1$.


$$b = \lim_{x \rightarrow \infty} (f(x) - kx) = \lim_{x \rightarrow \infty} \frac{x^2 - 3x + 3}{x - 1} - x = \lim_{x \rightarrow \infty} \frac{-2x + 3}{x - 1} = -2.$$

Demak, $y = x - 2$ —og' ma asimptota.

$$4) y' = \frac{-(2x - 3)(x - 1) - (x^2 - 3x + 3)}{(x - 1)^2} = \frac{x(x - 2)}{(x - 1)^2}.$$

$$\begin{cases} x = 0, \\ x = 1 \end{cases} \text{ da } y' = 0$$

$x = 1$ da y' mavjud emas.




$(0; -3)$ —funksiyaning maksimum nuqtasi

$(2; 1)$ —funksiyaning minimum nuqtasi.

$$5) y'' = \frac{(2x - 2)(x - 1)^2 - 2(x - 1)(x^2 - 2x)}{(x - 1)^4} = \frac{2}{(x - 1)^3},$$


$x = 1$ da y'' mavjud emas.



6) O'qlar bilan kesishish nuqtalarini topamiz:

$x = 0$ da $y = -3$.

$y = 0$ da kvadrat tenglama ildizlarga ega emas. Demak funksiyaning grafigi Ox oqi bilan kesishmaydi.



$$1. y = \frac{x^3 + 4}{x^2}.$$

$$2. y = \frac{x^2 - x + 1}{x - 1}.$$

$$3. y = \frac{2}{x^2 + 2x}.$$

$$4. y = \frac{4x^2}{3 + x^2}.$$

$$5. y = \frac{12x}{9 + x^2}.$$

$$6. y = \frac{4 - x^3}{x^2}.$$

$$7. y = \frac{x^2 - 4x + 1}{x - 4}.$$

$$8. y = \frac{2x^3 + 1}{x^2}.$$

$$9. y = \frac{(x - 1)^2}{x^2}.$$

$$10. y = \frac{x^2}{(x - 1)^2}.$$

$$11. y = \frac{x^2 + 1}{x^2}.$$

$$12. y = \frac{12 - 3x^2}{x^2 + 12}.$$

$$13. y = \frac{9 + 6x - 3x^2}{x^2 - 2x + 13}.$$

$$14. y = \frac{-8x}{x^2 + 4}.$$

$$15. y = \frac{x^2 - 1}{x^2 + 1}.$$

$$16. y = \frac{3x^4 + 1}{x^3}.$$

$$17. y = \frac{4x}{(x + 1)^2}.$$

$$18. y = \frac{8(x - 1)}{(x + 1)^2}.$$

$$19. y = \frac{1 - 2x^3}{x^2}.$$

$$20. y = \frac{4}{x^2 + 2x - 3}.$$

$$21. y = \frac{4}{3 + 2x - x^2}.$$

$$22. y = \frac{x^2 + 2x - 7}{x^2 + 2x - 3}.$$

$$23. y = \frac{1}{x^4 - 1}.$$

$$24. y = -\frac{x^2}{x^2 + 2}.$$

$$25. y = \frac{x^3 - 32}{x^2}.$$

$$26. y = \frac{4(x + 1)^2}{x^2 + 2x + 4}.$$

$$27. y = \frac{3x - 2}{x^3}.$$

$$28. y = \frac{x^2 - 6x + 9}{(x - 1)^2}. \quad 29. y = \frac{x^3 - 27x + 54}{x^3}. \quad 30. y = \frac{x^3 - 4}{x^2}.$$

Funksiyalarni tekshirish va grafiklarini chizish

7-masala. Quyidagi funktsiyani tekshiring va ularni grafigini yasang.

$$y = \frac{e^{2(x+1)}}{2(x+1)}.$$

1) $D(y) = (-\infty; 1) \cup (1; +\infty).$

2) Funktsiya juft ham, toq ham emas.

3. a) $\lim_{x \rightarrow -1-0} \frac{e^{2(x+1)}}{2(x+1)} = -\infty,$

$$\lim_{x \rightarrow -1+0} \frac{e^{2(x+1)}}{2(x+1)} = +\infty,$$

$x = -1$ – vertikal asimptota.

b) $k = \lim_{x \rightarrow -\infty} \frac{f(x)}{x} = \lim_{x \rightarrow -\infty} \frac{e^{2(x+1)}}{2(x+1)x} = -\infty,$

$$k = \lim_{x \rightarrow +\infty} \frac{f(x)}{x} = \lim_{x \rightarrow +\infty} \frac{e^{2(x+1)}}{2(x+1)x} = 0.$$


$$b = \lim_{x \rightarrow +\infty} \frac{e^{2(x+1)}}{2(x+1)} = 0..$$

Demak, $y = 0$ – gorizonttal asimptota.

4) $y' = \frac{1}{2} \times \frac{2(x+1)e^{2(x+1)} - e^{2(x+1)}}{(x+1)^2} = \frac{(2x+1)e^{2(x+1)}}{2(x+1)^2}.$

$x = -0,5$ da $y' = 0$,


$x = -1$ da y' mavjud emas.



5) $x = -\frac{1}{2}$; $e^{-\frac{1}{2}}$ funksiyaning minimum nuqtasi.

$$5) y = \frac{(2x^2 - 2x + 1)e^{2(x+1)}}{(x+1)^3},$$

$x = -1$ da y mavjud emas.



6) O'qlar bilan kesishish nuqtalarini topamiz:

$$x = 0 \text{ da } y = \frac{e^2}{2}.$$

$y = 0$ da kvadrat tenglama ildizlarga ega emas. Demak funksiyaning grafigi Ox oqi bilan kesishmaydi.

$$1. y = (2x + 3)e^{-2(x+1)}.$$

$$2. y = 3 \ln \frac{x}{x-3} - 1.$$

$$3. y = (3 - x)e^{x-2}.$$

$$4. y = \frac{e^{2-x}}{2-x}.$$

$$5. y = \ln \frac{x}{x+2} + 1.$$

$$6. y = (x-2)e^{3-x}.$$

$$7. y = \frac{e^{2(x-1)}}{2(x-1)}.$$

$$8. y = 3 - 3 \ln \frac{x}{x+4}.$$

$$9. y = -(2x+1)e^{2(x+1)}.$$

$$10. y = \frac{e^{2(x+2)}}{2(x+2)}.$$

$$11. y = \ln \frac{x}{x-2} - 2.$$

$$12. y = (2x+5)e^{-2(x+2)}.$$

$$13. y = \frac{e^{3-x}}{3-x}.$$

$$14. y = 2 \ln \frac{x}{x+1} - 1.$$

15. $y = (4 - x)e^{x-3}.$

16. $y = - \frac{e^{-2(x+2)}}{2(x+2)}.$

17. $y = 2 \ln \frac{x+3}{x} - 3.$

18. $y = (2x - 1)e^{2(1-x)}.$

19. $y = - \frac{e^{-(x+2)}}{x+2}.$

20. $y = 2 \ln \frac{x}{x-4} - 3.$

21. $y = -(x+1)e^{(x+2)}.$

22. $y = - \frac{e^{x+3}}{x+3}.$

23. $y = \ln \frac{x}{x+5} - 1.$

24. $y = -(2x+3)e^{2(x+2)}.$

25. $y = - \frac{e^{-2(x-1)}}{2(x-1)}.$

26. $y = \ln \frac{x-5}{x} + 2.$

27. $y = (x+4)e^{-(x+3)}.$

28. $y = - \frac{e^{x-3}}{x-3}.$

29. $y = \ln \frac{x+6}{x} - 1.$

30. $y = 2 \ln \frac{x-1}{x} + 1.$

Funksiyalarni tekshirish va grafiklarini chizish

8-masala. Quyidagi funktsiyani tekshiring va ularni grafigini yasang.

$$y = \sqrt[3]{(2-x)(x^2-4x+1)}.$$

1) $D(y) = (-\infty; +\infty).$

2) Funktsiya juft ham, toq ham emas.

3. a) vertikal asimptotalari yo'q.

$$b) \quad k = \lim_{x \rightarrow -\infty} \frac{f(x)}{x} = \lim_{x \rightarrow -\infty} \frac{\sqrt[3]{(2-x)(x^2-4x+1)}}{x} = -1,$$


$$\begin{aligned}
 b &= \lim_{x \rightarrow +\infty} (\sqrt[3]{(2-x)(x^2-4x+1)} - x) = \\
 &= \lim_{x \rightarrow +\infty} \frac{(2-x)(x^2-4x+1) - x^3}{\sqrt[3]{(2-x)^2(x^2-4x+1)^2} + x\sqrt[3]{(2-x)(x^2-4x+1)} + x^2} = \\
 &= \lim_{x \rightarrow +\infty} \frac{2-9x+6x^2-2x^3}{\sqrt[3]{(2-x)^2(x^2-4x+1)^2} + x\sqrt[3]{(2-x)(x^2-4x+1)} + x^2} = \frac{-2}{-1} = 2.
 \end{aligned}$$

Demak, $y = -x + 2$ – og' ma asimptota.

$$4) y'' = \frac{1}{3} \times \frac{-9+12x-3x^2}{\sqrt[3]{(2-x)^2(x^2-4x+1)^2}} = \frac{-x^2+4x-3}{\sqrt[3]{(2-x)^2(x^2-4x+1)^2}}.$$

$$\begin{aligned}
 &\downarrow x=1, \\
 &\uparrow x=3
 \end{aligned}
 \text{ da } y''=0,$$

$$\begin{aligned}
 &\downarrow x=2, \\
 &\uparrow x=2 \pm \sqrt{3}.
 \end{aligned}
 \text{ da } y'' \text{ mavjud emas.}$$




$(1; -\sqrt[3]{2})$ – funksiyaning minimum nuqtasi

$(3; \sqrt[3]{2})$ – funksiyaning maksimum nuqtasi .

$$5) y'' = \frac{-(4x^4 - 16x^3 + 14x^2 - 8x + 10)}{\sqrt[3]{(2-x)^5(x^2-4x+1)^5}},$$

$$\begin{aligned}
 &\downarrow x = -0,94; \\
 &\uparrow x = 0,73
 \end{aligned}
 \text{ ad } y'' = 0,$$


$$\begin{aligned}
 &\downarrow x = 2, \\
 &\uparrow x = 2 \pm \sqrt{3}.
 \end{aligned}
 \text{ da } y'' \text{ mavjud emas.}$$



6) O'qlar bilan kesishish nuqtalarini topamiz:

$$x = 0 \text{ da } y = \sqrt[3]{2}.$$

$$y = 0 \text{ da } \begin{cases} x = 2, \\ x = 2 \pm \sqrt{3}. \end{cases}$$



$$1. y = -\sqrt[3]{(x+3)(x^3+6x+6)}.$$

$$3. y = \sqrt[3]{(x+1)(x^2+2x-2)}.$$

$$5. y = \sqrt[3]{(x-3)(x^2-6x+6)}.$$

$$7. y = \sqrt[3]{x^2(x+2)^2}.$$

$$9. y = \sqrt[3]{(x^2-2x-3)^2}.$$

$$11. y = \sqrt[3]{x^2(x-4)^2}.$$

$$13. y = \sqrt[3]{(x-1)(x+2)^2}.$$

$$15. y = \sqrt[3]{(x+6)x^2}.$$

$$17. y = \sqrt[3]{(x-1)^2} - \sqrt[3]{(x-2)^2}.$$

$$19. y = \sqrt[3]{(x-3)x^2}.$$

$$21. y = \sqrt[3]{(x+2)(x-4)^2}.$$

$$23. y = \sqrt[3]{x^2} - \sqrt[3]{(x-1)^2}.$$

$$25. y = \sqrt[3]{x(x+3)^2}.$$

$$2. y = \sqrt[3]{(x+2)(x^2+4x+1)}.$$

$$4. y = \sqrt[3]{(x-1)(x^2-2x-2)}.$$

$$6. y = \sqrt[3]{(x^2-4x+3)^2}.$$

$$8. y = \sqrt[3]{x^2(x-2)^2}.$$

$$10. y = \sqrt[3]{x^2(x+4)^2}.$$

$$12. y = \sqrt[3]{(x+3)x^2}.$$

$$14. y = \sqrt[3]{(x-1)^2} - \sqrt[3]{x^2}.$$

$$16. y = \sqrt[3]{(x-4)(x+2)^2}.$$

$$18. y = \sqrt[3]{(x+1)(x-2)^2}.$$

$$20. y = \sqrt[3]{(x-2)^2} - \sqrt[3]{(x-3)^2}.$$

$$22. y = \sqrt[3]{(x-6)x^2}.$$

$$24. y = \sqrt[3]{x(x-3)^2}.$$

$$26. y = \sqrt[3]{(x+2)^2} - \sqrt[3]{(x+3)^2}.$$

$$27. y = \sqrt[3]{x(x-6)^2}.$$

$$28. y = \sqrt[3]{x(x+6)^2}.$$

$$29. y = \sqrt[3]{(x+1)^2} - \sqrt[3]{(x+2)^2}.$$

$$30. y = \sqrt[3]{x(x-1)^2}.$$

Funksiyalarni tekshirish va grafiklarini chizish

9–masala. Quyidagi funktsiyani tekshiring va ularni grafigini yasang.

$$y = e^{\sin x + \cos x}.$$

1) $D(y) = (-\infty; +\infty)$.

2) Funktsiya juft ham, toq ham emas.

3. a) – vertikal asimptotalari yo'q.

6) og'ma asimptotalari yo'q.

4) davriy funktsiya

$$T = -\frac{\rho}{4} + \rho n, \quad n \in \mathbb{Z}.$$


$$5) y = e^{\sin x + \cos x}.$$

$$y = e^{\sqrt{2} \cos(x - \frac{\rho}{4})},$$

$$y' = -\sqrt{2} \sin(x - \frac{\rho}{4}) e^{\sqrt{2} \cos(x - \frac{\rho}{4})}$$

$$y' = 0, \text{ u holda } \sin(x - \frac{\rho}{4}) = 0,$$

$$x = \frac{\rho}{4} + \rho k, \quad k \in \mathbb{Z}.$$



$$\begin{aligned}
 y'' &= -\sqrt{2} \cos \frac{\pi}{4} x - \frac{\rho}{4} \ddot{\varphi} e^{\sqrt{2} \cos \frac{\pi}{4} x - \frac{\rho}{4} \ddot{\varphi}} + \\
 6) \quad &+ \sqrt{2} \sin \frac{\pi}{4} x - \frac{\rho}{4} \ddot{\varphi} \times \sqrt{2} \sin \frac{\pi}{4} x - \frac{\rho}{4} \ddot{\varphi} e^{\sqrt{2} \cos \frac{\pi}{4} x - \frac{\rho}{4} \ddot{\varphi}} = \\
 &= \sqrt{2} e^{\sqrt{2} \cos \frac{\pi}{4} x - \frac{\rho}{4} \ddot{\varphi}} \left(-\cos \frac{\pi}{4} x - \frac{\rho}{4} \ddot{\varphi} + \sqrt{2} \sin^2 \frac{\pi}{4} x - \frac{\rho}{4} \ddot{\varphi} \right) = \\
 &= \sqrt{2} e^{\sqrt{2} \cos \frac{\pi}{4} x - \frac{\rho}{4} \ddot{\varphi}} \left(-\cos \frac{\pi}{4} x - \frac{\rho}{4} \ddot{\varphi} - \sqrt{2} \cos^2 \frac{\pi}{4} x - \frac{\rho}{4} \ddot{\varphi} \right)
 \end{aligned}$$

$$y'' = 0 \text{ при } x = \pm \frac{\rho}{4} + \frac{\rho}{4} + 2pn, \quad n \in \mathbb{Z},$$

$$\frac{\rho}{4} x = \frac{\rho}{4} + \frac{\rho}{4} + 2pn, \quad n \in \mathbb{Z}.$$

$$\frac{\rho}{2} x = \frac{\rho}{2} + 2pn, \quad n \in \mathbb{Z}.$$

$$\frac{\rho}{4} x = -\frac{\rho}{4} + \frac{\rho}{4} + 2pk, \quad k \in \mathbb{Z}.$$


$$\frac{\rho}{2} x = 2pk, \quad k \in \mathbb{Z}.$$

$$x \in \left[\frac{\rho}{4} + 2pn; 2pk + \frac{\rho}{2} \right] \text{ da funksiya botiq, chunki } y'' > 0.$$

$$x \in \left[\frac{\rho}{2} + 2pn; 2pk + \frac{\rho}{2} \right] \text{ da funksiya qavariq, chunki } y'' < 0.$$

Egillish nuqtasi:

$$\left(2pk; e^{\frac{\rho}{2} + 2pn} \right), \left(\frac{\rho}{2} + 2pn; e^{\frac{\sqrt{2}}{2} \ddot{\varphi}} \right).$$



$$1. y = \arctg \frac{\sin x + \cos x}{\sqrt{2}}$$

$$3. y = \frac{1}{\sin x + \cos x}$$

$$5. y = \arctg \sin x.$$

$$7. y = \frac{1}{\sin x - \cos x}$$

$$9. y = \arctg \frac{\sin x - \cos x}{\sqrt{2}}$$

$$11. y = \frac{1}{(\sin x + \cos x)^2}.$$

$$13. y = -\arctg \cos x.$$

$$15. y = \frac{1}{(\sin x - \cos x)^2}.$$

$$17. y = \sqrt[3]{\sin x}.$$

$$19. y = \sqrt{\frac{\sin x - \cos x}{\sqrt{2}}}.$$

$$y = \sqrt[3]{\cos x}$$

$$y = \sqrt{\cos x}.$$

$$y = \sqrt[3]{\frac{\sin x + \cos x}{\sqrt{2}}}.$$

$$27. y = \sqrt{\sin x}.$$

$$29. y = \sqrt{\frac{\sin x + \cos x}{\sqrt{2}}}.$$

$$2. y = \ln(\cos x + \sin x).$$

$$4. y = e^{\sqrt{2} \sin x}.$$

$$6. y = \ln(\sqrt{2} \sin x).$$

$$8. y = e^{\sin x - \cos x}.$$

$$10. y = \ln(\sin x - \cos x).$$

$$12. y = e^{-\sqrt{2} \cos x}.$$

$$14. y = \ln(-\sqrt{2} \cos x).$$

$$16. y = e^{-\sin x - \cos x}.$$

$$18. y = \ln(-\sin x - \cos x).$$

$$20. y = e^{-\sqrt{2} \sin x}$$

21.

$$22. y = \ln(-\sqrt{2} \sin x).$$

23.

$$24. y = e^{\cos x - \sin x}.$$

25.

$$26. y = \ln(\cos x - \sin x).$$

$$28. y = e^{\sqrt{2} \cos x}$$

$$30. y = \ln(\sqrt{2} \cos x).$$

V BOB. ANIQMAS INTEGRAL

ANIQMAS INTEGRALNING TA'RIFI VA XOSSALARI

Berilgan $F(x)$ funksiyani differensiallashda uning $f(x) = F'(x)$ hosilasini topish talab qilinadi. Masalan: $F(x) = x^3$, $f(x) = 3x^2$. Teskari masalani ko'raylik: berilgan $f(x)$ hosilasi bo'yicha shunday $F(x)$ funksiyani topingki, uning hosilasi $f(x)$ ga teng, ya'ni $F'(x) = f(x)$ bo'lsin.

1- ta'rif. Hosilasi $f(x)$ ga teng bo'lgan $F(x)$ funksiya $f(x)$ funksiyaning boshlang'ich funksiyasi (boshlang'ichi) deyiladi.

1- misol. Berilgan: $f(x) = 3x^2$. $F(x)$ boshlang'ich funksiyani toping.

Echish. $F(x) = x^3$, chunki

$$F'(x) = (x^3)' = 3x^2.$$

2- misol. Berilgan: $f(x) = \frac{1}{2\sqrt{x}}$. $F(x)$ ni toping.

Echish. $F(x) = \sqrt{x}$, chunki $F'(x) = (\sqrt{x})' = \frac{1}{2\sqrt{x}}$.

Ravshanki, agar $F(x)$ funksiya $f(x)$ funksiyaning boshlang'ichi bo'lsa, u holda $F(x) + C$ ko'rinishdagi istalgan funksiya ham (bu yerda C — ixtiyoriy o'zgarmas) $f(x)$ ning boshlang'ich funksiyasi bo'ladi, chunki $[F(x) + C]' = f(x)$.

Masalan, agar $f(x) = x^2$ bo'lsa, u holda $F(x) = \frac{x^3}{3} + 2$; $F(x) = \frac{x^3}{3} - 5$;

$$F(x) = \frac{x^3}{3} + \ln 6, \text{ chunki } \left(\frac{x^3}{3} + 2\right)' = \frac{x^3}{3}' + 2' = \frac{x^3}{3}' - 5' = \frac{x^3}{3}' + \ln 6' = x^2.$$

2- ta'rif. Agar $F(x)$ funksiya $f(x)$ funksiyaning boshlang'ichi bo'lsa, u holda $F(x) + C$ ifoda $f(x)$ funksiyaning aniqmas integrali deyiladi: va quyidagicha belgilanadi:

$$\frac{d}{dx} \int f(x) dx.$$

Shunday qilib, ta'rifga ko'ra agar $F'(x) = f(x)$ bo'lsa, $\int f(x) dx = F(x) + C$.

Bu yerda $f(x)$ — integral ostidagi funksiya, $f(x)dx$ — integral ostidagi ifoda, \int — integral belgisi, x — integrallash o'zgaruvchisi.

$f(x)$ ning boshlang'ich funksiyasini topish amali funksiyani *integrallash* (integral olish) deyiladi.

Aniqmas integralning xossalari

Agar $F'(x) = f(x)$ bo'lsa, u holda

1. $\frac{d}{dx} \int f(x) dx = f(x)$.
2. $d \int f(x) dx = f(x) dx$.
3. $\int F'(x) dx = F(x) + C$.
4. $\int A f(x) dx = A \int f(x) dx$, bu yerda $A = const$.
5. $\int [f_1(x) \pm f_2(x)] dx = \int f_1(x) dx \pm \int f_2(x) dx$.

Bu xossalarning to'g'riligi differensiallash orqali tekshiriladi.

Integrallashning asosiy usullarini qarab chiqishdan avval asosiy integrallar jadvalini jiddiy kengaytiradigan bir muhim integrallash qoidasini ko'rib chiqamiz. Agar $\int f(x) dx = F(x) + C$ va $z = j(x)$ bo'lsa, u

holda

$$\int f(z)dx = F(z) + C. \quad (1)$$

Bu qoida. integrallash formulasining ko'rinishi integrallash o'zgaruvchisining xarakteriga bog'liq emasligini bildiradi. Bu qoidaning to'g'riligi (1) tenglikning har ikki tomonini differensiallash orqali oson tekshiriladi. Jumladan,

$$1. \int f(ax)dx = \frac{1}{a} \int f(ax)d(ax) = \frac{1}{a} F(ax) + C.$$

$$2. \int f(ax \pm b)dx = \frac{1}{a} \int f(ax \pm b)d(ax \pm b) = \frac{1}{a} F(ax \pm b) + C.$$

Masalan:

$$\int \sin 3xdx = \frac{1}{3} \int \sin 3xd(3x) = -\frac{1}{3} \cos 3x + C.$$

$$\int e^{\frac{x}{2}} dx = 2 \int e^{\frac{x}{2}} d\left(\frac{x}{2}\right) = 2e^{\frac{x}{2}} + C.$$

$$\int \frac{dx}{3x-5} = \frac{1}{3} \int \frac{d(3x-5)}{3x-5} = \frac{1}{3} \ln |3x-5| + C.$$

INTEGRALLASHNING ASOSIY USULLARI

Quyidagilar integrallashning asosiy usullari hisoblanadi:

1. Yoyib integrallash usuli.
2. Bevosita integrallash usuli.
3. O'rniga qo'yish usuli.
4. Bo'laklab integrallash usuli.

Yoyish (integral ostidagi ifodani yoyib integrallash) usuli

Agar $f(x) = f_1(x) \pm f_2(x)$ bo'lsa, u holda 1- § dagi 5- xossaga ko'ra yozish mumkin:

$$\int f(x)dx = \int [f_1(x) \pm f_2(x)]dx = \int f_1(x)dx \pm \int f_2(x)dx.$$

$$\begin{aligned}
1. \int (2 - 3\sqrt{x})^3 dx &= \int (8 - 36\sqrt{x}dx + 54xdx - 27\sqrt{x^3})dx = \\
&= 8\int dx - 36\int \sqrt{x}dx + 54\int xdx - 27\int \sqrt{x^3}dx = \\
&= 8x - 36 \times \frac{x^{\frac{3}{2}}}{\frac{3}{2}} + 54 \times \frac{x^2}{2} - 27 \times \frac{x^{\frac{5}{2}}}{\frac{5}{2}} + C = \\
&= 8x - 24x\sqrt{x} + 27x^2 - \frac{54}{5}x^2\sqrt{x} + C.
\end{aligned}$$

Shuni qayd qilib o'taymizki, har qaysi qo'shiluvchini integrallagandan so'ng ixtiyoriy o'zgarmasni yozish shart emas, chunki bu o'zgarmlarning yig'indisi yana o'zgarmas bo'lib, uni biz eng oxirida yozishimiz yetarli.

$$\begin{aligned}
\int \frac{x^3 - 2x^2 + 3x + 1}{x^2} dx &= \int \left(x - 2 + \frac{3}{x} + \frac{1}{x^2} \right) dx = \\
2. &= \int x dx - 2 \int dx + 3 \int \frac{dx}{x} + \int \frac{dx}{x^2} = \\
&= \frac{x^2}{2} - 2x + 3 \ln |x| - \frac{1}{x} + C. \\
3. \int \cos^2 \frac{x}{2} dx &= \frac{1}{2} \int (1 + \cos x) dx = \frac{1}{2} \int dx + \frac{1}{2} \int \cos x dx = \frac{1}{2} (x + \sin x) + C. \\
4. \int \operatorname{ctg}^2 x dx &= \int \frac{1}{\sin^2 x} - 1 dx = \int \frac{1}{\sin^2 x} dx - \int dx = -\operatorname{ctg} x - x + C.
\end{aligned}$$

Bu misollardan ko'rinadiki, yoyish usuli bilan integrallaganimizda integral ostidagi funksiyani elementar matematika vositalari yordamida shunday qo'shiluvchilarga yoyganimizda ulardan olingan integral jadvaldagi integraldan iborat bo'lsin.

Bevosita integrallash usuli

Bu usul asosida (1) qoida yotadi. Unga ko'ra aniqmas integrallarni

hisoblaganda integrallash o'zgaruvchisi x erkli o'zgaruvchi yoki $z = j(x)$ funksiyadan iborat bo'lishidan qat'i nazar 1- § da bayon qilingan 1 – 5 xossalarni va I — XVI jadval integrallarni tatbiq qilish mumkin.

Bu usulni tadbig'ini quyidagi misollarda ko'rsatamiz.

$$1. \int \frac{dx}{3-x} = - \int \frac{d(3-x)}{3-x} = - \ln |3-x| + C.$$

$$2. \int \frac{dx}{\cos^2 7x} = \frac{1}{7} \int \frac{d(7x)}{\cos^2 7x} = \frac{1}{7} \operatorname{tg} 7x + C.$$

$$\begin{aligned} \int (2-3x)^5 dx &= - \frac{1}{3} \int (2-3x)^5 d(2-3x) = \\ 3. \quad &= - \frac{1}{3} \times \frac{(2-3x)^6}{6} + C = - \frac{1}{18} (2-3x)^6 + C. \end{aligned}$$

$$4. \int \frac{dx}{1+9x^2} = \frac{1}{3} \int \frac{d(3x)}{1+(3x)^2} = \frac{1}{3} \operatorname{arctg} 3x + C.$$

Ko'pchilik hollarda dastlab integral ostidagi funksiyani yoyib olib, so'ngra bevosita integrallashni tatbiq qilishga to'g'ri keladi.

$$1. \int \operatorname{tg}^2(ax) dx = \int \frac{1}{\cos^2(ax)} - 1 dx = \frac{1}{a} \operatorname{tg}(ax) - x + C.$$

$$\begin{aligned} 2. \int \frac{2x-1}{2x+3} dx &= \int \frac{2x+3-4}{2x+3} dx = \int \frac{1}{2x+3} - \frac{4}{2x+3} dx = \\ &= x - 4 \int \frac{dx}{2x+3} = x - 2 \ln |2x+3| + C. \end{aligned}$$

$$\begin{aligned} 3. \int \frac{x^2+3}{x-2} dx &= \int \frac{x^2-4+7}{x-2} dx = \int \frac{x+2}{x-2} + \frac{7}{x-2} dx = \\ &= \frac{(x+2)^2}{2} + 7 \ln |x-2| + C. \end{aligned}$$

Integral ostidagi kasrning surati maxrajining differensialidan iborat bo'lsa, integral maxrajning logarifmiga teng bo'ladi.

$$4. \quad \int \frac{(x^3 - 1)dx}{x^4 - 4x + 1} = \frac{1}{4} \int \frac{d(x^4 - 4x + 1)}{x^4 - 4x + 1} = \frac{1}{4} \ln|x^4 - 4x + 1| + C.$$

$$5. \quad \int \frac{dx}{x(1 + \ln x)} = \int \frac{d(1 + \ln x)}{1 + \ln x} = \ln|1 + \ln x| + C.$$

$$6. \quad \int \frac{x^3 - 1}{x + 1} dx = \int \frac{x^3 + 1 - 2}{x + 1} dx = \int \frac{(x + 1)(x^2 - x + 1) - 2}{x + 1} dx =$$

$$= \int \frac{x^2 - x + 1}{x + 1} - \frac{2}{x + 1} dx = \frac{x^3}{3} - \frac{x^2}{3} + x - 2 \ln|x + 1| + C.$$

1-masala. Aniqmas integralni toping.

$$\int \frac{1 - \cos x}{(x - \sin x)^2} dx = \left| \begin{array}{l} x - \sin x = t \\ (1 - \cos x)dx = dt \end{array} \right| = \int \frac{dt}{t^2} = -t^{-1} + C = -\frac{1}{x - \sin x} + C.$$

1. $\int (4 - 3x)e^{-3x} dx.$
2. $\int \arctg \sqrt{4x - 1} dx.$
3. $\int (3x + 4)e^{3x} dx.$
4. $\int (4x - 2) \cos 2x dx.$
5. $\int (4 - 16x) \sin 4x dx.$
6. $\int (5x - 2)e^{3x} dx.$
7. $\int (1 - 6x)e^{2x} dx.$
8. $\int \ln(x^2 + 4) dx.$
9. $\int (2 - 4x) \sin 2x dx.$
10. $\int \arctg \sqrt{6x - 1} dx.$
11. $\int (4x - 3)e^{-2x} dx.$
12. $\int (2 - 9x)e^{-3x} dx.$
13. $\int \arctg \sqrt{2x - 1} dx.$
14. $\int \arctg \sqrt{3x - 1} dx.$
15. $\int \arctg \sqrt{5x - 1} dx.$
16. $\int (5x + 6) \cos 2x dx.$
17. $\int (3x - 2) \cos 5x dx.$
18. $\int (x\sqrt{2} - 3) \cos 2x dx.$
19. $\int (4x + 7) \cos 3x dx.$
20. $\int (2x - 5) \cos 4x dx.$
21. $\int (8 - 3x) \cos 5x dx.$
22. $\int (x + 5) \sin 3x dx.$
23. $\int (2 - 3x) \sin 2x dx.$
24. $\int (4x + 3) \sin 5x dx.$
25. $\int (7x - 10) \sin 4x dx.$
26. $\int (\sqrt{2} - 8x) \sin 3x dx.$
27. $\int \frac{x}{\cos^2 x} dx.$
28. $\int \frac{x}{\sin^2 x} dx.$
29. $\int x \sin^2 x dx.$
30. $\int \frac{x \cos x}{\sin^3 x} dx.$

Javoblar. 1.1 - $\ln \left| \frac{1 + \sqrt{x^2 + 1}}{x} \right| + C$; 1.2 $\ln |x| + \frac{1}{2} \ln^2 x + C$; 1.3 - $\arcsin \frac{1}{|x|} + C$;

1.4 $\frac{x^2}{2} + \ln^2 x + C$; 1.5 $\frac{1}{2} \ln \left| x^2 + \frac{1}{2} + \sqrt{x^4 + x^2 + 1} \right| + C$;

1.6 - $\frac{1}{4} \times (\arccos x)^4 + \arccos x + C$; 1.7 - $\frac{\ln^2 \cos x}{2} + C$; 1.8 $\frac{1}{2 \cos^2(x+1)} + C$;

1.9 $\frac{1}{2} \times \ln(x^2 + 1) + \frac{1}{2(x^2 + 1)} + C$; 1.10 - $\frac{1}{x - \sin x} + C$; 1.11 - $\frac{1}{x \times \sin x} + C$;

1.12 $\frac{1}{4} \times \ln |x^4 + 1| + \frac{1}{2} \times \operatorname{arctg} x^2 + C$; 1.13 $\frac{1}{2} \times \ln \left| x^2 - \frac{1}{2} + \sqrt{x^4 - x^2 - 1} \right| + C$;

1.14 $\frac{3}{5} \times \sqrt[3]{(x-1)^5} + \frac{3}{2} \times \sqrt[3]{(x-1)^2} + C$; 1.15 $\ln(x-1) + \frac{1}{2} \ln^2(x-1) + C$;

1.16 - $\frac{1}{12} \times \frac{1}{(x^3 + 3x + 1)^4} + C$; 1.17 $\frac{1}{2} (4 \operatorname{arctg}^2 x - \ln(1 + x^2)) + C$;

1.18 $\frac{1}{2} x^2 - 2 \ln(x^2 + 4) + C$; 1.19 $\frac{1}{2} \times \ln |x^2 + 2 \sin x| + C$;

1.20 - $\frac{1}{2 \times (2 \sin x - 3 \cos x)^2} + C$; 1.21 $\ln |1 + 4x^2| - \frac{1}{4} \times \operatorname{arctg}^2 2x + C$;

1.22 - $\frac{1}{\sqrt{x} + x} + C$; 1.23 $\frac{1}{2} \times \operatorname{arctg} x^2 + C$; 1.24 $\sqrt{x^2 + 1} - \ln \left| \frac{1 + \sqrt{x^2 + 1}}{x} \right| + C$;

1.25 $\sqrt{x^2 + 1} - \frac{1}{2} \ln \left| \frac{\sqrt{x^2 + 1} - 1}{\sqrt{x^2 + 1} + 1} \right| + C$; 1.26 $\frac{1}{2} (\operatorname{arctg}^2 x + \ln(1 + x^2)) + C$;

1.27 $\frac{1}{2} \times \ln |1 + x^2| - \frac{(\operatorname{arctg} x)^5}{5} + C$; 1.28 $\frac{1}{2} (x^2 - \ln(x^2 + 1)) + C$;

1.29 $\frac{1}{3} \times (\arcsin x)^3 + \arcsin x + C$; 1.30 $2 \operatorname{arctg} \sqrt{x} - \ln |x + 1| + C$.

Bo'laklab integrallash usuli.

Bo'laklab integrallash usuli quyidagi formula orqali topiladi:

$$\int u dv = uv - \int v du, \quad (2)$$

bunda, $u(x)$, $v(x)$ lar uzluksiz differensiallanuvchi funksiyalar.

(2) formula bo'laklab integrallash formulasi deyiladi.

Bo'laklab integrallashning mohiyati shundan iboratki, berilgan integralni hisoblashda integral ostidagi $f(x)dx$ ifodani $u dv$ ko'paytma shaklida tasvirlab va (2) formulani tatbiq qilib, berilgan $\int u dv$ integralni $\int v du$ jadval itegrali yoki osongina olinadigan integral bilan almashtiriladi.

2-masala. Aniqmas integralni toping.

$$\begin{aligned} \int \ln(4x^2 + 1) dx &= \left| \begin{array}{ll} u = \ln(4x^2 + 1) & dv = dx \\ du = \frac{8x}{4x^2 + 1} & v = x \end{array} \right| = x \ln(4x^2 + 1) - \int \frac{x^2}{4x^2 + 1} dx = \\ &= x \ln(4x^2 + 1) - \frac{1}{2} \int \frac{1}{4x^2 + 1} dx = x \ln(4x^2 + 1) - \frac{1}{2} \arctg 2x + C = \\ &= x \ln(4x^2 + 1) + \arctg 2x - 2x + C. \end{aligned}$$

Maple 7 dasturi yordamida

`Int(ln(4*x^2+1),x)=int(ln(4*x^2+1),x);`

$$\int \ln(4x^2 + 1) dx = x \ln(4x^2 + 1) - 2x + \arctan(2x)$$

tekshirib ko'ring.

1. $\int \frac{dx}{x\sqrt{x^2 + 1}}.$

2. $\int \frac{1 + \ln x}{x} dx.$

3. $\int \frac{dx}{x\sqrt{x^2 - 1}}.$

4. $\int \frac{x^2 + \ln x^2}{x} dx.$

5. $\int \frac{x}{\sqrt{x^4 + x^2 + 1}} dx.$

6. $\int \frac{\arccos^3 x - 1}{\sqrt{1 - x^2}} dx.$

7. $\int \operatorname{tg} x \ln \cos x dx.$ 8. $\int \frac{\operatorname{tg}(x+1)}{\cos^2(x+1)} dx.$ 9. $\int \frac{x^3}{(x^2+1)^2} dx.$
10. $\int \frac{\sin x - \cos x}{(\cos x + \sin x)^5} dx.$ 11. $\int \frac{x \cos x + \sin x}{(x \sin x)^2} dx.$ 12. $\int \frac{x^3 + x}{x^4 + 1} dx.$
13. $\int \frac{x}{\sqrt{x^4 - x^2 - 1}} dx.$ 14. $\int \frac{x}{\sqrt{x^2 - 1}} dx.$ 15. $\int \frac{1 + \ln(x-1)}{x-1} dx.$
16. $\int \frac{(x^2+1)dx}{(x^3+3x+1)^5}.$ 17. $\int \frac{4 \operatorname{arctg} x - x}{1+x^2} dx.$ 18. $\int \frac{x^3}{x^2+4} dx.$
19. $\int \frac{x + \cos x}{x^2 + 2 \sin x} dx.$ 20. $\int \frac{2 \cos x + 3 \sin x}{(2 \sin x - 3 \cos x)^3} dx.$ 21. $\int \frac{8x - \operatorname{arctg} 2x}{1+4x^2} dx.$
22. $\int \frac{\frac{1}{2\sqrt{x}} + 1}{(\sqrt{x} + x)^2} dx.$ 23. $\int \frac{x}{x^4 + 1} dx.$ 24. $\int \frac{x + \frac{1}{x}}{\sqrt{x^2 + 1}} dx.$
25. $\int \frac{x - \frac{1}{x}}{\sqrt{x^2 + 1}} dx.$ 26. $\int \frac{\operatorname{arctg} x + x}{1+x^2} dx.$ 27. $\int \frac{x - (\operatorname{arctg} x)^4}{1+x^2} dx.$
28. $\int \frac{x^3}{x^2 + 1} dx.$ 29. $\int \frac{(\arcsin x)^2 + 1}{\sqrt{1-x^2}} dx.$ 30. $\int \frac{1 - \sqrt{x}}{\sqrt{x}(x+1)} dx.$

Javoblar. 2.1 $(x-1)e^{-3x} + C$; 2.2 $x \operatorname{arctg} \sqrt{4x-1} - \frac{1}{4} \sqrt{4x-1} + C$;

2.3 $(x+1)e^{3x} + C$; 2.4 $(2x-1)\sin 2x + \cos 2x + C$;

2.5 $(4x-1)\cos 4x - \sin 4x + C$; 2.6 $\frac{1}{9} \sqrt{15x-11} e^{3x} + C$;

2.7 $(2-3x)e^{2x} + C$; 2.8 $x \ln(x^2+4) - 2x + 4 \operatorname{arctg} \frac{x}{2} + C$;

2.9 $(2x-1)\cos 2x - \sin 2x + C$; 2.10 $x \operatorname{arctg} \sqrt{6x-1} - \frac{1}{6} \sqrt{6x-1} + C$;

2.11 $\frac{1}{2} \sqrt{1-4x} e^{-2x} + C$; 2.12 $\frac{1}{3} \sqrt{1+9x} e^{-3x} + C$;

2.13 $x \operatorname{arctg} \sqrt{2x-1} - \frac{1}{2} \sqrt{2x-1} + C$; 2.14 $x \operatorname{arctg} \sqrt{3x-1} - \frac{1}{3} \sqrt{3x-1} + C$;

$$\begin{aligned}
& 2.15 \ x \times \arctg \sqrt{5x-1} - \frac{1}{5} \times \sqrt{5x-1} + C; \quad 2.16 \ \frac{1}{2}(5x+6)\sin 2x + \frac{5}{4}\cos 2x + C; \\
& 2.17 \ \frac{1}{5}(3x-2)\sin 5x + \frac{3}{25}\cos 5x + C; \quad 2.18 \ \frac{1}{2}(x\sqrt{2}-3)\sin 2x + \frac{\sqrt{2}}{4}\cos 2x + C; \\
& 2.19 \ \frac{1}{3}(4x+7)\sin 3x + \frac{4}{9}\cos 3x + C; \quad 2.20 \ \frac{1}{4}(2x-5)\sin 4x + \frac{1}{8}\cos 4x + C; \\
& 2.21 \ \frac{1}{5}(8-3x)\sin 5x - \frac{3}{25}\cos 5x + C; \quad 2.22 \ -\frac{1}{3}(x+5)\cos 3x + \frac{1}{9}\sin 3x + C; \\
& 2.23 \ \frac{1}{2}(3x-2)\cos 2x - \frac{3}{4}\sin 2x + C; \quad 2.24 \ -\frac{1}{5}(4x+3)\cos 5x + \frac{4}{25}\sin 5x + C; \\
& 2.25 \ -\frac{7}{4}x\cos 4x + \frac{7}{16}\sin 4x + \frac{5}{2}\cos 4x + C; \quad 2.26 \ \frac{1}{3}(8x-\sqrt{2})\cos 3x - \frac{8}{9}\sin 3x + C; \\
& 2.27 \ x \times \lg x + \ln|\cos x| + C; \quad 2.28 \ -x \times \operatorname{ctg} x + \ln|\sin x| + C; \\
& 2.29 \ -\frac{x}{5} \times \sin 2x - \frac{1}{8} \times \cos 2x + \frac{x^2}{4} + C; \quad 2.30 \ -\frac{x + \cos x \times \sin x}{2\sin^2 x} + C
\end{aligned}$$

Kvadrat uchhadni o'z ichiga olgan funksiyalarning integrallari

Kvadrat uchhadni o'z ichiga olgan

$$\int \frac{Ax+B}{ax^2+bx+c} dx; \quad \int \frac{Ax+B}{\sqrt{ax^2+bx+c}} dx; \quad \int \sqrt{ax^2+bx+c} dx$$

funksiyalarni integrallash jadvalidagi formulalarga keltirib integrallash uchun avvalo kvadrat uchhaddan to'liq kvadratni ajratib olish kerak bo'ladi. Bu holda ax^2+bx+c kvadrat uchhad quyidagi ko'rinishga keladi:

$$\begin{aligned}
ax^2+bx+c &= a\left(x^2+\frac{b}{a}x+\frac{c}{a}\right) = a\left(x^2+\frac{b}{a}x+\frac{b^2}{4a^2}\right) + \frac{b}{a} + \frac{c}{a} - \frac{b^2}{4a^2} = \\
&= a\left(x+\frac{b}{2a}\right)^2 \pm k^2
\end{aligned}$$

So'ngra almashtirishlar yo'li bilan yuqoridagi integrallarni integrallash jadvalidagi formulalarga keltirish mumkin.

Masalan,

$$\begin{aligned} \int \frac{dx}{x^2 + 5x + 7} &= \int \frac{dx}{x^2 + 2 \times \frac{5}{2}x + \frac{25}{4} - \frac{25}{4} + 7} = \\ 1. &= \int \frac{dx}{x^2 + \frac{5}{2}x + \frac{3}{4}} = \frac{2}{\sqrt{3}} \arctg \frac{x + \frac{5}{2}}{\frac{\sqrt{3}}{2}} + C = \\ &= \frac{2}{\sqrt{3}} \arctg \frac{2x + 5}{\sqrt{3}} + C. \\ \int \frac{3x + 1}{\sqrt{-x^2 + x + 2}} dx &= -\frac{3}{2} \int \frac{-2x - \frac{3}{2}}{\sqrt{2 + x - x^2}} dx = \\ 2. &= -\frac{3}{2} \int \frac{1 - 2x - \frac{5}{2}}{\sqrt{2 + x - x^2}} dx = -\frac{3}{2} \int \frac{d(2 + x - x^2)}{\sqrt{2 + x - x^2}} + \frac{5}{3} \int \frac{dx}{\sqrt{\frac{9}{4} - x - \frac{1}{2}x^2}} = \\ &= -3\sqrt{2 + x - x^2} + \frac{5}{2} \arcsin \frac{2x - 1}{3} + C. \\ \int \sqrt{x^2 - 2x - 1} dx &= \int \sqrt{(x - 1)^2 - 2} dx = \\ 3. &= \frac{1}{2} (x - 1) \sqrt{x^2 - 2x - 1} - \ln |x - 1 + \sqrt{x^2 - 2x - 1}| + C. \end{aligned}$$

3-masala. $\int \frac{1}{x^2 + x - 2} dx$ аниқмас интегрални ҳисоблайлик:

$$\begin{aligned} \int \frac{1}{x^2 + x - 2} dx &= \int \frac{1}{(x - 1)(x + 2)} dx = \frac{1}{3} \int \frac{dx}{x - 1} - \int \frac{dx}{x + 2} = \\ &= \frac{1}{3} \ln |x - 1| - \frac{1}{3} \ln |x + 2| + C = \frac{1}{3} \ln \left| \frac{x - 1}{x + 2} \right| + C. \end{aligned}$$

Maple 7 dasturi yordamida

> **Int(1/(x^2+x-2),x)=int(1/(x^2+x-2),x);**

$$\int \frac{1}{x^2 + x - 2} dx = -\frac{1}{3} \ln(x+2) + \frac{1}{3} \ln(x-1)$$

tekshirib ko'ring.

1. $\int \frac{dx}{x^2 + x + 5}$

2. $\int \frac{dx}{\sqrt{6 - 4x - 2x^2}}$

3. $\int \sqrt{x^2 + 8x + 25} dx$

4. $\int \frac{x+2}{x^2 + 2x + 3} dx$

5. $\int \frac{dx}{\sqrt{5 - 4x - x^2}}$

6. $\int \sqrt{8 + 2x - x^2} dx$

7. $\int \frac{5x-7}{x^2 + 3x + 8} dx$

8. $\int \frac{dx}{\sqrt{2 + 3x - 2x^2}}$

9. $\int \frac{x+7}{x^2 + 11x + 42} dx$

10. $\int \frac{dx}{x^2 - x + 14}$

11. $\int \frac{dx}{x^2 + 2x + 6}$

12. $\int \frac{x}{3x^2 - 8x + 9} dx$

13. $\int \frac{7x+4}{x^2 + x + 9} dx$

14. $\int \frac{dx}{\sqrt{3x^2 - 6x + 9}}$

15. $\int \sqrt{x^2 + 4x + 13} dx$

16. $\int \frac{3x-11}{x^2 + 8x + 18} dx$

17. $\int \frac{7x-8}{x^2 + 5x + 17} dx$

18. $\int \frac{2x-3}{x^2 + x + 5} dx$

19. $\int \frac{x+2}{x^2 - 2x - 3} dx$

20. $\int \frac{dx}{\sqrt{x^2 - 6x + 3}}$

21. $\int \sqrt{5 + 4x - x^2} dx$

22. $\int \frac{x+2}{x^2 + 2x - 3} dx$

23. $\int \frac{3x+4}{x^2 + 7x + 14} dx$

24. $\int \frac{x}{3x^2 + 4x + 5} dx$

25. $\int \frac{dx}{x^2 + 4x - 12}$

26. $\int \frac{dx}{\sqrt{3x^2 - 6x + 12}}$

27. $\int \frac{x-3}{x^2 - 9x + 23} dx$

28. $\int \frac{dx}{x^2 - 6x + 34}$

29. $\int \frac{x+7}{x^2 + 11x + 42} dx$

30. $\int \frac{dx}{x^2 + 6x + 34}$

Javoblar. 3.1 $\frac{2}{\sqrt{19}} \operatorname{arctg} \frac{2x+1}{\sqrt{19}} + C$. 3.2 $\frac{1}{\sqrt{2}} \arcsin \frac{x+1}{2} + C$.

3.3 $\frac{x+4}{2} \sqrt{x^2 + 8x + 25} + \frac{9}{2} \ln \left| x+4 + \sqrt{x^2 + 8x + 25} \right| + C$.

$$3.4 \frac{1}{2} \ln(x^2 + 2x + 3) + \frac{1}{\sqrt{2}} \operatorname{arctg} \frac{x+1}{\sqrt{2}} + C. \quad 3.5 \arcsin \frac{x+2}{3} + C. \quad 3.6$$

$$\frac{x-1}{2} \sqrt{8+2x-x^2} + \frac{9}{2} \arcsin \frac{x-1}{2} + C.$$

$$3.7 \frac{5}{2} \ln(x^2 + 3x + 8) - \frac{29}{\sqrt{23}} \operatorname{arctg} \frac{2x+3}{\sqrt{23}} + C. \quad 3.8 \frac{1}{\sqrt{2}} \arcsin \frac{4x-3}{5} + C.$$

$$3.9 \frac{1}{2} \ln(x^2 + 11x + 42) + \frac{3}{\sqrt{47}} \operatorname{arctg} \frac{2x+11}{\sqrt{47}} + C. \quad 3.10 \frac{2}{\sqrt{55}} \operatorname{arctg} \frac{2x-1}{\sqrt{55}} + C.$$

$$3.11 \frac{1}{\sqrt{5}} \operatorname{arctg} \frac{x+1}{\sqrt{5}} + C. \quad 3.12 \frac{1}{6} \ln(3x^2 - 8x + 9) + \frac{4}{\sqrt{11}} \operatorname{arctg} \frac{6x-8}{\sqrt{11}} + C.$$

$$3.13 \frac{7}{2} \ln(x^2 + x + 9) + \frac{1}{\sqrt{35}} \operatorname{arctg} \frac{2x+1}{\sqrt{35}} + C. \quad 3.14 \frac{1}{\sqrt{3}} \ln|x-1 + \sqrt{3x^2 - 6x + 9}| + C.$$

$$3.15 \frac{x+2}{2} \sqrt{x^2 + 4x + 13} + \frac{9}{2} \ln|x+2 + \sqrt{x^2 + 4x + 13}| + C.$$

$$3.16 \frac{3}{2} \ln(x^2 + 8x + 18) - \frac{23}{\sqrt{2}} \operatorname{arctg} \frac{x+4}{\sqrt{2}} + C.$$

$$3.17 \frac{7}{2} \ln(x^2 + 5x + 17) - \frac{51}{\sqrt{43}} \operatorname{arctg} \frac{2x+5}{\sqrt{43}} + C.$$

$$3.18 \ln(x^2 + x + 5) - \frac{8}{\sqrt{19}} \operatorname{arctg} \frac{2x+1}{\sqrt{19}} + C.$$

$$3.19 - \frac{1}{4} \ln|x+1| + \frac{5}{4} \ln|x-3| + C. \quad 3.20 \ln|x-3 + \sqrt{x^2 - 6x + 3}| + C.$$

$$3.21 \frac{x-2}{2} \sqrt{5+4x-x^2} + \frac{9}{2} \arcsin \frac{x-2}{2} + C. \quad 3.22 \frac{1}{4} \ln|x+3| + \frac{3}{4} \ln|x-1| + C.$$

$$3.23 \frac{3}{2} \ln(x^2 + 7x + 14) - \frac{13}{\sqrt{7}} \operatorname{arctg} \frac{2x+7}{\sqrt{7}} + C.$$

$$3.24 \frac{1}{6} \ln(3x^2 + 4x + 5) - \frac{2}{\sqrt{11}} \operatorname{arctg} \frac{6x+4}{\sqrt{11}} + C. \quad 3.25 \frac{1}{8} \ln \left| \frac{x-2}{x+6} \right| + C$$

$$3.26 \frac{1}{\sqrt{3}} \ln|x-1 + \sqrt{3x^2 - 6x + 12}| + C.$$

$$3.27 \frac{1}{2} \ln(x^2 - 9x + 23) + \frac{3}{\sqrt{11}} \operatorname{arctg} \frac{2x-9}{\sqrt{11}} + C.$$

$$3.28 \frac{1}{5} \operatorname{arctg} \frac{x-3}{5} + C. \quad 3.29 \frac{1}{2} \ln(x^2 + 11x + 42) + \frac{3}{\sqrt{47}} \operatorname{arctg} \frac{2x+11}{\sqrt{47}} + C.$$

$$3.30 \frac{1}{5} \operatorname{arctg} \frac{x+3}{5} + C.$$

Ratsional kasrlarni integrallash.

Ratsional kasr deb,

$$\frac{Q_m(x)}{P_n(x)} = \frac{b_0x^m + b_1x^{m-1} + \dots + b_m}{a_0x^n + a_1x^{n-1} + \dots + a_n},$$

Ko'rinishdagi kasrga aytiladi, bu yerda $P_n(x)$ va $Q_m(x)$ – darajalari mos ravishda n va m ga teng bo'lgan x ga nisbatan butun ko'phadlar.

Agar $n \geq m$ bo'lsa, ratsional kasr noto'g'ri, $n < m$ bo'lsa, to'g'ri kasr deyiladi.

Har qanday noto'g'ri kasrning suratini maxrajiga bo'lish natijasida butun qismini ajratib, uni biror ko'phad va to'g'ri kasr yig'indisi shaklida yozish mumkin:

$$\frac{Q_m(x)}{P_n(x)} = q(x) + \frac{Q_k(x)}{P_n(x)}, \quad k < n.$$

Masalan, $\frac{x^4 + 4}{x^2 + 3x - 1}$ noto'g'ri kasrning suratini maxrajiga bo'g'sak, quyidagiga ega bo'lamiz:

$$\frac{x^4 + 4}{x^2 + 3x - 1} = x^2 - 3x + 10 + \frac{-3x + 14}{x^2 + 3x - 1}.$$

4–masala. Aniqmas integralni toping.

$$\int \frac{x^3 - 3x^2 - 12}{(x - 4)(x - 3)(x - 2)} dx.$$

Kasrni bo'lamiz

$$\begin{array}{r|l} x^3 - 3x^2 - 12 & x^3 - 9x^2 + 26x - 24 \\ x^3 - 9x^2 + 26x - 24 & 1 \\ \hline 6x^2 - 26x + 12 & \end{array}$$

$$\int \frac{x^3 - 3x^2 - 12}{(x-4)(x-3)(x-2)} dx = \int \frac{1}{(x-4)(x-3)(x-2)} dx + \int \frac{6x^2 - 26x + 12}{(x-4)(x-3)(x-2)} dx$$

$\frac{6x^2 - 26x + 12}{(x-4)(x-3)(x-2)}$ to'g'ri kasrni sodda ratsional kasrlar yig'indisi

ko'rinishida yozamiz:

$$\begin{aligned} \frac{6x^2 - 26x + 12}{(x-4)(x-3)(x-2)} &= \frac{A}{x-4} + \frac{B}{x-3} + \frac{C}{x-2} = \\ &= \frac{A(x-3)(x-2) + B(x-4)(x-2) + C(x-4)(x-3)}{(x-4)(x-3)(x-2)}. \end{aligned}$$

$$A(x-3)(x-2) + B(x-4)(x-2) + C(x-4)(x-3) = 6x^2 - 26x + 12.$$

A, B, C noma'lum koeffitsiyentlarni topish uchun maxrajni nolga

aylantiradigan son qiymatlarni x ning o'rniga qo'yish bilan topamiz.

Odatda, bu usulni xususiy qiymatlar usuli deyiladi.

$$x = 4 \text{ da, } 2A = 4 \Rightarrow A = 2;$$

$$x = 3 \text{ da, } -B = -12 \Rightarrow B = 12;$$

$$x = 2 \text{ da, } 2C = -16 \Rightarrow C = -8;$$

Bundan

$$\int \frac{1}{(x-4)(x-3)(x-2)} dx = \int \frac{2}{x-4} + \frac{12}{x-3} - \frac{8}{x-2} dx =$$

$$= x + 2 \ln|x - 4| + 12 \ln|x - 3| - 8 \ln|x - 2| + C.$$

Maple 7 dasturi yordamida

> Int((1+(6*x^2-26*x+12)/((x-4)*(x-3)*(x-2))),x)=int((1+(6*x^2-26*x+12)/((x-4)*(x-3)*(x-2))),x);

$$\int 1 + \frac{6x^2 - 26x + 12}{(x-4)(x-3)(x-2)} dx = x + 2 \ln(x-4) + 12 \ln(x-3) - 8 \ln(x-2)$$

tekshirib ko'ring.

1. $\int \frac{x^3 + 1}{x^2 - x} dx.$

2. $\int \frac{3x^3 + 1}{x^2 - 1} dx.$

3. $\int \frac{x^3 - 17}{x^2 - 4x + 3} dx.$

4. $\int \frac{2x^3 + 5}{x^2 - x - 2} dx.$

5. $\int \frac{2x^3 - 1}{x^2 + x - 6} dx.$

6. $\int \frac{3x^3 + 25}{x^2 + 3x + 2} dx.$

7. $\int \frac{x^3 + 2x^2 + 3}{(x-1)(x-2)(x-3)} dx.$

8. $\int \frac{3x^3 + 2x^2 + 1}{(x+2)(x-2)(x-1)} dx.$

9. $\int \frac{x^3}{(x-1)(x+1)(x+2)} dx.$

10. $\int \frac{x^3 - 3x^2 - 12}{(x-4)(x-3)x} dx.$

11. $\int \frac{4x^3 + x^2 + 2}{x(x-1)(x-2)} dx.$

12. $\int \frac{3x^2 - 2}{x^3 - x} dx.$

13. $\int \frac{x^3 - 3x^2 - 12}{(x-4)(x-2)x} dx.$

14. $\int \frac{x^5 - x^3 + 1}{x^2 - x} dx.$

15. $\int \frac{x^5 + 3x^3 - 1}{x^2 + x} dx.$

16. $\int \frac{2x^5 - 8x^3 + 3}{x^2 - 2x} dx.$

17. $\int \frac{3x^5 - 12x^3 - 7}{x^2 + 2x} dx.$

18. $\int \frac{-x^5 + 9x^3 + 4}{x^2 + 3x} dx.$

19. $\int \frac{-x^5 + 25x^3 + 1}{x^2 + 5x} dx.$

20. $\int \frac{x^3 - 5x^2 + 5x + 23}{(x-1)(x+1)(x-5)} dx.$ 21.

$$\text{20. } \int \frac{x^5 + 2x^4 - 2x^3 + 5x^2 - 7x + 9}{(x-3)(x-1)x} dx.$$

$$\text{22. } \int \frac{2x^4 - 5x^2 - 8x - 8}{x(x-2)(x+2)} dx.$$

$$\text{23. } \int \frac{4x^4 + 2x^2 - x - 3}{x(x-1)(x+1)} dx.$$

$$\text{24. } \int \frac{3x^4 + 3x^3 - 5x^2 + 2}{x(x-1)(x+2)} dx.$$

$$\text{25. } \int \frac{2x^4 + 2x^3 - 41x^2 + 20}{x(x-4)(x+5)} dx.$$

$$\text{26. } \int \frac{x^5 - x^4 - 6x^3 + 13x + 6}{x(x-3)(x+2)} dx.$$

$$\text{27. } \int \frac{3x^3 - x^2 - 12x - 2}{x(x+1)(x-2)} dx.$$

$$\text{28. } \int \frac{2x^4 + 2x^3 - 3x^2 + 2x - 9}{x(x-1)(x+3)} dx.$$

$$\text{29. } \int \frac{2x^3 - x^2 - 7x - 12}{x(x-3)(x+1)} dx.$$

$$\text{30. } \int \frac{2x^3 - 40x - 8}{x(x+4)(x-2)} dx.$$

Javoblar. 4.1 $\frac{x^2}{2} + x - \ln|x| + 2\ln|x-1| + C;$

4.2 $\frac{3x^2}{2} + 2\ln|x-1| + \ln|x+1| + C;$

4.3 $\frac{x^2}{2} + 4x + 8\ln|x-1| + 5\ln|x-3| + C;$

4.4 $x^2 + 2x + 7\ln|x-2| - \ln|x+1| + C;$

4.5 $x^2 - 2x + 11\ln|x+3| + 3\ln|x-2| + C;$

4.6 $\frac{3x^2}{2} - 9x + 22\ln|x+1| - \ln|x+2| + C;$

4.7 $x + 3\ln|x-1| - 19\ln|x-2| + 24\ln|x-3| + C;$

4.8 $3x - 1,25\ln|x+2| + 8,25\ln|x-2| - 2\ln|x-1| + C;$

4.9 $x + \frac{1}{6}\ln|x-1| + \frac{1}{2}\ln|x+1| - \frac{3}{8}\ln|x+2| + C;$

4.10 $x + \ln|x-4| + 4\ln|x-3| - \ln|x| + C;$

4.11 $4x + \ln|x| - 7\ln|x-1| - 19\ln|x-2| + C;$

4.12 $3x + 2\ln|x| + \frac{1}{2}\ln|x-1| - \frac{5}{2}\ln|x+1| + C;$

4.13 $x + \frac{1}{2}\ln|x-4| + 4\ln|x-2| - \frac{3}{2}\ln|x| + C;$

$$4.14 \frac{x^4}{4} + \frac{x^3}{3} - \ln|x| + \ln|x-1| + C;$$

$$4.15 \frac{x^4}{4} - \frac{x^3}{3} + 2x^2 - 4x - \ln|x| + 5\ln|x+1| + C;$$

$$4.16 \frac{x^4}{2} + \frac{4x^3}{3} - \frac{3}{2}\ln|x| + \frac{3}{2}\ln|x-2| + C;$$

4.17

$$\frac{3x^4}{4} - 2x^3 - \frac{7}{2}\ln|x| + \frac{7}{2}\ln|x+2| + C;$$

$$4.18 - \frac{x^4}{4} + x^3 + \frac{4}{3}\ln|x| - \frac{4}{3}\ln|x+3| + C;$$

$$4.19 - \frac{x^4}{4} + \frac{5x^3}{3} + \frac{1}{5}\ln|x| - \frac{1}{5}\ln|x+5| + C;$$

$$4.20 x - 3 > \ln|x-1| + \ln|x+1| + 2 > \ln|x-5| + C;$$

$$4.21 \frac{x^3}{3} + x + 4 \times \ln|x+3| + 2 \times \ln|x-1| - 3 \times \ln|x| + C;$$

$$4.22 x^2 + 2 \times \ln|x| - \frac{3}{2} \times \ln|x-2| - \frac{5}{2} \times \ln|x+2| + C;$$

$$4.23 2x^2 + 3 \times \ln|x| + \ln|x-1| + 2\ln|x+1| + C;$$

$$4.24 \frac{3x^2}{2} - \ln|x| + \ln|x-1| + \ln|x+2| + C;$$

$$4.25 x^2 - \ln|x| + \frac{1}{9} \times \ln|x-4| - \frac{1}{9} \times \ln|x+5| + C;$$

$$4.26 \frac{x^3}{3} - \ln|x| + 3 \times \ln|x-3| - 2 \times \ln|x+2| + C;$$

$$4.27 3x + \ln|x| + 2 > \ln|x+1| - \ln|x-2| + C;$$

$$4.28 x^2 - 2x + 3 \times \ln|x| - \frac{3}{2} \times \ln|x-1| + \frac{11}{2} \times \ln|x+3| + C;$$

$$4.29 2x + 4 > \ln|x| + \ln|x-3| - 2 > \ln|x+1| + C;$$

$$4.30 2x + \ln|x| + \ln|x+4| - 6 > \ln|x-2| + C.$$

Noma'lum koeffitsiyentlar usuli.

Agar integral ostidagi kasr to'g'ri ($n > m$) bo'lsa, quyidagicha ish tutamiz:

1. $Q_m(x)$ maxrajni ko'paytuvchilarga yoyamiz. Aytaylik,

$$Q_m(x) = (x - a) \times (x - b)^a \times (x^2 + p_1x + q_1) \times (x^2 + p_2x + q_2)^b$$

bo'lsin, bu yerda a - sodda haqiqiy ildiz, b - karraligi a bo'lgan haqiqiy ildiz, $x^2 + p_1x + q_1$ va $x^2 + p_2x + q_2$ - okmpleks qo'shma ildizlarga ega bo'lgan uchhadlar.

Alebradagi teoremaga ko'ra to'g'ri ratsional $\frac{Q_m(x)}{P_n(x)}$ kasrni

quyidagicha yozish mumkin:

$$\begin{aligned} \frac{Q_m(x)}{P_n(x)} = & \frac{A}{x - a} + \frac{B_1}{x - b} + \frac{B_2}{(x - b)^2} + \dots + \frac{B_a}{(x - b)^a} + \frac{Mx + N}{x^2 + p_1x + q_1} + \\ & + \frac{M_1x + N_1}{x^2 + p_gx + q_g} + \frac{M_2x + N_2}{(x^2 + p_gx + q_g)^2} + \dots + \frac{M_bx + N_b}{(x^2 + p_gx + q_g)^b}. \end{aligned}$$

Masalan, agar $Q_m(x) = (x - 1)(x + 2)^3(x^2 + 2x + 2)(x^2 + 3x + 5)^2$ bo'lsa, u holda yoyilma quyidagicha bo'ladi:

$$\begin{aligned} \frac{Q_m(x)}{P_n(x)} = & \frac{A}{x - 1} + \frac{B_1}{x + 2} + \frac{B_2}{(x + 2)^2} + \frac{B_3}{(x + 2)^3} + \frac{Mx + N}{x^2 + 2x + 2} + \\ & + \frac{M_1x + N_1}{x^2 + 3x + 5} + \frac{M_2x + N_2}{(x^2 + 3x + 5)^2}. \end{aligned}$$

Bu yerda $A, B_1, B_2, B_3, M, N, M_1, N_1, M_2, N_2$ - hozircha noma'lum koeffitsiyentlar. Bu koeffitsiyentlarni topish uchun yoyilmaning o'ng

tomonini umumiy maxrajga keltiramiz va chap hamda o'ng tomondagi suratlarni aynan tenglaymiz. Hosil bo'lgan ayniyatda x ning o'ngdagi va chapdagi bir xil darajalari oldidagi koeffitsiyentlarni tenglab, noma'lum koeffitsiyentlarni topish uchun tenglamalar sistemasi tuziladi. Tenglamalar sistemasini yechish bilan noma'lum koeffitsiyentlar topiladi. Bu usulni odatda *noma'lum koeffitsiyentlar usuli* deyiladi.

5–masala. Aniqmas integralni toping.

$$\int \frac{x^3 + 6x^2 + 13x + 9}{(x+1)(x+2)^3} dx.$$

$\frac{x^3 + 6x^2 + 13x + 9}{(x+1)(x+2)^3}$ to'g'ri kasrni sodda ratsional kasrlar yig'indisi

ko'rinishida yozamiz:

$$\begin{aligned} \frac{x^3 + 6x^2 + 13x + 9}{(x+1)(x+2)^3} &= \frac{A}{x+1} + \frac{B_1}{x+2} + \frac{B_2}{(x+2)^2} + \frac{B_3}{(x+2)^3} = \\ &= \frac{A(x+2)^3 + B_1(x+1)(x+2)^2 + B_2(x+1)(x+2) + B_3(x+1)}{(x+1)(x+2)^3}. \end{aligned}$$

$$A(x+2)^3 + B_1(x+1)(x+2)^2 + B_2(x+1)(x+2) + B_3(x+1) = x^3 + 6x^2 + 13x + 9$$

$$x = -1 \text{ da, } A = 1;$$

$$x = -2 \text{ da, } -B_3 = -1 \Rightarrow B_3 = 1;$$

x ning bir xil darajalari oldidagi koeffitsiyentlarni tenglaymiz:

$$x^3: \quad A + B_1 = 1 \Rightarrow B_1 = 0;$$

$$x^0: \quad 8A + 4B_1 + 2B_2 + B_3 = 9 \Rightarrow B_2 = 0;$$

Demak,

$$\frac{x^3 + 6x^2 + 13x + 9}{(x+1)(x+2)^3} = \frac{1}{x+1} + \frac{1}{(x+2)^3}.$$

Bundan:

$$\int \frac{1}{x+1} + \frac{1}{(x+2)^3} dx = \ln|x+1| - \frac{1}{2(x+2)^2} + C.$$

Maple 7 dasturi yordamida

Int((x^3+6*x^2+13*x+9)/((x+1)*(x+2)^3),x)=int((x^3+6*x^2+13*x+9)/((x+1)*(x+2)^3),x);

$$\frac{x^3 + 6x^2 + 13x + 9}{(x+1)(x+2)^3} dx = \ln(x+1) - \frac{1}{2(x+2)^2}$$

tekshirib ko'ring.

1. $\int \frac{x^3 + 6x^2 + 13x + 8}{x(x+2)^3} dx.$

2. $\int \frac{x^3 - 6x^2 + 13x - 6}{(x+2)(x-2)^3} dx.$

3. $\int \frac{x^3 + 6x^2 + 14x + 10}{(x+1)(x+2)^3} dx.$

4. $\int \frac{x^3 - 6x^2 + 11x - 10}{(x+2)(x-2)^3} dx.$

5. $\int \frac{x^3 + 6x^2 + 11x + 7}{(x+1)(x+2)^3} dx.$

6. $\int \frac{2x^3 + 6x^2 + 7x + 1}{(x-1)(x+1)^3} dx.$

7. $\int \frac{x^3 + 6x^2 + 10x + 10}{(x-1)(x+2)^3} dx.$

8. $\int \frac{2x^3 + 6x^2 + 7x + 2}{x(x+1)^3} dx.$

9. $\int \frac{x^3 - 6x^2 + 13x - 8}{x(x-2)^3} dx.$

10. $\int \frac{x^3 - 6x^2 + 13x - 7}{(x+1)(x-2)^3} dx.$

11. $\int \frac{x^3 - 6x^2 + 14x - 6}{(x+1)(x-2)^3} dx.$

12. $\int \frac{x^3 - 6x^2 + 10x - 10}{(x+1)(x-2)^3} dx.$

13. $\int \frac{x^3 + x + 2}{(x+2)x^3} dx.$

14. $\int \frac{3x^3 + 9x^2 + 10x + 2}{(x-1)(x+1)^3} dx.$

$$15. \int \frac{2x^3 + x + 1}{(x+1)x^3} dx.$$

$$16. \int \frac{2x^3 + 6x^2 + 7x + 4}{(x+2)(x+1)^3} dx.$$

$$17. \int \frac{2x^3 + 6x^2 + 5x}{(x+2)(x+1)^3} dx.$$

$$18. \int \frac{2x^3 + 6x^2 + 7x}{(x-2)(x+1)^3} dx.$$

$$19. \int \frac{2x^3 + 6x^2 + 5x + 4}{(x-2)(x+1)^3} dx.$$

$$20. \int \frac{x^3 + 6x^2 + 4x + 24}{(x-2)(x+2)^3} dx.$$

$$21. \int \frac{x^3 + 6x^2 + 14x + 4}{(x-2)(x+2)^3} dx.$$

$$22. \int \frac{x^3 + 6x^2 + 18x - 4}{(x-2)(x+2)^3} dx.$$

$$23. \int \frac{x^3 + 6x^2 + 10x + 12}{(x-2)(x+2)^3} dx.$$

$$24. \int \frac{x^3 - 6x^2 + 14x - 4}{(x+2)(x-2)^3} dx.$$

$$25. \int \frac{x^3 + 6x^2 + 15x + 2}{(x-2)(x+2)^3} dx.$$

$$26. \int \frac{2x^3 - 6x^2 + 7x - 4}{(x-2)(x-1)^3} dx.$$

$$27. \int \frac{2x^3 - 6x^2 + 7x}{(x+2)(x-1)^3} dx.$$

$$28. \int \frac{x^3 + 6x^2 - 10x + 52}{(x-2)(x+2)^3} dx.$$

$$29. \int \frac{x^3 - 6x^2 + 13x - 6}{(x-2)(x+2)^3} dx.$$

$$30. \int \frac{x^3 + 6x^2 + 13x + 6}{(x-2)(x+2)^3} dx.$$

Javoblar. 5.1 $\ln |x| - \frac{1}{2(x+2)^2} + C$; 5.2 $\ln |x+2| - \frac{1}{2(x-2)^2} + C$;

5.3 $\ln |x+1| - \frac{1}{(x+2)^2} + C$; 5.4 $\ln |x+2| + \frac{1}{2(x-2)^2} + C$;

5.5 $\ln |x+1| + \frac{1}{2(x+2)^2} + C$; 5.6 $2\ln |x-1| - \frac{1}{2(x+1)^2} + C$;

5.7 $\ln |x-1| + \frac{1}{(x+2)^2} + C$; 5.8 $2\ln |x| - \frac{1}{2(x+1)^2} + C$;

5.9 $\ln |x| - \frac{1}{2(x-2)^2} + C$; 5.10 $\ln |x+1| - \frac{1}{2(x-2)^2} + C$;

$$5.11 \ln |x+1| - \frac{1}{(x-2)^2} + C; 5.12 \ln |x+1| + \frac{1}{(x-2)^2} + C;$$

$$5.13 \ln |x+2| - \frac{1}{2x^2} + C; 5.14 3 \ln |x-1| - \frac{1}{2(x+1)^2} + C;$$

$$5.15 2 \ln |x+1| - \frac{1}{2x^2} + C; 5.16 2 \ln |x+2| - \frac{1}{2(x+1)^2} + C;$$

$$5.17 2 \ln |x+2| + \frac{1}{2(x+1)^2} + C; 5.18 2 \ln |x-2| - \frac{1}{2(x+1)^2} + C;$$

$$5.19 2 \ln |x-2| + \frac{1}{2(x+1)^2} + C; 5.20 \ln |x-2| + \frac{4}{(x+2)^2} + C; 5.21$$

$$2 \ln |x-2| - \frac{1}{2(x+1)^2} + C; 5.22 \ln |x-2| - \frac{3}{(x+2)^2} + C;$$

$$5.23 \ln |x-2| + \frac{1}{(x+2)^2} + C; 5.24 \ln |x+2| - \frac{1}{(x-2)^2} + C;$$

$$5.25 \ln |x-2| - \frac{3}{2(x+2)^2} + C; 5.26 2 \ln |x-2| - \frac{1}{2(x-1)^2} + C;$$

$$5.27 2 \ln |x+2| - \frac{1}{2(x-1)^2} + C; 5.28 \ln |x-2| + \frac{11}{(x+2)^2} + C;$$

$$5.29 \frac{1}{16} \ln |x-2| + \frac{15}{16} \ln |x+2| + \frac{33x+34}{4(x+2)^2} + C;$$

$$5.30 \ln |x-2| - \frac{1}{2(x+2)^2} + C.$$

Maxrajining ildizlari kompleks va karrali.

6–masala. Aniqmas integralni toping.

$$\int \frac{x^3 + 5x^2 + 12x + 4}{(x+2)^2(x^2+4)} dx.$$

$\frac{x^3 + 5x^2 + 12x + 4}{(x+2)^2(x^2+4)}$ to'g'ri kasrni sodda ratsional kasrlar yig'indisi

ko'rinishida yozamiz:

$$\begin{aligned}\frac{x^3 + 5x^2 + 12x + 4}{(x+2)^2(x^2+4)} &= \frac{A}{x+2} + \frac{B}{(x+2)^2} + \frac{Cx+D}{x^2+4} = \\ &= \frac{A(x+2)(x^2+4) + B(x^2+4) + (Cx+D)(x+2)^2}{(x+2)^2(x^2+4)}.\end{aligned}$$

$$A(x+2)(x^2+4) + B(x^2+4) + (Cx+D)(x^2+4x+4) = x^3 + 5x^2 + 12x + 4.$$

$$x = -2 \text{ da, } 8B = -8 \Rightarrow B = -1;$$

x ning bir xil darajalari oldidagi koeffitsiyentlarni tenglaymiz:

$$x^3: \quad A + C = 1 \Rightarrow A = 0;$$

$$x: \quad 4A + 4C + 4D = 12 \Rightarrow C = 1;$$

$$x^0: \quad 8A + 4B + 4D = 4 \Rightarrow D = 2;$$

Bundan:

$$\begin{aligned}\int \frac{1}{(x+2)^2} + \frac{x+2}{x^2+4} dx &= \frac{1}{x+2} + \frac{1}{2} \int \frac{2x}{x^2+4} + 2 \int \frac{dx}{x^2+4} = \\ &= \frac{1}{x+2} + \frac{1}{2} \ln|x^2+4| + \arctg \frac{x}{2} + C.\end{aligned}$$

Maple 7 dasturi yordamida

Int((x^3+5*x^2+12*x+4)/((x+2)^2*(x^2+4)),x)=int((x^3+5*x^2+12*x+4)/((x+2)^2*(x^2+4)),x);

$$\int \frac{x^3 + 5x^2 + 12x + 4}{(x+2)^2(x^2+4)} dx = \frac{1}{x+2} + \frac{1}{2} \ln(x^2+4) + \arctan \frac{x}{2} + C$$

tekshirib ko'ring.

$$1. \int \frac{x^3 + 4x^2 + 4x + 2}{(x+1)^2(x^2+x+1)} dx.$$

$$2. \int \frac{x^3 + 4x^2 + 3x + 2}{(x+1)^2(x^2+1)} dx.$$

$$3. \int \frac{2x^3 + 7x^2 + 7x - 1}{(x+2)^2(x^2 + x + 1)} dx.$$

$$4. \int \frac{2x^3 + 4x^2 + 2x - 1}{(x+1)^2(x^2 + 2x + 2)} dx.$$

$$5. \int \frac{x^3 + 6x^2 + 9x + 6}{(x+1)^2(x^2 + 2x + 2)} dx.$$

$$6. \int \frac{2x^3 + 11x^2 + 16x + 10}{(x+2)^2(x^2 + 2x + 3)} dx.$$

$$7. \int \frac{3x^3 + 6x^2 + 5x - 1}{(x+1)^2(x^2 + 2)} dx.$$

$$8. \int \frac{x^3 + 9x^2 + 21x + 21}{(x+3)^2(x^2 + 3)} dx.$$

$$9. \int \frac{x^3 + 6x^2 + 8x + 8}{(x+2)^2(x^2 + 4)} dx.$$

$$10. \int \frac{2x^3 - 4x^2 - 16x - 12}{(x-1)^2(x^2 + 4x + 5)} dx.$$

$$11. \int \frac{-3x^3 + 13x^2 - 13x + 1}{(x-2)^2(x^2 - x + 1)} dx.$$

$$12. \int \frac{x^3 + 2x^2 + 10x}{(x+1)^2(x^2 - x + 1)} dx.$$

$$13. \int \frac{3x^3 + x + 46}{(x-1)^2(x^2 + 9)} dx.$$

$$14. \int \frac{4x^3 + 24x^2 + 20x - 28}{(x+3)^2(x^2 + 2x + 2)} dx.$$

$$15. \int \frac{2x^3 + 3x^2 + 3x + 2}{(x^2 + x + 1)(x^2 + 1)} dx.$$

$$16. \int \frac{x^3 + x + 1}{(x^2 + x + 1)(x^2 + 1)} dx.$$

$$17. \int \frac{x^2 + x + 3}{(x^2 + x + 1)(x^2 + 1)} dx.$$

$$18. \int \frac{2x^3 + 4x^2 + 2x + 2}{(x^2 + x + 1)(x^2 + x + 2)} dx.$$

$$19. \int \frac{2x^3 + 7x^2 + 7x + 9}{(x^2 + x + 1)(x^2 + x + 2)} dx.$$

$$20. \int \frac{4x^2 + 3x + 4}{(x^2 + 1)(x^2 + x + 1)} dx.$$

$$21. \int \frac{3x^3 + 4x^2 + 6x}{(x^2 + 2)(x^2 + 2x + 2)} dx.$$

$$22. \int \frac{2x^2 - x + 1}{(x^2 - x + 1)(x^2 + 1)} dx.$$

$$23. \int \frac{x^3 + x^2 + 1}{(x^2 + 1)(x^2 - x + 1)} dx.$$

$$24. \int \frac{x^3 + x + 1}{(x^2 + 1)(x^2 - x + 1)} dx.$$

$$25. \int \frac{2x^3 + 2x + 1}{(x^2 + 1)(x^2 - x + 1)} dx.$$

$$26. \int \frac{x^3 + 2x^2 + x + 1}{(x^2 + 1)(x^2 + x + 1)} dx.$$

$$27. \int \frac{x+4}{(x^2+2)(x^2+x+2)} dx.$$

$$28. \int \frac{2x^3+2x^2+2x+1}{(x^2+1)(x^2+x+1)} dx.$$

$$29. \int \frac{3x^3+7x^2+12x+6}{(x^2+x+3)(x^2+2x+3)} dx.$$

$$30. \int \frac{2x^3+3x^2+3x+2}{(x^2+1)(x^2+x+1)} dx.$$

Javoblar. 6.1 - $\frac{1}{x+1} + \frac{1}{2} \ln(x^2+x+1) + \frac{1}{\sqrt{3}} \arctg \frac{2x+1}{\sqrt{3}} + C;$

$$6.2 - \frac{1}{x+1} + \frac{1}{2} \ln|x^2+1| + \arctg x + C;$$

$$6.3 \frac{1}{x+2} + \ln(x^2+x+1) - \frac{2}{\sqrt{3}} \arctg \frac{2x+1}{\sqrt{3}} + C;$$

$$6.4 \frac{1}{x+1} + \ln(x^2+2x+2) - \arctg(x+1) + C;$$

$$6.5 - \frac{2}{x+1} + \frac{1}{2} \ln(x^2+2x+2) + \arctg(x+1) + C;$$

$$6.6 - \frac{2}{x+2} + \ln(x^2+2x+3) - \frac{1}{\sqrt{2}} \arctg \frac{x+1}{\sqrt{2}} + C;$$

$$6.7 \frac{1}{x+1} + \frac{3}{2} \ln(x^2+2) + \frac{1}{\sqrt{2}} \arctg \frac{x}{\sqrt{2}} + C;$$

$$6.8 - \frac{1}{x+1} + \frac{1}{2} \ln|x^2+3| + \frac{2}{\sqrt{3}} \arctg \frac{x}{\sqrt{3}} + C;$$

$$6.9 - \frac{1}{x+2} + \frac{1}{2} \ln(x^2+2) + \frac{1}{2} \arctg \frac{x}{2} + C;$$

$$6.6 \frac{3}{x-1} + \ln(x^2+4x+5) - \arctg(x+2) + C;$$

$$6.11 - \frac{1}{x-2} - \frac{3}{2} \ln(x^2-x+1) - \sqrt{3} \arctg \frac{2x-1}{\sqrt{3}} + C;$$

$$6.12 \frac{3}{x+1} + \frac{1}{2} \ln(x^2-x+1) + \frac{7}{\sqrt{3}} \arctg \frac{2x-1}{\sqrt{3}} + C;$$

$$6.13 - \frac{5}{x-1} + \frac{3}{2} \ln(x^2+9) + \frac{1}{3} \arctg \frac{x}{3} + C;$$

$$6.14 \quad -\frac{4}{x+3} + 2\ln(x^2 + 2x + 2) - 8\operatorname{arctg}(x+1) + C;$$

$$6.15 \quad \frac{1}{2} \times \ln(x^2 + x + 1) + \frac{1}{\sqrt{3}} \times \operatorname{arctg} \frac{x+1}{\sqrt{3}} + \frac{1}{2} \times \ln(x^2 + 1) + \operatorname{arctg} x + C;$$

$$6.16 \quad \ln(x^2 + x + 1) - \frac{1}{2} \times \ln(x^2 + 1) + C; \quad 6.17$$

$$\ln(x^2 + x + 1) + \frac{2}{\sqrt{3}} \times \operatorname{arctg} \frac{x+1}{\sqrt{3}} - \ln(x^2 + 1) + \operatorname{arctg} x + C; \quad 6.18$$

$$- \ln(x^2 + x + 1) + \frac{2}{\sqrt{3}} \times \operatorname{arctg} \frac{x+1}{\sqrt{3}} + 2\ln(x^2 + x + 2) + C;$$

$$6.19 \quad \frac{8}{\sqrt{3}} \times \operatorname{arctg} \frac{x+1}{\sqrt{3}} + \ln(x^2 + x + 2) + C;$$

$$6.20 \quad 3 \times \operatorname{arctg} x + \frac{2}{\sqrt{3}} \times \operatorname{arctg} \frac{x+1}{\sqrt{3}} + C;$$

$$6.21 \quad \ln(x^2 + 2) + \frac{1}{2} \times \ln(x^2 + 2x + 2) - \operatorname{arctg}(x+1) + C;$$

$$6.22 \quad \frac{1}{2} \times \ln(x^2 - x + 1) + \frac{1}{\sqrt{3}} \times \operatorname{arctg} \frac{x-1}{\sqrt{3}} - \frac{1}{2} \times \ln(x^2 + 1) + \operatorname{arctg} x + C;$$

$$6.23 \quad \frac{1}{2} \times \ln(x^2 - x + 1) + \frac{1}{\sqrt{3}} \times \operatorname{arctg} \frac{x-1}{\sqrt{3}} + \operatorname{arctg} x + C;$$

$$6.24 \quad \frac{2}{\sqrt{3}} \times \operatorname{arctg} \frac{x-1}{\sqrt{3}} + \frac{1}{2} \ln(x^2 + 1) + C;$$

$$6.25 \quad \frac{1}{2} \times \ln(x^2 - x + 1) - \frac{1}{2} \times \ln(x^2 + 1) + \sqrt{3} \operatorname{arctg} \frac{x-1}{\sqrt{3}} + C;$$

$$6.26 \quad \frac{2}{\sqrt{3}} \times \operatorname{arctg} \frac{x+1}{\sqrt{3}} + \frac{1}{2} \times \ln(x^2 + 1) + C;$$

$$6.27 \quad \ln(x^2 + x + 2) - \ln(x^2 + 2) + \frac{1}{\sqrt{2}} \times \operatorname{arctg} \frac{x}{\sqrt{2}} + C;$$

$$6.28 \quad \frac{1}{2} \times \ln(x^2 + x + 1) + \frac{1}{\sqrt{3}} \times \operatorname{arctg} \frac{x+1}{\sqrt{3}} + \frac{1}{2} \times \ln(x^2 + 1) + C;$$

$$6.29 \quad \ln(x^2 + x + 3) + \frac{1}{2} \times \ln(x^2 + x + 3) + C;$$

$$6.30 \quad \frac{1}{2} \ln(x^2 + x + 1) + \frac{1}{\sqrt{3}} \arctg \frac{2x+1}{\sqrt{3}} + \frac{1}{2} \ln(x^2 + 1) + \arctg x + C.$$

$\int x^m (a + bx^n)^p dx$ differensial binomlari integrali,

bu yerda a, b - o'zgarmas sonlar, m, n va p - ratsional sonlar.

$\int x^m (a + bx^n)^p dx$ ko'rinishdagi integrallarni hisoblash m, n va p

ratsional sonlarga bog'liqligini rus matematigi P.L.Chebyshev ko'rsatgan va uchta holdagina elementar funksiyalar orqali ifodalanadi.

1. Agar p - butun son bo'lsa, u holda integral $x = t^s$ o'rniga qo'yish yordamida (bunda s - kasrlar maxrajining m va n ning eng kichik umumiy karralisi) ratsional funksiya integraliga keltiriladi.

$\int \sqrt[3]{x}(2 + \sqrt{x})^2 dx$ integralda $p = 2$ - butun son. Hisoblaymiz

$$\begin{aligned} \int x^{\frac{1}{3}}(x + x^{\frac{1}{2}})dx &= \left| \begin{array}{l} m = \frac{1}{3}; \quad n = \frac{1}{2} \\ x = t^6, \quad dx = 6t^5 dt \end{array} \right| = \int t^2(2 + t^3)^2 6t^5 dt = \\ &= 6 \int (4t^7 + 4t^{10} + t^{13})dt = 6 \left(\frac{4}{8}t^8 + \frac{4}{11}t^{11} + \frac{1}{14}t^{14} \right) + C = \\ &= \left| t = \sqrt[6]{x} \right| = 3\sqrt[3]{x^4} + \frac{24}{11}\sqrt[6]{x^{11}} + \frac{3}{7}\sqrt[3]{x^7} + C. \end{aligned}$$

2. Agar $\frac{m+1}{n}$ - butun son bo'lsa, u holda integral $a + bx^n = t^s$ o'rniga

qo'yish bilan ratsional funksiya integrallanadi, bunda s son p kasrning maxraji.

$\int x^5 \sqrt[3]{(1+x^3)^2} dx$ integralda $m = 5, n = 3, p = \frac{2}{3}; \frac{m+1}{n} = \frac{5+1}{3} = 2$ - butun

son. $s = 3$, ya'ni p kasrning maxraji.

Tegishli o'rniga qo'yishdan, $1 + x^3 = t^3, x = (t^3 - 1)^{\frac{1}{3}}, dx = t^2(t^3 - 1)^{-\frac{2}{3}} dt$.

Demak,

$$\int (t^3 - 1)^{\frac{5}{3}} \cdot 2 \cdot (t^3 - 1)^{-\frac{2}{3}} dt = \int (t^3 - 1) t^4 dt = \int (t^7 - t^4) dt =$$

$$= \frac{t^8}{8} - \frac{t^5}{5} + C = \frac{(t^3 - 1)^{\frac{5}{3}}}{8} - \frac{1}{5} + C.$$

x o'zgaruvchiga qaytib, uzil-kesil topamiz:

$$\int x^5 \sqrt[3]{(1+x^3)^2} dx = (1+x^3)^{\frac{5}{3}} \frac{1+x^3}{8} - \frac{1}{5} + C.$$

3. $\frac{m+1}{n} + p$ –butun son bo'lganda $ax^{-n} + b = t^s$ o'rniga qo'yish bilan, bu yerda s son p kasrning maxraji.

7–masala. Aniqmas integralni toping.

$$\int \frac{\sqrt[5]{1+\sqrt[3]{x}}}{x\sqrt[5]{x^2}} dx.$$

Berilgan integralni $\int \frac{\sqrt[5]{1+\sqrt[3]{x}}}{x\sqrt[5]{x^2}} dx = \int x^{-\frac{7}{5}} (1+x^{\frac{1}{3}})^{\frac{1}{5}} dx$ ko'rinishda yozib

olamiz. Integral ositidagi ifoda $x^m (a+bx^n)^p$ ekanligidan

$$m = -\frac{7}{5}, \quad n = \frac{1}{3}, \quad p = \frac{1}{5}, \quad \frac{m+1}{n} + p = -1.$$

$x^{-\frac{1}{3}} + 1 = t^5$ o'rniga qo'yishdan foydalanamiz.

$$x = (t^5 - 1)^{-3}, \quad dx = -3(t^5 - 1)^{-4} 5t^4 dt \quad \text{ga egamiz.}$$

$$\int x^{-\frac{7}{5}} (1+x^{\frac{1}{3}})^{\frac{1}{5}} dx = \int (t^5 - 1)^{\frac{21}{5}} (1 + (t^5 - 1)^{-1})^{\frac{1}{5}} \frac{(-15)t^4}{(t^5 - 1)^4} dt =$$

$$= -\frac{15}{6} t^6 + C = -\frac{5}{2} (x^{-\frac{1}{3}} + 1)^{\frac{6}{5}} + C = -\frac{5}{2} \frac{1 + \sqrt[3]{x}}{\sqrt[3]{x}} + C.$$

$$1. \int \frac{\sqrt{1+\sqrt{x}}}{x\sqrt[4]{x^3}} dx.$$

$$2. \int \frac{\sqrt[3]{1+\sqrt{x}}}{x\sqrt[3]{x^2}} dx.$$

$$3. \int \frac{\sqrt{1+\sqrt[3]{x}}}{x\sqrt{x}} dx.$$

$$4. \int \frac{\sqrt[3]{1+\sqrt[3]{x}}}{x\sqrt[9]{x^4}} dx.$$

$$5. \int \frac{\sqrt[3]{1+\sqrt[3]{x^2}}}{x\sqrt[9]{x^8}} dx.$$

$$6. \int \frac{\sqrt[3]{(1+\sqrt[3]{x})^2}}{x\sqrt[9]{x^5}} dx.$$

$$7. \int \frac{\sqrt[3]{(1+\sqrt[3]{x^2})^2}}{x^2\sqrt[9]{x}} dx.$$

$$8. \int \frac{\sqrt{1+\sqrt[3]{x^2}}}{x^2} dx.$$

$$9. \int \frac{\sqrt{1+x}}{x^2\sqrt{x}} dx.$$

$$10. \int \frac{\sqrt[4]{(1+\sqrt{x})^3}}{x\sqrt[8]{x^7}} dx.$$

$$11. \int \frac{\sqrt[4]{(1+\sqrt[3]{x})^3}}{x\sqrt[12]{x^7}} dx.$$

$$12. \int \frac{\sqrt[4]{(1+\sqrt[3]{x^2})^3}}{x^2\sqrt[6]{x}} dx.$$

$$13. \int \frac{\sqrt{1+\sqrt[4]{x^3}}}{x^2\sqrt[8]{x}} dx.$$

$$14. \int \frac{\sqrt[3]{1+\sqrt[4]{x^3}}}{x^2} dx.$$

$$15. \int \frac{\sqrt[3]{(1+\sqrt[4]{x^3})^2}}{x^2\sqrt[4]{x}} dx.$$

$$16. \int \frac{\sqrt[5]{(1+\sqrt{x})^4}}{x\sqrt[10]{x^9}} dx.$$

$$17. \int \frac{\sqrt[5]{(1+\sqrt[3]{x})^4}}{x\sqrt[5]{x^3}} dx.$$

$$18. \int \frac{\sqrt[5]{(1+\sqrt[3]{x^2})^4}}{x^2\sqrt[5]{x}} dx.$$

$$19. \int \frac{\sqrt[5]{(1+\sqrt[4]{x^3})^4}}{x^2\sqrt[20]{x^7}} dx.$$

$$20. \int \frac{\sqrt[5]{(1+\sqrt[5]{x^4})}}{x^2\sqrt[25]{x^{11}}} dx.$$

$$21. \int \frac{\sqrt{1+\sqrt[5]{x^4}}}{x^2\sqrt[5]{x}} dx.$$

$$22. \int \frac{\sqrt[3]{1+\sqrt[5]{x^4}}}{x^2\sqrt[15]{x}} dx.$$

$$23. \int \frac{\sqrt[3]{(1+\sqrt[5]{x^4})^2}}{x^2\sqrt[3]{x}} dx.$$

$$24. \int \frac{\sqrt[4]{(1+\sqrt[5]{x^4})^3}}{x^2\sqrt[5]{x^2}} dx.$$

$$25. \int \frac{\sqrt[3]{1+\sqrt[4]{x}}}{x\sqrt[3]{x}} dx.$$

$$26. \int \frac{\sqrt[3]{(1+\sqrt[4]{x})^2}}{x\sqrt[12]{x^5}} dx.$$

$$27. \int \frac{\sqrt[4]{1+\sqrt[3]{x}}}{x\sqrt[12]{x^5}} dx.$$

$$28. \int \frac{\sqrt[4]{1+\sqrt[3]{x^2}}}{x\sqrt[6]{x^5}} dx.$$

$$29. \int \frac{\sqrt[3]{1+\sqrt[5]{x}}}{x\sqrt[15]{x^4}} dx.$$

$$30. \int \frac{\sqrt[3]{(1+\sqrt{x})^2}}{x\sqrt[6]{x^5}} dx.$$

Javoblar. 7.1 - $\frac{4\sqrt{(1+\sqrt{x})^3}}{3\sqrt[4]{x}} + C$; 7.2 - $\frac{3\sqrt[3]{(1+\sqrt{x})^4}}{2\sqrt[3]{x^2}} + C$; 7.3 - $2\sqrt{\frac{(1+\sqrt[3]{x})^3}{x}} + C$;

$$\begin{aligned}
7.4 - \frac{9}{4} \frac{\sqrt[3]{1+\sqrt[3]{x}}}{\sqrt[3]{x}} \frac{\ddot{\circ}^4}{\circ} + C; 7.5 - \frac{9}{8} \frac{\sqrt[3]{1+\sqrt[3]{x^2}}}{\sqrt[3]{x^2}} \frac{\ddot{\circ}^4}{\circ} + C; 7.6 - \frac{9}{5} \frac{\sqrt[3]{1+\sqrt[3]{x}}}{\sqrt[3]{x}} \frac{\ddot{\circ}^5}{\circ} + C; \\
7.7 - \frac{9}{10} \frac{\sqrt[3]{1+\sqrt[3]{x^2}}}{\sqrt[3]{x^2}} \frac{\ddot{\circ}^5}{\circ} + C; 7.8 - \frac{\sqrt{(1+\sqrt[3]{x^2})^3}}{x} + C; 7.9 - \frac{2}{3} \frac{\sqrt[3]{1+x}}{x} \frac{\ddot{\circ}^3}{\circ} + C; \\
7.10 - \frac{8}{7} \frac{\sqrt[3]{1+\sqrt{x}}}{\sqrt{x}} \frac{\ddot{\circ}^7}{\circ} + C; 7.11 - \frac{12}{7} \frac{\sqrt[3]{1+\sqrt[3]{x}}}{\sqrt[3]{x}} \frac{\ddot{\circ}^7}{\circ} + C; 7.12 - \frac{6 \sqrt[4]{(1+\sqrt[3]{x^2})^7}}{7 \sqrt[6]{x^7}} + C; \\
7.13 - \frac{8}{9} \frac{\sqrt[3]{1+\sqrt[4]{x^3}}}{\sqrt[4]{x^3}} \frac{\ddot{\circ}^3}{\circ} + C; 7.14 - \frac{\sqrt[3]{(1+\sqrt[4]{x^3})^4}}{x} + C; 7.15 - \frac{4}{5} \frac{\sqrt[3]{x}}{x}^{-\frac{3}{4}} + 1 \frac{\ddot{\circ}^{\frac{5}{3}}}{\circ} + C; 7.16 \\
- \frac{10}{9} \frac{\sqrt[3]{1+\sqrt{x}}}{\sqrt{x}} \frac{\ddot{\circ}^9}{\circ} + C; 7.17 - \frac{5}{3} \frac{\sqrt[3]{1+\sqrt[3]{x}}}{\sqrt[3]{x}} \frac{\ddot{\circ}^9}{\circ} + C; 7.18 - \frac{5}{6} \frac{\sqrt[3]{1+\sqrt[3]{x^2}}}{\sqrt[3]{x^2}} \frac{\ddot{\circ}^9}{\circ} + C; 7.19 \\
- \frac{20}{27} \frac{\sqrt[3]{1+\sqrt[4]{x^3}}}{\sqrt[4]{x^3}} \frac{\ddot{\circ}^9}{\circ} + C; 7.20 - \frac{25}{36} \frac{\sqrt[3]{1+\sqrt[5]{x^4}}}{\sqrt[5]{x^4}} \frac{\ddot{\circ}^9}{\circ} + C; \\
7.21 - \frac{5}{6} \frac{\sqrt{(1+\sqrt[5]{x^4})^3}}{x \sqrt[5]{x}} + C; 7.22 - \frac{15(1+\sqrt[5]{x^4})^{\frac{4}{3}}}{16 \sqrt[15]{x^{16}}} + C; 7.23 - \frac{3(1+\sqrt[5]{x^4})^{\frac{5}{3}}}{4 \sqrt[3]{x^4}} + C; \\
7.24 - \frac{5}{7} \frac{\sqrt[3]{x}}{x}^{-\frac{4}{5}} + 1 \frac{\ddot{\circ}^{\frac{7}{4}}}{\circ} + C; 7.25 - \frac{1}{3} \frac{\sqrt[3]{x}}{x}^{-\frac{1}{4}} + 1 \frac{\ddot{\circ}^{\frac{4}{3}}}{\circ} + C; 7.26 - \frac{12}{5} \frac{\sqrt[3]{1+\sqrt[4]{x}}}{\sqrt[4]{x}} \frac{\ddot{\circ}^5}{\circ} + C; \\
7.27 - \frac{12}{5} \frac{\sqrt[3]{1+\sqrt[3]{x}}}{\sqrt[3]{x}} \frac{\ddot{\circ}^5}{\circ} + C; 7.28 - \frac{6 \sqrt[4]{(1+\sqrt[3]{x^2})^5}}{5 \sqrt[6]{x^5}} + C; 7.29 - \frac{15(1+\sqrt{x})^{\frac{4}{3}}}{4 \sqrt[15]{x^4}} + C; \\
7.30 - \frac{6}{5} \frac{\sqrt[3]{1+\frac{1}{\sqrt{x}}}}{\sqrt{x}} \frac{\ddot{\circ}^5}{\circ} + C.
\end{aligned}$$

VI BOB. Aniq integral.

Aniq integral mavzusida Nyuton–Leybnits formulasi bilan tanishasiz va uni aniq integrallarni yechishdagi tatbiqi, hamda boshlang'ich funksiyalarni topishda qo'llaysiz. Undan tashqari geometrik masalalarni aniq integrallar yordamida yassi figuralarning yuzi, egri chiziqli yoyi

uzunligi va jismning hajmini topish bilan bog'liq masalalarni o'zlashtirasiz.

Agar $F(x)$ funksiya $[a, b]$ da uzluksiz $f(x)$ funksiyaning boshlang'ichi bo'lsa, u holda quyidagi formula o'rinlidir:

$$\int_a^b f(x) dx = F(x) \Big|_a^b = F(b) - F(a). \quad (1)$$

Nyuton–leybnits formulasi deb ataluvchi (1) formula aniqmas integral bilan aniq integral o'rtasidagi bog'lanishni ifodalaydi.

Aniq integralda o'zgaruvchini almashtirish.

Bu holda ushbu formula o'rinli bo'ladi:

$$\int_a^b f(x) dx = \int_{j(a)}^{j(b)} f(j(t)) j'(t) dt,$$

bu yerda $j(t)$ va $j'(t)$ lar, $[a, b]$ kesmada uzluksiz funksiyalar, $a = j(a)$, $b = j(b)$.

Masalan, $\int_3^8 \frac{x dx}{\sqrt{1+x}}$ integralni topish talab etilsin. $\sqrt{1+x} = t$ belgilash

kiritamiz, u holda $1+x = t^2$, $dx = 2t dt$. Yangi o'zgaruvchi t ning o'zgarish chegaralarini topamiz: x o'zgaruvchi $[3, 8]$ intervalda o'zgaradi. $x = 3$ da $t = \sqrt{1+3} = 2$; $x = 8$ da $t = \sqrt{1+8} = 3$.

Demak,

$$\int_3^8 \frac{x dx}{\sqrt{1+x}} = \int_2^3 \frac{(t^2 - 1) 2t dt}{t} = 2 \int_2^3 (t^2 - 1) dt = 2 \left[\frac{t^3}{3} - t \right]_2^3 = \frac{32}{3}.$$

1–masala. Aniq integralni hisoblang.

$$\begin{aligned} \int_0^{1/2} \frac{8x - \operatorname{arctg} 2x}{1+4x^2} dx &= \int_0^{1/2} \frac{8x}{1+4x^2} dx - \int_0^{1/2} \operatorname{arctg} 2x d(\operatorname{arctg} 2x) = \\ &= \ln|1+4x^2| \Big|_0^{1/2} - \frac{1}{2} \operatorname{arctg}^2 2x \Big|_0^{1/2} = \ln 2 - 0 - \frac{1}{2} \times \frac{\rho^2}{16} + 0 = \ln 2 - \frac{\rho^2}{32}. \end{aligned}$$

$$1. \int_{e+1}^{e^2+1} \frac{1 + \ln(x-1)}{x-1} dx.$$

$$2. \int_0^1 \frac{(x^2+1)dx}{(x^3+3x+1)^2}.$$

$$3. \int_0^1 \frac{4\operatorname{arctg} x - x}{1+x^2} dx.$$

$$4. \int_0^2 \frac{x^3 dx}{x^2+4}.$$

$$5. \int_{\rho}^{2\rho} \frac{x + \cos x}{x^2 + 2 \sin x} dx.$$

$$6. \int_0^{\rho/4} \frac{2 \cos x + 3 \sin x}{(2 \sin x - 3 \cos x)^3} dx.$$

$$7. \int_1^4 \frac{\frac{1}{2\sqrt{x}} + 1}{(\sqrt{x} + x)^2} dx.$$

$$8. \int_0^1 \frac{x dx}{x^4 + 1}.$$

$$9. \int_{\sqrt{3}}^{\sqrt{8}} \frac{x + \frac{1}{x}}{\sqrt{x^2 + 1}} dx.$$

$$10. \int_{\sqrt{3}}^{\sqrt{8}} \frac{x - \frac{1}{x}}{\sqrt{x^2 + 1}} dx.$$

$$11. \int_0^{\sqrt{3}} \frac{\operatorname{arctg} x + x}{1+x^2} dx.$$

$$12. \int_0^{\sqrt{3}} \frac{x - (\operatorname{arctg} x)^4}{1+x^2} dx.$$

$$13. \int_0^1 \frac{x^3}{x^2+1} dx.$$

$$14. \int_0^{\sin 1} \frac{(\arcsin x)^2 + 1}{\sqrt{1-x^2}} dx.$$

$$15. \int_1^3 \frac{1 - \sqrt{x}}{\sqrt{x}(x+1)} dx.$$

$$16. \int_{\sqrt{3}}^{\sqrt{8}} \frac{dx}{x\sqrt{x^2+1}}.$$

$$17. \int_1^e \frac{1 + \ln x}{x} dx.$$

$$18. \int_{\sqrt{2}}^2 \frac{dx}{x\sqrt{x^2-1}}.$$

$$19. \int_1^e \frac{x^2 + \ln x^2}{x} dx.$$

$$20. \int_0^1 \frac{x}{\sqrt{x^4+x^2+1}} dx.$$

$$21. \int_0^1 \frac{x^3 dx}{(x^2 + 1)^2}.$$

$$23. \int_{-1}^0 \frac{\operatorname{tg}(x+1)}{\cos^2(x+1)} dx.$$

$$25. \int_p^{2p} \frac{1 - \cos x}{(x - \sin x)^2} dx.$$

$$27. \int_{p/4}^{p/2} \frac{x \cos x + \sin x}{(x \sin x)^2} dx.$$

$$29. \int_{\sqrt{2}}^{\sqrt{3}} \frac{xdx}{\sqrt{x^4 - x^2 - 1}}.$$

$$22. \int_0^{p/4} \operatorname{tg} x \ln \cos x dx.$$

$$24. \int_0^{1/\sqrt{2}} \frac{(\arccos x)^3 - 1}{\sqrt{1 - x^2}} dx.$$

$$26. \int_0^{p/4} \frac{\sin x - \cos x}{(x - \sin x)^2} dx.$$

$$28. \int_0^1 \frac{x^3 + x}{x^4 + 1} dx.$$

$$30. \int_2^9 \frac{xdx}{\sqrt[3]{x-1}}.$$

Javoblar. 1.1 $2\frac{1}{2}$; 1.2 $\frac{4}{15}$; 1.3 $\frac{p^2 - 4 \ln 2}{8}$; 1.4 $2 - 2 \ln 2$; 1.5 $\ln 2$; 1.6 $-\frac{17}{18}$; 1.7 $\frac{1}{3}$; 1.8 $\frac{p}{8}$;

1.9 $1 + \ln \sqrt{\frac{3}{2}}$; 1.10 $1 + \ln \sqrt{\frac{3}{2}}$; 1.11 $\frac{p^2}{18} + \ln 2$; 1.12 $\ln 2 - \frac{p^5}{5 \cdot 3^5}$; 1.13 $\frac{1 - \ln 2}{2}$; 1.14 $\frac{4}{3}$;

1.15 $\frac{p}{6} - \ln 2$; 1.16 $\ln \sqrt{\frac{3}{2}}$; 1.17 $\frac{3}{2}$; 1.18 $\frac{p}{12}$; 1.19 $\frac{e^2 + 1}{2}$; 1.20 $\ln \sqrt{\frac{3 + 2\sqrt{3}}{3}}$;

1.21 $\frac{\ln 4 - 1}{4}$; 1.22 $-\frac{1}{2} \ln^2 \frac{\sqrt{2}}{2}$; 1.23 $\frac{\operatorname{tg}^2 1}{2}$; 1.24 $\frac{15p^4}{2^{10}} - \frac{p}{4}$; 1.25 $\frac{1}{2p}$; 1.26 $-\frac{3}{16}$;

1.27 $\frac{4\sqrt{2} - 2}{p}$; 1.28 $\frac{\ln 4 + p}{8}$; 1.29 $\ln \sqrt{\frac{5 + 2\sqrt{5}}{5}}$; 1.30 23,1.

Bo'laklab integrallash.

Agar $u(x)$ va $v(x)$ lar $[a, b]$ kesmada differensiallanuvchi funksiyalar bo'lsa, u holda ushbu formula o'rinalidir:

$$\int_a^b u dv = uv \Big|_a^b - \int_a^b v du.$$

2–masala. Aniq integralni hisoblang.

$$\begin{aligned}
 \int_{-2}^0 (x^2 - 4) \cos 3x dx &= \left| \begin{array}{ll} u = x^2 - 4 & dv = \cos 3x dx \\ du = 2x dx & v = \frac{1}{3} \sin 3x \end{array} \right| = \\
 &= \frac{1}{3} (x^2 - 4) \sin 3x \Big|_{-2}^0 - \frac{2}{3} \int_{-2}^0 x \sin 3x dx = \left| \begin{array}{ll} u = x & dv = \sin 3x dx \\ du = dx & v = -\frac{1}{3} \cos 3x \end{array} \right| = \\
 &= -\frac{2}{3} \left(-\frac{1}{3} x \cos 3x \right) \Big|_{-2}^0 + \frac{1}{3} \int_{-2}^0 \cos 3x dx = \\
 &= -\frac{2}{3} \left(-\frac{2}{3} \cos 6 + \frac{1}{9} \sin 3x \right) \Big|_{-2}^0 = \frac{4}{9} \cos 6 - \frac{2}{27} \sin 6.
 \end{aligned}$$

1. $\int_{-2}^0 (x^2 + 5x + 6) \cos 2x dx.$

2. $\int_{-1}^0 (x^2 + 4x + 3) \cos x dx.$

3. $\int_{-2}^0 (x + 2)^2 \cos 3x dx.$

4. $\int_{-4}^0 (x^2 + 7x + 12) \cos x dx.$

5. $\int_0^p (2x^2 + 4x + 7) \cos 2x dx.$

6. $\int_0^p (9x^2 + 9x + 11) \cos 3x dx.$

7. $\int_0^p (8x^2 + 16x + 17) \cos 4x dx.$

8. $\int_0^{2p} (3x^2 + 5) \cos 2x dx.$

9. $\int_0^{2p} (2x^2 - 15) \cos 3x dx.$

10. $\int_0^{2p} (3 - 7x^2) \cos 2x dx.$

11. $\int_0^{2p} (1 - 8x^2) \cos 4x dx.$

12. $\int_{-1}^0 (x^2 + 2x + 1) \sin 3x dx.$

13. $\int_0^3 (x^2 - 2x) \sin 2x dx.$

14. $\int_0^p (x^2 - 3x + 2) \sin x dx.$

$$15. \int_0^{p/2} (x^2 - 5x + 6) \sin 3x dx.$$

$$16. \int_{-3}^0 (x^2 + 6x + 9) \sin 2x dx.$$

$$17. \int_0^{p/4} (x^2 + 17,5) \sin 2x dx.$$

$$18. \int_0^{p/2} (1 - 5x^2) \sin x dx.$$

$$19. \int_{p/4}^3 (3x - x^2) \sin 2x dx.$$

$$20. \int_1^2 x \ln^2 x dx.$$

$$21. \int_1^{e^2} \frac{\ln^2 x dx}{\sqrt{x}}.$$

$$22. \int_1^8 \frac{\ln^2 x dx}{\sqrt[3]{x^2}}.$$

$$23. \int_0^1 (x+1) \ln^2 (x+1) dx.$$

$$24. \int_2^3 (x-1)^3 \ln^2 (x-1) dx.$$

$$25. \int_{-1}^0 (x+2)^3 \ln^2 (x+2) dx.$$

$$26. \int_0^2 (x+1)^2 \ln^2 (x+1) dx.$$

$$27. \int_1^e \sqrt{x} \ln^2 x dx.$$

$$28. \int_{-1}^1 x^2 e^{-x/2} dx.$$

$$29. \int_0^1 x^2 e^{3x} dx.$$

$$30. \int_{-2}^0 (x^2 + 2) e^{x/2} dx.$$

Javoblar. 2.1 $\frac{5 - \cos 4 - \sin 4}{4}$; 2.2 $4 - 2 \cos 1 - 2 \sin 1$; 2.3 $\frac{12 - 2 \sin 6}{27}$;

2.4 $7 + \cos 4 - 2 \sin 4$; 2.5 p ; 2.6 $-2p - 2$; 2.7 p ; 2.8 $2p$; 2.9 $\frac{8p}{9}$; 2.10 $-7p$; 2.11 $-2p$;

2.12 $-\frac{7 + 2 \cos 3}{27}$; 2.13 $\frac{3 \sin 6 + \cos 6 - 1}{4}$; 2.14 $p^2 - 3p$; 2.15 $\frac{67 - 3p}{27}$; 2.16 $-\frac{17 + \cos 6}{4}$;

2.17 $\frac{17}{2} + \frac{p}{8}$; 2.18 $11 - 5p$; 2.19 $\frac{p - 6 + 2 \cos 6 - 6 \sin 6}{8}$; 2.20 $2 \ln^2 2 - 2 \ln 2 + \frac{3}{4}$;

2.21 $8e - 16$; 2.22 $6 \ln^2 8 - 36 \ln 8 + 54$; 2.23 $2 \ln^2 2 - 2 \ln 2 + \frac{3}{4}$;

2.24 $4 \ln^2 2 - 2 \ln 2 + \frac{15}{32}$; 2.25 $4 \ln^2 2 - 2 \ln 2 + \frac{15}{32}$; 2.26 $9 \ln^2 3 - 6 \ln 3 + 1 \frac{25}{27}$;

$$2.27 \frac{10e\sqrt{e} - 16}{27}; 2.28 - \frac{26}{\sqrt{e}} + 10\sqrt{e}; 2.29 \frac{5e^3 - 2}{27}; 2.30 20 - \frac{44}{e}.$$

$R(\sin x, \cos x)dx$ ko'rinishdagi integrallar

$R(\sin x, \cos x)dx$ ko'rinishdagi integralar (R - $\sin x$ va $\cos x$ larga nisbatan ratsional funksiya) $\operatorname{tg} \frac{x}{2} = t$ almashtirish yordamida ratsional funksiylarning integrallariga keltiriladi.

$$\sin x = \frac{2t \operatorname{tg} \frac{x}{2}}{1 + t^2 \operatorname{tg}^2 \frac{x}{2}} = \frac{2t}{1 + t^2};$$

$$\cos x = \frac{1 - t^2 \operatorname{tg}^2 \frac{x}{2}}{1 + t^2 \operatorname{tg}^2 \frac{x}{2}} = \frac{1 - t^2}{1 + t^2};$$

$$x = 2 \arctg t; \quad dx = \frac{2}{1 + t^2} dt.$$

$R(\sin x, \cos x)dx = R\left(\frac{2t}{1+t^2}, \frac{1-t^2}{1+t^2}\right) \frac{2dt}{1+t^2}$ ko'rinishga keladi. Bunday almashtirish *universal almashtirish* deyiladi.

3-masala. Aniq integralni hisoblang.

$$\int_0^{p/2} \frac{\cos x - \sin x}{(1 + \sin x)^2} dx = \left| \begin{array}{l} \operatorname{tg} \frac{x}{2} = t \quad \cos x = \frac{1-t^2}{1+t^2} \\ dx = \frac{2}{1+t^2} dt \quad \sin x = \frac{2t}{1+t^2} \end{array} \right| = \int_0^1 \frac{1-t^2}{1+t^2} - \frac{2t}{1+t^2} \times \frac{2dt}{1+t^2} =$$

$$= \int_0^1 \frac{2(1-2t-t^2)}{(1+t)^4} dt.$$

$\frac{2(1-2t-t^2)}{(1+t)^4}$ to'g'ri kasrni sodda ratsional kasrlar yig'indisi ko'rinishida

yozamiz:

$$\frac{2-4t-2t^2}{(1+t)^4} = \frac{A}{1+t} + \frac{B}{(1+t)^2} + \frac{C}{(1+t)^3} + \frac{D}{(1+t)^4} =$$

$$= \frac{A(1+t)^3 + B(1+t)^2 + C(1+t) + D}{(1+t)^4}.$$

$$A(1+t)^3 + B(1+t)^2 + C(1+t) + D = 2 - 4t - 2t^2.$$

$$t = -1 \text{ da, } D = 4;$$

x ning bir xil darajalari oldidagi koeffitsiyentlarni tenglaymiz:

$$t^3: \quad A = 0;$$

$$t^2: \quad 3A + B = -2 \Rightarrow B = -2;$$

$$t: \quad 3A + 2B + C = -4 \Rightarrow C = 0;$$

Demak,

$$\int_0^1 \frac{4}{(1+t)^4} - \frac{2}{(1+t)^2} dt = \int_0^1 \frac{4}{3(1+t)^3} + \frac{2}{1+t} dt = -\frac{4}{3 \times 8} + 1 + \frac{4}{3} - 2 = \frac{1}{6}.$$

$$1. \int_0^{2 \arctg 2} \frac{dx}{\sin^2 x (1 - \cos x)}.$$

$$2. \int_0^{p/2} \frac{\cos x dx}{2 + \cos x}.$$

$$3. \int_0^{2\operatorname{arctg} 2} \frac{dx}{\sin^2 x (1 + \cos x)^{p/2}}.$$

$$5. \int_0^{2\operatorname{arctg} 3} \frac{dx}{2\operatorname{arctg} 2 \cos x (1 - \cos x)}.$$

$$7. \int_0^{p/2} \frac{dx}{2\operatorname{arctg} \frac{1}{2} (1 + \sin x - \cos x)^2}.$$

$$9. \int_0^{2p/3} \frac{1 + \sin x}{1 + \cos x + \sin x} dx.$$

$$11. \int_0^{p/2} \frac{(1 + \cos x) dx}{1 + \cos x + \sin x}.$$

$$13. \int_0^{2\operatorname{arctg} \frac{1}{2}} \frac{1 + \sin x}{(1 - \sin x)^2} dx.$$

$$15. \int_0^{2\operatorname{arctg} \frac{1}{3}} \frac{\cos x dx}{(1 + \cos x)(1 - \sin x)}.$$

$$17. \int_0^{p/2} \frac{\cos x dx}{(1 + \cos x - \sin x)^2}.$$

$$19. \int_0^{2\operatorname{arctg} \frac{1}{2}} \frac{(1 - \sin x) dx}{\cos x (1 + \cos x)}.$$

$$21. \int_0^{p/2} \frac{\sin x dx}{(1 + \sin x + \cos x)^2}.$$

$$23. \int_0^{p/2} \frac{\cos^2 x dx}{(1 + \cos x - \sin x)^2}.$$

$$25. \int_0^{2p/3} \frac{\cos^2 x dx}{(1 + \cos x + \sin x)^2}.$$

$$4. \int_0^{p/2} \frac{\cos x dx}{2\operatorname{arctg} \frac{1}{2} (1 - \cos x)^3}.$$

$$6. \int_0^{2\operatorname{arctg} \frac{1}{2}} \frac{dx}{2\operatorname{arctg} \frac{1}{3} \sin x (1 - \sin x)}.$$

$$8. \int_0^{p/2} \frac{\cos x dx}{5 + 4 \cos x}.$$

$$10. \int_0^{p/2} \frac{\cos x dx}{p/3 (1 + \sin x - \cos x)}.$$

$$12. \int_0^{p/2} \frac{\sin x dx}{1 + \cos x + \sin x}.$$

$$14. \int_0^{\frac{p}{2}} \frac{\cos x}{1 + \cos x + \sin x} dx.$$

$$16. \int_0^{p/2} \frac{\cos x dx}{(1 + \cos x - \sin x)^2}.$$

$$18. \int_0^{p/2} \frac{\cos x dx}{(1 + \cos x + \sin x)^2}.$$

$$20. \int_0^{p/2} \frac{\sin x dx}{(1 + \sin x)^2}.$$

$$22. \int_0^{p/2} \frac{\sin x dx}{(1 + \cos x - \sin x)^2}.$$

$$24. \int_0^{p/2} \frac{\sin^2 x dx}{(1 + \cos x + \sin x)^2}.$$

$$26. \int_0^{2\operatorname{arctg} 2} \frac{dx}{p/2 \sin x (1 + \sin x)}.$$

$$27. \int_0^{p/2} \frac{dx}{(1 + \sin x + \cos x)^2}.$$

$$28. \int_0^{p/2} \frac{\sin x dx}{2 + \sin x}.$$

$$29. \int_0^{p/4} \frac{dx}{\cos x(1 + \cos x)}.$$

$$30. \int_0^{p/2} \frac{\sin x dx}{5 + 3 \sin x}.$$

Javoblar. 3.1 $\frac{55}{96}$; 3.2 $\frac{(9 - 4\sqrt{3})p}{18}$; 3.3 $1\frac{5}{24}$; 3.4 -4 ; 3.5 $\frac{1}{6} + \ln 2 - \ln 3$; 3.6 $\ln 3 - \ln 2 + 1$;

3.7 $\frac{2}{3} - \ln \frac{3}{2}$; 3.8 $\frac{p}{8} - \frac{5}{6} \arctg \frac{1}{3}$; 3.9 $\frac{p}{3} + \ln 2$; 3.10 $\frac{1}{2} \ln 2 - \frac{p}{12}$; 3.11 $\frac{1}{2} \ln 2 + \frac{p}{4}$;

3.12 $-\frac{1}{2} \ln 2 + \frac{p}{4}$; 3.13 $\frac{26}{3}$; 3.14 $\frac{p}{4} - \frac{1}{2} \ln 2$; 3.15 $-\frac{1}{3} - 2 \ln \frac{2}{3}$; 3.16 $\frac{p}{3} - \ln 2$;

3.17 $-\frac{1}{2} + \ln 2$; 3.18 $-\frac{1}{2} + \ln 2$; 3.19 $-\frac{1}{2} + 2 \ln \frac{3}{2}$; 3.20 $\frac{1}{3}$; 3.21 $\ln 2 - \frac{1}{2}$; 3.22 $\frac{1}{2} - \ln 2$;

3.23 $\frac{\sqrt{3}}{2} - \ln 2$; 3.24 $\frac{1}{2} - \frac{1}{2} \ln 2$; 3.25 $\frac{\sqrt{3}}{2} - \ln 2$; 3.26 $\ln 2 - \frac{1}{3}$;

3.27 $1 - \ln 2$; 3.28 $\frac{p}{2} - \frac{2p}{3\sqrt{3}}$; 3.29 $\frac{\sqrt{2} - 2}{\sqrt{2}} - \ln(\sqrt{2} - 1)$; 3.30 $\frac{p - 5 \arctg 2 + \arctg \frac{3}{4}}{6}$.

$\int \operatorname{tg}^m x dx$ va $\int \operatorname{ctg}^m x dx$ (bu yerda m - butun musbat son) ko'rinishdagi integrallarda mos ravishda

$$\operatorname{tg} t = t, \quad dx = \frac{dt}{1+t^2}$$

$$\operatorname{ctg} t = t, \quad dx = -\frac{dt}{1+t^2}$$

$$\sin 2x = \frac{2 \operatorname{tg} x}{1 + \operatorname{tg}^2 x} = \frac{2t}{1+t^2}$$

o'rniga qo'yish orqali hisoblanadi.

4-masala. Aniq integralni hisoblang.

$$\int_{\rho/4}^{\arctg 3} \frac{dx}{(3\operatorname{tg} x + 5) \sin 2x} = \left| \begin{array}{l} \operatorname{tg} x = t \\ dx = \frac{dt}{1+t^2} \end{array} \quad \sin 2x = \frac{2t}{1+t^2} \right| =$$

$$= \int_1^3 \frac{dt}{(3t+5) \frac{2t}{1+t^2}} = \frac{1}{2} \int_1^3 \frac{dt}{t(3t+5)}.$$

$$\frac{1}{t(3t+5)} = \frac{A}{t} + \frac{B}{3t+5} = \frac{A(3t+5) + Bt}{t(3t+5)},$$

$$A(3t+5) + Bt = 1.$$

$$t=0 \text{ da, } A = \frac{1}{5};$$

$$t = -\frac{5}{3} \text{ da, } B = -\frac{3}{5};$$

Shunday qilib,

$$\frac{1}{10} \int_1^3 \frac{dt}{t(3t+5)} = \frac{1}{10} \left(\ln|t| - \ln|3t+5| \right) \Big|_1^3 = \frac{1}{10} (\ln 3 - \ln 14 - 0 + \ln 8) =$$

$$= \frac{1}{10} \ln \frac{24}{14} = \frac{1}{10} \ln \frac{12}{7}.$$

1. $\int_{\arccos(4/\sqrt{17})}^{\rho/4} \frac{2\operatorname{ctg} x + 1}{(2\sin x + \cos x)^2} dx.$
2. $\int_0^{\arccos(4/\sqrt{17})} \frac{3 + 2\operatorname{tg} x}{2\sin^2 x + 3\cos^2 x - 1} dx.$
3. $\int_{\rho/4}^{\arctg 3} \frac{4\operatorname{tg} x - 5}{1 - \sin 2x + 4\cos^2 x} dx.$
4. $\int_0^{\arctg \frac{1}{3}} \frac{(8 + \operatorname{tg} x)}{18\sin^2 x + 2\cos^2 x} dx.$
5. $\int_0^{\arccos \sqrt{2/3}} \frac{\operatorname{tg} x + 2}{\sin^2 x + 2\cos^2 x - 3} dx.$
6. $\int_{\arcsin(1/\sqrt{37})}^{\rho/4} \frac{6\operatorname{tg} x dx}{3\sin 2x + 5\cos^2 x}.$
7. $\int_0^{\rho/4} \frac{2\operatorname{tg}^2 x - 11\operatorname{tg} x - 22}{4 - \operatorname{tg} x} dx.$
8. $\int_{-\arctg(1/3)}^0 \frac{3\operatorname{tg} x + 1}{2\sin 2x - 5\cos 2x + 1} dx.$

$$9. \int_{\rho/4}^{\arctg 3} \frac{1 + \operatorname{ctg} x}{(\sin x + 2 \cos x)^2} dx.$$

$$10. \int_{\rho/4}^{\arccos(1/\sqrt{3})} \frac{\operatorname{tg} x}{\sin^2 x - 5 \cos^2 x + 4} dx.$$

$$11. \int_0^{\rho/4} \frac{6 \sin^2 x}{3 \cos 2x - 4} dx.$$

$$12. \int_0^{\arctg 3} \frac{4 + \operatorname{tg} x}{2 \sin^2 x + 18 \cos^2 x} dx.$$

$$13. \int_0^{\arctg 2} \frac{12 + \operatorname{tg} x}{3 \sin^2 x + 12 \cos^2 x} dx.$$

$$14. \int_0^{\arctg(2/3)} \frac{6 + \operatorname{tg} x}{9 \sin^2 x + 4 \cos^2 x} dx.$$

$$15. \int_0^{\arcsin \sqrt{3/7}} \frac{\operatorname{tg}^2 x dx}{3 \sin^2 x + 4 \cos^2 x - 7}.$$

$$16. \int_0^{\rho/4} \frac{7 + 3 \operatorname{tg} x}{(\sin x + 2 \cos x)^2} dx.$$

$$17. \int_{\arcsin(2/\sqrt{5})}^{\arcsin(3/\sqrt{10})} \frac{2 \operatorname{tg} x + 5}{(5 - \operatorname{tg} x) \sin 2x} dx.$$

$$18. \int_{-\arccos(1/\sqrt{10})}^0 \frac{3 \operatorname{tg}^2 x - 50}{2 \operatorname{tg} x + 7} dx.$$

$$19. \int_0^{\rho/4} \frac{5 \operatorname{tg} x + 2}{2 \sin 2x + 5} dx.$$

$$20. \int_{\rho/4}^{\arcsin(2/\sqrt{5})} \frac{4 \operatorname{tg} x - 5}{4 \cos^2 x - \sin 2x + 1} dx.$$

$$21. \int_0^{\arcsin \sqrt{7/8}} \frac{6 \sin^2 x dx}{4 + 3 \cos 2x}.$$

$$22. \int_{-\arccos(1/\sqrt{5})}^0 \frac{11 - 3 \operatorname{tg} x}{\operatorname{tg} x + 3} dx.$$

$$23. \int_0^{\arcsin(3/\sqrt{10})} \frac{2 \operatorname{tg} x - 5}{(4 \cos x - \sin x)^2} dx.$$

$$24. \int_{\rho/4}^{\arccos(1/\sqrt{26})} \frac{36 dx}{(6 - \operatorname{tg} x) \sin 2x}.$$

$$25. \int_0^{\rho/4} \frac{4 - 7 \operatorname{tg} x}{2 + 3 \operatorname{tg} x} dx.$$

$$26. \int_{-\arcsin(2/\sqrt{5})}^{\rho/4} \frac{2 - \operatorname{tg} x}{(\sin x + 3 \cos x)^2} dx.$$

$$27. \int_{\rho/4}^{\arcsin \sqrt{2/3}} \frac{8 \operatorname{tg} x dx}{3 \cos^2 x + 8 \sin 2x - 7}.$$

$$28. \int_{\arccos(1/\sqrt{10})}^{\arccos(1/\sqrt{26})} \frac{12 dx}{(6 + 5 \operatorname{tg} x) \sin 2x}.$$

$$29. \int_0^{\rho/3} \frac{\operatorname{tg}^2 x}{4 + 3 \cos 2x} dx.$$

$$30. \int_0^{\arccos(1/\sqrt{6})} \frac{3 \operatorname{tg}^2 x - 1}{\operatorname{tg}^2 x + 5} dx.$$

Javoblar. 4.1 $2 \ln 2 - \frac{1}{2}$; 4.2 $\frac{3}{\sqrt{2}} \arctg \frac{1}{4\sqrt{2}} + \ln \frac{33}{32}$; 4.3 $2 \ln 2 - \frac{\rho}{8}$; 4.4 $\frac{\rho}{3} + \frac{\ln 2}{36}$;

4.5 $-\frac{\ln 2 + \sqrt{2}\rho}{4}$; 4.6 $\frac{5}{6} \ln \frac{6}{11}$; 4.7 $2 \ln 3 - 6 \ln 2 - 5 \frac{\rho}{4}$; 4.8 $\frac{1}{4} \ln \frac{6}{7}$; 4.9 $\frac{1}{4} \ln \frac{9}{5} + \frac{1}{15}$;

$$\begin{aligned}
& 4.10 \frac{1}{10} \ln \frac{9}{4}; 4.11 - \frac{\rho}{4} + \frac{1}{\sqrt{7}} \operatorname{arctg} \sqrt{7}; 4.12 \frac{\rho}{6} + \frac{\ln 2}{4}; 4.13 \frac{\rho}{2} + \frac{\ln 2}{6}; 4.14 \frac{\rho}{4} + \frac{\ln 2}{18}; \\
& 4.15 - \frac{\sqrt{3}}{8} + \frac{\sqrt{3}\rho}{32}; 4.16 3 \ln \frac{3}{2} + \frac{1}{6}; 4.17 2 \ln \frac{3}{2}; 4.18 - \frac{1}{2} \ln 7 - \ln 10 - 7 \operatorname{arctg} 3; \\
& 4.19 \frac{1}{2} \ln \frac{14}{5}; 4.20 2 \ln \frac{5}{4} - \frac{1}{2} \operatorname{arctg} \frac{1}{2}; 4.21 \frac{\sqrt{7}\rho}{4} - \operatorname{arctg} \sqrt{7}; 4.22 \ln 45 + 3 \operatorname{arctg} 2; \\
& 4.23 \frac{9}{4} - \ln 16; 4.24 \frac{\ln 5}{6}; 4.25 2 \ln 5 - 3 \ln 2 - \frac{\rho}{4}; 4.26 \frac{15}{4} - \ln 4; \\
& 4.27 \frac{4}{21} \times \ln \left| \frac{7\sqrt{2} - 2}{5} \right| - \frac{4}{3} \times \ln |2 - \sqrt{2}|; 4.28 \ln \frac{105}{93}; 4.29 \sqrt{3} - \sqrt{7} \times \operatorname{arctg} \sqrt{\frac{3}{7}}; \\
& 4.30 \frac{\rho}{\sqrt{5}} - \operatorname{arctg} \sqrt{5}.
\end{aligned}$$

Ösin^m x × cosⁿ x dx ko'rinishdagi integrallar,

bu yerda m va n – butun sonlar.

1. Agar m va n sonlarning hech bo'lmaganda bittasi toq musbat son, masalan, $m = 2k + 1$ bo'lsa bo'lsa, u holda quyidagicha yo'l tutamiz:

$$\begin{aligned}
& \text{Ösin}^{2k+1} x \times \cos^n x dx = \text{Ösin}^{2k} x \times \cos^n x \times \sin x dx = \\
& = - \text{Ö}(1 - \cos^2 x)^k \times \cos^n x d(\cos x).
\end{aligned}$$

Masalan,

$$\begin{aligned}
& \text{Ösin}^5 x \times \cos^2 x dx = \text{Ösin}^4 x \times \cos^2 x \times \sin x dx = \\
& = - \text{Ö}(1 - \cos^2 x)^2 \times \cos^2 x d(\cos x) = \\
& = - \frac{1}{3} \cos^3 x + \frac{2}{5} \cos^5 x + \frac{1}{7} \cos^7 x + C.
\end{aligned}$$

Agar m va n sonlardan biri toq musbat son, boshqasi istalgan haqiqiy son bo'lsa ham xuddi yuqoridek yo'l tutamiz.

2. Agar m va n juft musbat sonlar bo'lsa, integralni

$$\sin x \times \cos x = \frac{1}{2} \sin 2x, \quad \sin^2 x = \frac{1 - \cos 2x}{2}, \quad \cos^2 x = \frac{1 + \cos 2x}{2}$$

Trigonometrik formulalar yordamida hisoblaymiz.

Masalan,

$$\begin{aligned} \int \sin^2 x \cos^4 x dx &= \int (\sin x \times \cos x)^2 \cos^2 x dx = \\ &= \int \frac{\sin^2 2x}{4} \times \frac{1 + \cos 2x}{2} dx = \frac{1}{8} \int (\sin^2 2x + \sin^2 2x \times \cos 2x) dx = \\ &= \frac{1}{8} \int \frac{1 - \cos 4x}{2} dx + \frac{1}{8 \times 2} \int \sin^2 2x d(\sin 2x) = \\ &= \frac{1}{16} x - \frac{\sin 4x}{4} + \frac{1}{48} \sin^3 2x + C = \\ &= \frac{1}{16} x - \frac{\sin 4x}{4} + \frac{\sin^3 2x}{3} + C. \end{aligned}$$

3. Agar m va n juft-toqligi bir xil bo'lgan butun manfiy sonlar bo'lsa, integral

$$1 + \operatorname{tg}^2 x = \frac{1}{\cos^2 x}, \quad 1 - \operatorname{ctg}^2 x = \frac{1}{\sin^2 x} \text{ yoki } \frac{1}{\sin^2 x} = \frac{1 + \operatorname{tg}^2 x}{\operatorname{tg}^2 x}$$

formulalar yordamida hisoblanadi.

$$\begin{aligned} \int \frac{dx}{\cos^4 x} &= \int \frac{1}{\cos^2 x} \times \frac{dx}{\cos^2 x} = \int (1 + \operatorname{tg}^2 x)^2 d(\operatorname{tg} x) = \\ &= \operatorname{tg} x + \frac{\operatorname{tg}^3 x}{3} + C. \end{aligned}$$

5–masala. Aniq integralni hisoblang.

$$\begin{aligned}
& \int_0^p 2^4 \cos^8 x \, dx = \int_0^p \cos^2 x \, dx = \frac{1}{2} \int_0^p (1 + \cos x) \, dx = \\
& = \int_0^p (1 + \cos x)^4 \, dx = \int_0^p (1 + 2 \cos x + \cos^2 x)^2 \, dx = \\
& = \int_0^p (1 + 3 \cos x + 6 \cos^2 x + 4 \cos^3 x + \cos^4 x) \, dx = \\
& = \int_0^p \left(\frac{35}{8} + 3 \cos x + \frac{7}{2} \cos 2x + \frac{1}{8} \cos 4x \right) \, dx + 4 \int_0^p (1 - \sin^2 x) \cos x \, dx = \\
& = \frac{35}{8} x + 3 \sin x + \frac{7}{4} \sin 2x + \frac{1}{32} \sin 4x \Big|_0^p + 4 \int_0^p (1 - \sin^2 x) d(\sin x) = \\
& = \frac{35}{8} p + 4 \left(\sin x - \frac{1}{3} \sin^3 x \right) \Big|_0^p = \frac{35}{8} p.
\end{aligned}$$

$$1. \int_0^p 2^8 \sin^8 x \, dx.$$

$$2. \int_0^p 2^4 \sin^6 x \cos^2 x \, dx.$$

$$3. \int_0^{2p} \sin^4 x \cos^4 x \, dx.$$

$$4. \int_0^{2p} \sin^2 x \cos^6 x \, dx.$$

$$5. \int_{-p/2}^0 2^8 \sin^8 x \, dx.$$

$$6. \int_{p/2}^p 2^4 \sin^6 x \cos^2 x \, dx.$$

$$7. \int_0^p 2^4 \sin^4 x \cos^4 x \, dx.$$

$$8. \int_0^{2p} \sin^2 x \cos^6 x \, dx.$$

$$9. \int_0^{2p} \cos^8 x \, dx.$$

$$10. \int_0^p 2^4 \sin^8 x \, dx.$$

$$11. \int_{-p}^0 2^8 \sin^6 x \cos^2 x \, dx.$$

$$12. \int_{p/2}^{2p} 2^8 \sin^4 x \cos^4 x \, dx.$$

$$13. \int_0^p 2^4 \sin^2 x \cos^6 x \, dx.$$

$$14. \int_0^{2p} \cos^8 x \, dx.$$

$$15. \int_0^{2p} \sin^8 \frac{x}{4} dx.$$

$$17. \int_{-p/2}^0 \sin^4 x \cos^4 x dx.$$

$$19. \int_0^p \cos^8 x dx.$$

$$21. \int_0^{2p} \sin^6 \frac{x}{4} \cos^2 \frac{x}{4} dx.$$

$$23. \int_{-p/2}^0 \sin^2 x \cos^6 x dx.$$

$$25. \int_0^p \sin^8 x dx.$$

$$27. \int_0^{2p} \sin^4 \frac{x}{4} \cos^4 \frac{x}{4} dx.$$

$$29. \int_{-p/2}^0 \cos^8 x dx.$$

$$16. \int_0^p \sin^6 \frac{x}{2} \cos^2 \frac{x}{2} dx.$$

$$18. \int_{p/2}^p \sin^2 x \cos^6 x dx.$$

$$20. \int_0^{2p} \sin^8 x dx.$$

$$22. \int_0^p \sin^4 \frac{x}{2} \cos^4 \frac{x}{2} dx.$$

$$24. \int_{p/2}^p \cos^8 x dx.$$

$$26. \int_0^{2p} \sin^6 x \cos^2 x dx.$$

$$28. \int_0^p \sin^2 \frac{x}{2} \cos^6 \frac{x}{2} dx.$$

$$30. \int_0^{2p} \sin^4 3x \cos^4 3x dx.$$

Javoblar. 5.1 $35p$; 5.2 $\frac{5p}{8}$; 5.3 $\frac{3p}{64}$; 5.4 $\frac{5p}{64}$; 5.5 $35p$; 5.6 $\frac{5p}{16}$; 5.7 $\frac{3p}{4}$; 5.8 $\frac{5p}{2^6}$;

5.9 $\frac{35p}{64}$; 5.10 $\frac{35p}{8}$; 5.11 $10p$; 5.12 $9p$; 5.13 $\frac{5p}{8}$; 5.14 $\frac{35p}{64}$; 5.15 $\frac{35p}{64}$;

5.16 $\frac{5p}{8}$; 5.17 $3p$; 5.18 $5p$; 5.19 $\frac{35p}{8}$; 5.20 $\frac{35p}{64}$; 5.21 $\frac{5p}{64}$; 5.22 $\frac{3p}{8}$;

5.23 $5p$; 5.24 $105p$; 5.25 $\frac{35p}{8}$; 5.26 $\frac{5p}{64}$; 5.27 $\frac{3p}{64}$; 5.28 $\frac{5p}{8}$; 5.29 $35p$; 5.30 $\frac{3p}{64}$.

$\int_0^R \frac{ax+b}{cx+d} \frac{dx}{x^2+q_1^2}, \frac{ax+b}{cx+d} \frac{dx}{x^2+q_2^2}, \dots, \frac{dx}{x^2+q_n^2}$ ko'rinishdagi integrallar, bu

yerda $p_1, q_1, p_2, q_2, \dots$ - butun sonlar. Agar barcha q_1, q_2, \dots maxrajlarining

eng kichik karralisi k bo'lsa, u holda ushbu integral $\frac{ax+b}{cx+d} = t^k$ o'rniga qo'yish yordamida ratsional funksiya dan olingan integrallarga keltiriladi.

6-masala. Aniq integralni hisoblang.

$$\begin{aligned}
 \int_0^9 \sqrt{\frac{9-2x}{2x-21}} dx &= \int_{dx=\frac{12t}{(t^2+1)} dt}^{\frac{9-2x}{2x-21}=t^2} = 12 \int \frac{t}{(t^2+1)^2} dt = \\
 &= 12 \int \frac{t^2}{(t^2+1)^2} = \int_{dt=\frac{da}{\cos^2 a}}^{t=\operatorname{tg} a} = 12 \int g^2 \cos^2 a da = \\
 &= 12 \int \sin^2 a da = 6 \int (1 - \cos 2a) da = 6 \operatorname{arctg} T - 3 \sin(2 \operatorname{arctg} t) = \\
 &= 6 \operatorname{arctg} \sqrt{\frac{9-2x}{2x-21}} - 3 \sin(2 \operatorname{arctg} \sqrt{\frac{9-2x}{2x-21}}) = \\
 &= 6 \operatorname{arctg} \sqrt{3} - 3 \sin(2 \operatorname{arctg} \sqrt{3}) - 6 \operatorname{arctg} \frac{1}{3} + 3 \sin(2 \operatorname{arctg} \frac{1}{3}) = \\
 &= 2p - 3 \sin \frac{2p}{3} - p + 3 \sin \frac{p}{3} = p - 3 \frac{\sqrt{3}}{2} + 3 \frac{\sqrt{3}}{2} = p.
 \end{aligned}$$

1. $\int_0^1 \frac{4\sqrt{1-x} - \sqrt{3x+1}}{(\sqrt{3x+1} + 4\sqrt{1-x})(3x+1)^2} dx.$
2. $\int_1^{64} \frac{1 - \sqrt[6]{x} + 2\sqrt[3]{x}}{x + 2\sqrt{x^3} + \sqrt[3]{x^4}} dx.$
3. $\int_{-14/15}^{-7/8} \frac{6\sqrt{x+2}}{(x+2)^2 \sqrt{x+1}} dx.$
4. $\int_0^5 \sqrt{\frac{5-x}{5+x}} \frac{dx}{(5+x)\sqrt{25-x^2}}.$
5. $\int_8^{12} \sqrt{\frac{6-x}{x-14}} dx.$
6. $\int_0^1 \sqrt{\frac{1-x}{1+x}} \frac{dx}{(1+x)\sqrt{1-x^2}}.$

$$7. \int_{5/2}^{10/3} \frac{\sqrt{x+2} + \sqrt{x-2}}{(\sqrt{x+2} - \sqrt{x-2})(x-2)^2} dx.$$

$$8. \int_1^8 \frac{5\sqrt{x+24}}{(x+24)^2 \sqrt{x}} dx.$$

$$9. \int_1^2 \frac{x + \sqrt{3x-2} - 10}{\sqrt{3x-2} + 7} dx.$$

$$10. \int_6^{10} \sqrt{\frac{4-x}{x-12}} dx.$$

$$11. \int_0^2 \frac{(4\sqrt{2-x} - \sqrt{2x+2})}{(\sqrt{2x+2} + 4\sqrt{2-x})(2x+2)^2} dx. \quad 12. \int_{-1/2}^0 \frac{xdx}{2 + \sqrt{2x+1}}.$$

$$13. \int_0^4 e^{\sqrt{\frac{4-x}{4+x}}} \frac{dx}{(4+x)\sqrt{16-x^2}}.$$

$$14. \int_{1/8}^1 \frac{15\sqrt{x+3}}{(x+3)^2 \sqrt{x}} dx.$$

$$15. \int_{-5/3}^1 \frac{\sqrt[3]{3x+5} + 2}{1 + \sqrt[3]{3x+5}} dx.$$

$$16. \int_2^3 \sqrt{\frac{3-2x}{2x-7}} dx.$$

$$17. \int_0^7 \frac{\sqrt{x+25} dx}{(x+25)^2 \sqrt{x+1}}.$$

$$18. \int_0^2 \frac{(4\sqrt{2-x} - \sqrt{3x+2}) dx}{(\sqrt{3x+2} + 4\sqrt{2-x})(3x+2)^2}.$$

$$19. \int_0^2 e^{\sqrt{\frac{2-x}{2+x}}} \frac{dx}{(2+x)\sqrt{4-x^2}}.$$

$$20. \int_3^5 \sqrt{\frac{2-x}{x-6}} dx.$$

$$21. \int_{1/24}^{1/3} \frac{5\sqrt{x+1}}{(x+1)^2 \sqrt{x}} dx.$$

$$22. \int_9^{15} \sqrt{\frac{6-x}{x-18}} dx.$$

$$23. \int_0^1 \frac{(4\sqrt{1-x} - \sqrt{2x+1}) dx}{(\sqrt{2x+1} + 4\sqrt{1-x})(2x+1)^2} dx. \quad 24. \int_1^{64} \frac{(2 + \sqrt[3]{x}) dx}{(\sqrt[6]{x} + 2\sqrt[3]{x} + \sqrt{x})\sqrt{x}}.$$

$$25. \int_{16/15}^{4/3} \frac{4\sqrt{x}}{x^2 \sqrt{x-1}} dx.$$

$$26. \int_0^6 \frac{e^{\sqrt{(6-x)/(6+x)}} dx}{(6+x)\sqrt{36-x^2}}.$$

$$27. \int_1^{64} \frac{6 - \sqrt{x} + \sqrt[4]{x}}{\sqrt{x^3 - 7x} - 6\sqrt[4]{x^3}} dx.$$

$$28. \int_0^1 \frac{(4\sqrt{1-x} - \sqrt{x+1}) dx}{(\sqrt{x+1} + 4\sqrt{1-x})(x+1)^2}.$$

$$29. \int_0^3 \frac{e^{\sqrt{(3-x)/(3+x)}} dx}{(3+x)\sqrt{9-x^2}}.$$

$$30. \int_0^2 \frac{(4\sqrt{2-x} - \sqrt{x+2}) dx}{(\sqrt{x+2} + 4\sqrt{2-x})(x+2)^2}.$$

Javoblar. 6.1 $\frac{1}{16} \ln 5$; 6.2 $6 \ln \frac{4}{3}$; 6.3 1; 6.4 $\frac{e-1}{5}$; 6.5 $\frac{4p}{3}$; 6.6 $e-1$; 6.7 $\frac{1}{2} + \ln 2$;
 6.8 $\frac{1}{8}$; 6.9 $-\frac{22}{27}$; 6.10 $\frac{4p}{3}$; 6.11 $\frac{1}{24} \ln 5$; 6.12 $\frac{7}{6} - 3 \ln \frac{3}{2}$; 6.13 $\frac{1}{4} \times (e-1)$; 6.14 3;
 6.15 $\frac{8}{3} + \ln 3$; 6.16 $\frac{p}{3}$; 6.17 $\frac{1}{40}$; 6.18 $\frac{1}{32} \ln 5$; 6.19 $\frac{e-1}{2}$; 6.20 $\frac{2p}{3}$; 6.21 3;
 6.22 $2p$; 6.23 $\frac{1}{12} \ln 5$; 6.24 $30 \ln \frac{3}{2} - 6$; 6.25 2; 6.26 $\frac{e-1}{6}$; 6.27 $4 \ln \frac{2}{2\sqrt{2}+1}$; 6.28 $\frac{1}{8} \ln 5$;
 6.29 $\frac{e-1}{3}$; 6.30 $\frac{1}{4} \ln 5$; 6.1 $\frac{1}{16} \ln 5$; 6.2 $6 \ln \frac{4}{3}$; 6.3 1; 6.4 $\frac{e-1}{5}$; 6.5 $\frac{4p}{3}$; 6.6 $e-1$; 6.7 $\frac{1}{2} + \ln 2$;
 6.8 $\frac{1}{8}$; 6.9 $-\frac{22}{27}$; 6.10 $\frac{4p}{3}$; 6.11 $\frac{1}{24} \ln 5$; 6.12 $\frac{7}{6} - 3 \ln \frac{3}{2}$; 6.13 $\frac{1}{4} \times (e-1)$; 6.14 3;
 6.15 $\frac{8}{3} + \ln 3$; 6.16 $\frac{p}{3}$; 6.17 $\frac{1}{40}$; 6.18 $\frac{1}{32} \ln 5$; 6.19 $\frac{e-1}{2}$; 6.20 $\frac{2p}{3}$; 6.21 3;
 6.22 $2p$; 6.23 $\frac{1}{12} \ln 5$; 6.24 $30 \ln \frac{3}{2} - 6$; 6.25 2; 6.26 $\frac{e-1}{6}$; 6.27 $4 \ln \frac{2}{2\sqrt{2}+1}$; 6.28 $\frac{1}{8} \ln 5$;
 6.29 $\frac{e-1}{3}$; 6.30 $\frac{1}{4} \ln 5$.

$R(x, \sqrt{a^2 \pm x^2})$ va $R(x, \sqrt{x^2 - a^2})$ ko'rinishdagi integrallar.

1. $\int_a^b R(x, \sqrt{a^2 - x^2}) dx$;

2. $\int_a^b R(x, \sqrt{a^2 + x^2}) dx$;

3. $\int_a^b R(x, \sqrt{x^2 - a^2}) dx$;

bu yerda R - ratsional funksiya.

Agar a) $x = a \sin t$ yoki $x = a \cos t$

b) $x = a \tan t$ yoki $x = a \cot t$

c) $x = a \sec t$ yoki $x = a \csc t$

trigonometrik o'rniga qo'yishlardan foydanilsa, bu integrallar

$\int R(\sin t, \cos t) dt$ ko'rinishdagi integrallarga keltiriladi.

7-masala. Aniq integralni hisoblang.

$$\begin{aligned} \int_0^3 \frac{dx}{(9+x^2)^{3/2}} &= \left| \begin{array}{l} x = 3 \operatorname{tg} t \\ dx = \frac{3dt}{\cos^2 t} \end{array} \right| = \int_0^{p/4} \frac{3dt}{(9+9\operatorname{tg}^2 t)^{3/2} \cos^2 t} = \\ &= \frac{3}{27} \int_0^{p/4} \frac{\cos^3 t}{\cos^2 t} dt = \frac{3}{27} \int_0^{p/4} \cos t dt = \frac{3}{27} \sin t \Big|_0^{p/4} = \frac{\sqrt{2}}{18}. \end{aligned}$$

1. $\int_0^{16} \sqrt{256 - x^2} dx.$

2. $\int_0^1 x^2 \sqrt{1 - x^2} dx.$

3. $\int_0^5 \frac{dx}{(25+x^2)\sqrt{25+x^2}}.$

4. $\int_0^{\sqrt{5}/2} \frac{dx}{\sqrt{(5-x^2)^3}}.$

5. $\int_1^2 \frac{\sqrt{x^2 - 1}}{x^4} dx.$

6. $\int_0^{\sqrt{2}/2} \frac{x^4 dx}{\sqrt{(1-x^2)^3}}.$

7. $\int_0^{\sqrt{3}} \frac{dx}{\sqrt{(4-x^2)^3}}.$

8. $\int_0^1 \frac{x^4 dx}{(2-x^2)^{3/2}}.$

9. $\int_0^2 \frac{x^2 dx}{\sqrt{16-x^2}}.$

10. $\int_0^2 \sqrt{4-x^2} dx.$

11. $\int_0^4 \frac{dx}{(16+x^2)^{3/2}}.$

12. $\int_0^4 x^2 \sqrt{16-x^2} dx.$

13. $\int_0^{5/2} \frac{x^2 dx}{\sqrt{25-x^2}} dx.$

14. $\int_0^5 x^2 \sqrt{25-x^2} dx.$

$$15. \int_0^4 \sqrt{16 - x^2} dx.$$

$$16. \int_0^{4\sqrt{3}} \frac{dx}{\sqrt{(64 - x^2)^3}}.$$

$$17. \int_{\sqrt{2}}^{2\sqrt{2}} \frac{\sqrt{x^2 - 2}}{x^4} dx.$$

$$18. \int_0^{2\sqrt{2}} \frac{x^4 dx}{(16 - x^2)\sqrt{16 - x^2}}.$$

$$19. \int_{-3}^3 x^2 \sqrt{9 - x^2} dx.$$

$$20. \int_1^{\sqrt{3}} \frac{dx}{\sqrt{(1 + x^2)^3}}.$$

$$21. \int_0^2 \frac{dx}{\sqrt{(16 - x^2)^3}}.$$

$$22. \int_0^2 \frac{x^4 dx}{\sqrt{(8 - x^2)^3}}.$$

$$23. \int_3^6 \frac{x^2 - 9}{x^4} dx.$$

$$24. \int_0^1 \sqrt{4 - x^2} dx.$$

$$25. \int_2^4 \frac{\sqrt{x^2 - 4}}{x^4} dx.$$

$$26. \int_0^2 \frac{dx}{(4 + x^2)\sqrt{4 + x^2}}.$$

$$27. \int_0^{\sqrt{2}} \frac{x^4 dx}{(4 - x^2)^{3/2}}.$$

$$28. \int_0^{1/\sqrt{2}} \frac{dx}{(1 - x^2)\sqrt{1 - x^2}}.$$

$$29. \int_0^1 \frac{x^2 dx}{\sqrt{4 - x^2}}.$$

$$30. \int_0^{3/2} \frac{x^2 dx}{\sqrt{9 - x^2}}.$$

Javoblar. 7.1 64ρ ; 7.2 $\frac{\rho}{16}$; 7.3 $\frac{\sqrt{2}}{50}$; 7.4 $\frac{\sqrt{3}}{15}$; 7.5 $\frac{\sqrt{3}}{8}$; 7.6 $\frac{5}{4} - \frac{3\rho}{8}$; 7.7 $\frac{\sqrt{3}}{4}$; 7.8 $\frac{5}{2} - \frac{3\rho}{4}$;

7.9 $\frac{4\rho}{3} - 2\sqrt{3}$; 7.10 ρ ; 7.11 $\frac{\sqrt{2}}{32}$; 7.12 16ρ ; 7.13 $\frac{25\rho}{12} - \frac{25\sqrt{3}}{8}$; 7.14 $\frac{625\rho}{16}$; 7.15 4ρ ;

7.16 $\frac{\sqrt{3}}{64}$; 7.17 $\frac{\sqrt{3}}{16}$; 7.18 $20 - 6\rho$; 7.19 $\frac{81\rho}{8}$; 7.20 $\frac{\sqrt{3} - \sqrt{2}}{2}$; 7.21 $\frac{\sqrt{3}}{48}$; 7.22 $10 - 3\rho$; 7.23 $\frac{\sqrt{3}}{72}$;

7.24 $\frac{\rho}{3} + \frac{\sqrt{3}}{2}$; 7.25 $\frac{\sqrt{3}}{32}$; 7.26 $\frac{\sqrt{2}}{8}$; 7.27 $5 - \frac{3\rho}{2}$; 7.28 1 ; 7.29 $\frac{\rho}{3} - \frac{\sqrt{3}}{2}$; 7.30 $\frac{3\rho}{4} - \frac{9\sqrt{3}}{8}$.

Yassi figuralar yuzlarini hisoblash

1. Uzluksiz $y = f(x)$ ($f(x) \geq 0$) egri chiziq, $x = a$, $x = b$ to'g'ri chiziqlar hamda Ox o'qning $[a, b]$ kesmasi bilan chegaralangan egri chizikli trapetsiyaning yuzi

$$S = \int_a^b f(x) dx$$


formula bilan hisoblanadi.

2. Uzluksiz $x = j(y)$ ($j(y) \geq 0$) egri chiziq, $y = c$, $y = d$ to'g'ri chiziqlar hamda Oy o'qning $[c, d]$ kesmasi bilan chegaralangan egri chizikli trapetsiyaning yuzi

$$S = \int_c^d j(y) dy$$

formula bilan hisoblanadi.

8–masala. $y = (x - 2)^3$, $y = 4x - 8$ chiziqlar bilan chegaralangan figuraning yuzini toping.



$$\begin{aligned}
 S &= 2 \int_0^2 (4x - 8 - (x - 2)^3) dx = 2 \int_0^2 (4x - 8 - x^3 + 6x^2 - 12x + 8) dx = \\
 &= 2 \int_0^2 (6x^2 - x^3 - 8x) dx = 2 \left(2x^3 - \frac{1}{4}x^4 - 4x^2 \right) \Big|_0^2 = 4 \times 2^3 - \frac{1}{2} \times 2^4 - 8 \times 2^2 = 8.
 \end{aligned}$$

1. $y = x\sqrt{9 - x^2}$, $y = 0$, $(0 \leq x \leq 3)$.

2. $y = 4 - x^2$, $y = x^2 - 2x$.

3. $y = \sin x \cos^2 x$, $y = 0$, $(0 \leq x \leq \frac{\pi}{2})$.

4. $y = \sqrt{4 - x^2}$, $y = 0$, $x = 0$, $x = 1$.

5. $y = x^2 \sqrt{4 - x^2}$, $y = 0$, $(0 \leq x \leq 2)$.

6. $y = \cos x \sin^2 x$, $y = 0$, $(0 \leq x \leq \frac{\pi}{2})$.

7. $y = \sqrt{e^x - 1}$, $y = 0$, $0 \leq x \leq \ln 2$.

8. $y = \frac{1}{x\sqrt{1 + \ln x}}$, $y = 0$, $x = 1$, $x = e^3$.

9. $y = \arccos x$, $y = 0$, $x = 0$.

10. $y = (x + 1)^2$, $y^2 = x + 1$.

11. $y = 2x - x^2 + 3$, $y = x^2 - 4x + 3$.

12. $y = x\sqrt{36 - x^2}$, $y = 0$, $(0 \leq x \leq 6)$.

13. $y = \arccos y$, $x = 0$, $y = 0$.

14. $y = x \arctg x$, $y = 0$, $x = \sqrt{3}$.

15. $y = x^2 \sqrt{8 - x^2}$, $y = 0$ $(0 \leq x \leq 2\sqrt{2})$.

16. $y = \sqrt{e^y - 1}$, $x = 0$, $y = \ln 2$.

17. $y = x\sqrt{4 - x^2}$, $y = 0$ $(0 \leq x \leq 2)$.

$$18. y = \frac{x}{1 + \sqrt{x}}, \quad y = 0, \quad x = 1.$$

$$19. y = \frac{1}{1 + \cos x}, \quad y = 0, \quad x = \frac{\rho}{2}, \quad x = -\frac{\rho}{2}.$$

$$20. x = (y - 2)^3, \quad x = 4y - 8.$$

$$21. y = \cos^5 x \sin 2x, \quad y = 0, \quad (0 \leq x \leq \frac{\rho}{2}).$$

$$22. y = \frac{x}{(x^2 + 1)^2}, \quad y = 0, \quad x = 1.$$

$$23. x = 4 - y^2, \quad x = y^2 - 2y.$$

$$24. x = \frac{1}{y\sqrt{1 + \ln y}}, \quad x = 0, \quad y = 1, \quad y = e^3.$$

$$25. y = \frac{e^{1/x}}{x^2}, \quad y = 0, \quad x = 2, \quad x = 1.$$

$$26. y = x^2 \sqrt{16 - x^2}, \quad y = 0 \quad (0 \leq x \leq 4).$$

$$27. x = \sqrt{4 - y^2}, \quad x = 0, \quad y = 0, \quad y = 1.$$

$$28. y = (x - 1)^2, \quad y^2 = x - 1.$$

$$29. y = x^2 \cos x, \quad y = 0, \quad (0 \leq x \leq \rho/2).$$

$$30. x = 4 - (y - 1)^2, \quad x = y^2 - 4y + 3.$$

$$\text{Javoblar. } 8.1 \ 9; 8.2 \ 9; 8.3 \ \frac{1}{3}; 8.4 \ \frac{\rho}{3} + \frac{\sqrt{3}}{2}; 8.5 \ \rho; 8.6 \ \frac{1}{3}; 8.7 \ \gg 0,43; 8.8 \ 2; 8.9 \ 1; 8.10 \ \frac{1}{3};$$

$$8.11 \ 9; 8.12 \ 72; 8.13 \ 1; 8.14 \ \frac{\rho}{\sqrt{3}} - \ln 2; 8.15 \ 4\rho; 8.16 \ \gg 0,429; 8.17 \ \frac{8}{3}; 8.18 \ \frac{5}{3} - 2\ln 2; 8.19 \ 2;$$

$$8.20 \ 8; 8.21 \ \frac{2}{7}; 8.22 \ \frac{1}{4}; 8.23 \ 9; 8.24 \ 2; 8.25 \ e - \sqrt{e}; 8.26 \ 16\rho; 8.27 \ \frac{\sqrt{3}}{2} + \frac{\rho}{3}; 8.28 \ \frac{1}{3};$$

$$8.29 \ \frac{\rho^2}{4} - 2; 8.30 \ 9.$$

Yassi figuralar yuzlarini hisoblash

3. Uzlüksiz $y = f_1(x)$ va $y = f_2(x)$ ($f_1(x) \leq f_2(x)$) egri chiziqlar hamda , $x = a$, $x = b$ ($a < b$) to'g'ri chiziqlar bilan chegaralangan figuraning yuzi

$$S = \int_a^b (f_2(x) - f_1(x)) dx$$

formula bilan hisoblanadi.


4. Uzlüksiz $y = j_1(x)$ va $y = j_2(x)$ ($j_1(x) \leq j_2(x)$) egri chiziqlar hamda , $y = c$, $y = d$ ($c < d$) to'g'ri chiziqlar bilan chegaralangan figuraning yuzi

$$S = \int_c^d (j_2(x) - j_1(x)) dx$$

formula bilan hisoblanadi.

9–masala. Quyidagi chiziqlar bilan chegaralangan shaklning yuzasini hisoblang.

$$\begin{aligned} \vec{r} x &= 2\sqrt{2} \cos^3 t, \\ \vec{r} y &= \sqrt{2} \sin^3 t, \\ x &= 1(x^3 - 1). \end{aligned}$$



$$S = \int_a^b y(t) x'(t) dt.$$

$$2\sqrt{2} \cos^3 t \cdot 1 \cdot (-\sin t) \Big|_{-\pi/4}^{\pi/4} = \frac{\rho}{4} + 2\rho n; \frac{\rho}{4} + 2\rho n \Big|_{-\pi/4}^{\pi/4}.$$

$$S = S_{ABCDE} - S_{ABDE} = \int_{-\pi/4}^{\pi/4} \sqrt{2} \sin^3 t \cdot 6\sqrt{2} \cos^2 t \cdot (-\sin t) dt = 1 \cdot 1 =$$

$$12 \int_{-\pi/4}^{\pi/4} \sin^4 t \cos^2 t dt - 1 = 12 \int_{-\pi/4}^{\pi/4} \frac{1}{8} (\cos 4t - 4 \cos 2t + 3) \cdot \frac{1}{2} (1 + \cos 2t) dt - 1 =$$

$$= \frac{12}{16} \int_{-\pi/4}^{\pi/4} (-\cos 4t - \frac{1}{2} \cos 2t + \frac{1}{2} \cos 6t + 1) dt - 1 =$$

$$= \frac{12}{16} \left[-\frac{1}{4} \sin 4t - \frac{1}{4} \sin 2t + \frac{1}{12} \sin 6t + t \right]_{-\pi/4}^{\pi/4} - 1 = 1,7.$$

$$\begin{aligned} 1. \quad & \begin{cases} \dot{x} = 4\sqrt{2} \cos^3 t, \\ \dot{y} = 2\sqrt{2} \sin^3 t, \\ x = 2(x^3 - 2). \end{cases} \end{aligned}$$

$$\begin{aligned} 2. \quad & \begin{cases} \dot{x} = \sqrt{2} \cos t, \\ \dot{y} = 2\sqrt{2} \sin t, \\ y = 2(y^3 - 2). \end{cases} \end{aligned}$$

$$\begin{aligned} 3. \quad & \begin{cases} \dot{x} = 4(t - \sin t), \\ \dot{y} = 4(1 - \cos t), \\ y = 4(0 < x < 8\rho, y^3 - 4). \end{cases} \end{aligned}$$

$$\begin{aligned} 4. \quad & \begin{cases} \dot{x} = 16 \cos^3 t, \\ \dot{y} = 2 \sin^3 t, \\ x = 2(x^3 - 2). \end{cases} \end{aligned}$$

$$\begin{aligned} \dot{x} &= 2 \cos t, \\ \dot{y} &= 6 \sin t, \\ y &= 3(y^3 - 3). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 16 \cos^3 t, \\ \dot{y} &= \sin^3 t, \\ x &= 6\sqrt{3}(x^3 - 6\sqrt{3}). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 3(t - \sin t), \\ \dot{y} &= 3(1 - \cos t), \\ y &= 3(0 < x < 6\rho, y^3 - 3). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 2\sqrt{2} \cos t, \\ \dot{y} &= 3\sqrt{2} \sin t, \\ y &= 3(y^3 - 3). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 32 \cos^3 t, \\ \dot{y} &= \sin^3 t, \\ x &= 4(x^3 - 4). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 6(t - \sin t), \\ \dot{y} &= 6(1 - \cos t), \\ y &= 6(0 < x < 12\rho, y^3 - 6). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 6 \cos^3 t, \\ \dot{y} &= 4 \sin^3 t, \\ y &= 2\sqrt{3}(y^3 - 2\sqrt{3}). \end{aligned}$$

$$\begin{aligned} \dot{x} &= \sqrt{2} \cos t, \\ \dot{y} &= 4\sqrt{2} \sin t, \\ y &= 4(y^3 - 4). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 2(t - \sin t), \\ \dot{y} &= 2(1 - \cos t), \\ y &= 3(0 < x < 4\rho, y^3 - 3). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 6 \cos t, \\ \dot{y} &= 2 \sin t, \\ y &= \sqrt{3}(y^3 - \sqrt{3}). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 8\sqrt{2} \cos^3 t, \\ \dot{y} &= \sqrt{2} \sin^3 t, \\ x &= 4(x^3 - 4). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 6(t - \sin t), \\ \dot{y} &= 6(t - \cos t), \\ y &= 9(0 < x < 12\rho, y^3 - 9). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 3 \cos t, \\ \dot{y} &= 8 \sin t, \\ y &= 4(y^3 - 4). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 8 \cos^3 t, \\ \dot{y} &= 4 \sin^3 t, \\ x &= 3\sqrt{3}(x^3 - 3\sqrt{3}). \end{aligned}$$

$$\begin{aligned} \dot{x} &= 10(t - \sin t), \\ \dot{y} &= 10(1 - \cos t), \\ y &= 15(0 < x < 20\rho, y^3 - 15). \end{aligned}$$

$$\begin{aligned} \dot{x} &= t - \sin t, \\ \dot{y} &= 1 - \cos t, \\ y &= 1(0 < x < 2\rho, y^3 - 1). \end{aligned}$$

$$\begin{aligned} 21. \begin{cases} \dot{x} = 8 \cos^3 t, \\ \dot{y} = 8 \sin^3 t, \end{cases} \\ x = 1(x^3 - 1). \end{aligned}$$

$$\begin{aligned} 22. \begin{cases} \dot{x} = 9 \cos t, \\ \dot{y} = 4 \sin t, \end{cases} \\ y = 2(y^3 - 2). \end{aligned}$$

$$\begin{aligned} 23. \begin{cases} \dot{x} = 8(t - \sin t), \\ \dot{y} = 8(1 - \cos t), \end{cases} \\ y = 12(0 < x < 16\rho, y^3 - 12). \end{aligned}$$

$$\begin{aligned} 24. \begin{cases} \dot{x} = 24 \cos^3 t, \\ \dot{y} = 2 \sin^3 t, \end{cases} \\ x = 9\sqrt{3}(x^3 - 9\sqrt{3}). \end{aligned}$$

$$\begin{aligned} 25. \begin{cases} \dot{x} = 3 \cos t, \\ \dot{y} = 8 \sin t, \end{cases} \\ x = 4\sqrt{3}(y^3 - 4\sqrt{3}). \end{aligned}$$

$$\begin{aligned} 26. \begin{cases} \dot{x} = 2(t - \sin t), \\ \dot{y} = 2(1 - \cos t), \end{cases} \\ y = 2(0 < x < 4\rho, y^3 - 2). \end{aligned}$$

$$\begin{aligned} 27. \begin{cases} \dot{x} = 4\sqrt{2} \cos^3 t, \\ \dot{y} = \sqrt{2} \sin^3 t, \end{cases} \\ x = 2(x^3 - 2). \end{aligned}$$

$$\begin{aligned} 28. \begin{cases} \dot{x} = 2\sqrt{2} \cos t, \\ \dot{y} = 5\sqrt{2} \sin t, \end{cases} \\ y = 5(y^3 - 5). \end{aligned}$$

$$\begin{aligned} 29. \begin{cases} \dot{x} = 4(t - \sin t), \\ \dot{y} = 4(1 - \cos t), \end{cases} \\ y = 6(0 < x < 8\rho, y^3 - 6). \end{aligned}$$

$$\begin{aligned} 30. \begin{cases} \dot{x} = 32 \cos^3 t, \\ \dot{y} = 3 \sin^3 t, \end{cases} \\ x = 12\sqrt{3}(x^3 - 12\sqrt{3}). \end{aligned}$$

Javoblar. 9.1 $\frac{3}{2}\rho - 2$; 9.2 $\rho - 2$; 9.3 $24\rho + 64$; 9.4 4ρ ; 9.5 $4\rho - 3\sqrt{3}$; 9.6 $2\sqrt{3}$;

9.7 ρ ; 9.8 $\gg 1, 12$; 9.9 $\frac{27}{2}\rho + 36$; 9.10 $\frac{3}{2}\rho + 2$; 9.11 $3\rho - 6$; 9.12 $36\rho + 81\sqrt{3}$;

9.13 $4\rho + 3\sqrt{3}$; 9.14 $8\rho - 6\sqrt{3}$; 9.15 $18\rho + 72$; 9.16 $2\rho - 3\sqrt{3}$; 9.17 $4\rho - 6\sqrt{3}$;

9.18 $100\rho + 225\sqrt{3}$; 9.19 $2\rho - 4$; 9.20 $\frac{\rho}{2} + 2$; 9.21 8ρ ; 9.22 $12\rho - 9\sqrt{3}$;

9.23 $48\sqrt{3}$; 9.24 $3\rho - \frac{9\sqrt{3}}{2}$; 9.25 $2, 174$; 9.26 $2\rho + 8$; 9.27 $\frac{3\rho}{4}$; 9.28 $5\rho - 10$;

9.29 $16\rho + 36\sqrt{3}$; 9.30 $6\rho - 9\sqrt{3}$.

Figuraning yuzini qutb koordinalar sistemasida hisoblash


Qutb koordinatalar sistemasida berilgan uzluksiz $r = r(j)$ egri chiziq va $j = a, j = b$ ($a < b$) nurlar bilan chegaralangan figuraning yuzi

$$S = \frac{1}{2} \int_a^b \dot{r}^2(j) dj$$

formula bilan hisoblanadi.

10–masala. Tenglamalari qutb koordinatalari sistemasida berilgan chiziqlar bilan chegaralangan shaklning yuzasini hisoblang.

$$r = 4 \cos j, \quad r = 2 \quad (r^3 = 2).$$




$$4 \cos 3j = 2,$$

$$\cos 3j = \frac{1}{2}.$$

Demak

$$-\frac{\rho}{3} + 2\rho n \leq 3j \leq \frac{\rho}{3} + 2\rho n, n \in \mathbb{Z}$$

$$-\frac{\rho}{9} + \frac{2\rho n}{3} \leq j \leq \frac{\rho}{9} + \frac{2\rho n}{3}, n \in \mathbb{Z},$$



$$S = \frac{1}{2} \int_a^b r^2(j) dj ,$$

$$\begin{aligned} S &= 6 \times \frac{1}{2} \int_{-p/3}^0 16 \cos^2 3j \, dj = 24 \int_{-p/3}^0 (1 + \cos 6j) \, dj = \\ &= 24 \left(j + \frac{1}{6} \sin 6j \right) \Big|_{-p/3}^0 = 24 \left(0 + 0 + \frac{p}{3} + \frac{1}{6} \times 0 \right) = 8p. \end{aligned}$$

1. $r = \cos 2j$.

2. $r = \sqrt{3} \cos j$, $r = \sin j$ ($0 \leq j \leq p/2$).

3. $r = 4 \sin 3j$, $r = 2$ ($r^3 = 2$).

4. $r = 2 \cos j$, $r = 2\sqrt{3} \sin j$ ($0 \leq j \leq p/2$).

5. $r = \sin 3j$.

6. $r = 6 \sin 3j$, $r = 3$ ($r^3 = 3$).

7. $r = \cos 3j$.

8. $r = \cos j$, $r = \sqrt{2} \sin(j - p/4)$ ($-p/4 \leq j \leq p/2$).

9. $r = \sin j$, $r = \sqrt{2} \cos(j - p/4)$ ($0 \leq j \leq 3p/4$).

10. $r = 6 \cos 3j$, $r = 3$ ($r^3 = 3$).

11. $r = \frac{1}{2} + \sin j$.

12. $r = \cos j$, $r = \sin j$ ($0 \leq j \leq p/2$).

$$13. r = \sqrt{2} \cos j - p/4, \quad r = \sqrt{2} \sin j - p/4 \quad (p/4 \leq j \leq 3p/4).$$

$$14. r = \cos j, \quad r = 2 \cos j.$$

$$15. r = \sin j, \quad r = 2 \sin j.$$

$$16. r = 1 + \sqrt{2} \cos j.$$

$$17. r = \frac{1}{2} + \cos j.$$

$$18. r = 1 + \sqrt{2} \sin j.$$

$$19. r = \frac{5}{2} \sin j, \quad r = \frac{3}{2} \sin j.$$

$$20. r = \frac{3}{2} \cos j, \quad r = \frac{5}{2} \cos j.$$

$$21. r = 4 \cos 4j.$$

$$22. r = \sin 6j.$$

$$23. r = 2 \cos j, \quad r = 3 \cos j.$$

$$24. r = \cos j + \sin j.$$

$$25. r = 2 \sin 4j.$$

$$26. r = 2 \cos 6j.$$

$$27. r = \cos j - \sin j.$$

$$28. r = 3 \sin j, \quad r = 5 \sin j.$$

$$29. r = 2 \sin j, \quad r = 4 \sin j.$$

$$30. r = 6 \sin j, \quad r = 4 \sin j.$$

Javoblar. 10.1 $\frac{p}{2}$; 10.2 $\frac{5p}{24} - \frac{\sqrt{3}}{4}$; 10.3 $\frac{4p}{3} + 2\sqrt{3}$; 10.4 $\frac{5p}{6} - \sqrt{3} \gg 0,88$; 10.5 $\frac{p}{4}$;

10.6 $3p + \frac{9\sqrt{3}}{2}$; 10.7 $\frac{p}{4} \gg 0,78$; 10.8 $\frac{p}{4} - \frac{1}{4}$; 10.9 $\frac{p-1}{4}$; 10.10 $\frac{9p}{2}$; 10.11 $\frac{3p}{4}$; 10.12 $\frac{1}{2}$;

10.13 $\frac{p+2}{4}$; 10.14 $\frac{3p}{4}$; 10.15 $\frac{3p}{4}$; 10.16 11,94; 10.17 $\frac{3p}{4}$; 10.18 $2p$; 10.19 p ; 10.20 p ;

10.21 8ρ ; 10.22 $\frac{\rho}{4}$; 10.23 $\frac{5\rho}{4}$; 10.24 $\frac{\rho}{2}$; 10.25 $\frac{\rho}{2}$; 10.26 2ρ ; 10.27 $\frac{\rho}{2}$; 10.28 4ρ ; 10.29 3ρ ;
10.30 5ρ .

Yoy uzunligini hisoblash

Agar $y = f(x)$ funksiya $[a, b]$ da silliq egri chiziq (ya'ni $f'(x)$ uzluksiz) bo'lsa, u holda uning yoyi uzunligi

$$l = \int_a^b \sqrt{1 + y'^2} dx$$

formula bilan hisoblanadi. Bunda a va b yoy uchlarining absissalaridir ($a < b$).

Agar egri chiziq $x = j(y)$ ($c \leq y \leq d$) ko'rinishda berilgan bo'lsa, yoy uzunligi

$$l = \int_c^d \sqrt{1 + x'^2} dy$$

formula bilan hisoblanadi.

11–masala. Tenglamalari to'g'ri burchakli koordinatalar sistemasida berilgan egri chiziq yoyining uzunligini hisoblang.

$$y = \sqrt{1 - x^2} + \arcsin x, \quad 0 \leq x \leq \frac{8}{9}.$$

$$y' = -\frac{x}{\sqrt{1 - x^2}} - \frac{1}{\sqrt{1 - x^2}} = \frac{-1 - x}{\sqrt{1 - x^2}}.$$

$$l = \int_a^b \sqrt{1 + (y')^2} dx,$$

$$\begin{aligned}
 I &= \int_0^{8/9} \sqrt{1 + \frac{(x+1)^2}{1-x^2}} dx = \int_0^{8/9} \sqrt{\frac{1-x^2+x^2+2x+1}{1-x^2}} dx = \int_0^{8/9} \sqrt{\frac{2+2x}{1-x^2}} dx = \\
 &= \int_0^{8/9} \sqrt{\frac{2}{1-x}} dx = \sqrt{2} \int_0^{8/9} \frac{dx}{\sqrt{1-x}} = \left. \sqrt{2} \sqrt{1-x} \right|_0^{8/9} = \\
 &= -2(\sqrt{2/9} - \sqrt{2}) = -2\sqrt{2/9} + 2\sqrt{2} = \frac{4\sqrt{2}}{3}.
 \end{aligned}$$

1. $y = \ln x, \quad \sqrt{3} \leq x \leq \sqrt{15}.$

2. $y = \frac{x^2}{4} - \frac{\ln x}{2}, \quad 1 \leq x \leq 2.$

3. $y = \sqrt{1-x^2} + \arcsin x, \quad 0 \leq x \leq \frac{7}{9}.$

4. $y = \ln \frac{5}{2x}, \quad \sqrt{3} \leq x \leq \sqrt{8}.$

5. $y = -\ln \cos x, \quad 0 \leq x \leq \pi/6.$

6. $y = e^x + 6, \quad \ln \sqrt{8} \leq x \leq \ln \sqrt{15}.$

7. $y = 2 + \arcsin \sqrt{x} + \sqrt{x-x^2}, \quad \frac{1}{4} \leq x \leq 1.$

8. $y = \ln(x^2 - 1), \quad 2 \leq x \leq 3.$

9. $y = \ln(1-x^2), \quad 0 \leq x \leq \frac{1}{4}.$

10. $y = 2 + \cosh x, \quad 0 \leq x \leq 1.$

11. $y = 1 - \ln \cos x, \quad 0 \leq x \leq \pi/6.$

12. $y = e^x + 13, \quad \ln \sqrt{15} \leq x \leq \ln \sqrt{24}.$

13. $y = -\arccos \sqrt{x} + \sqrt{x-x^2}, \quad 0 \leq x \leq \frac{1}{4}.$

14. $y = 2 - e^x, \quad \ln \sqrt{3} \leq x \leq \ln \sqrt{8}.$

$$15. y = \arcsin x - \sqrt{1 - x^2}, \quad 0 \leq x \leq \frac{15}{16}.$$

$$16. y = 1 - \ln \sin x, \quad \rho/3 \leq x \leq \rho/2.$$

$$17. y = 1 - \ln(x^2 - 1), \quad 3 \leq x \leq 4.$$

$$18. y = \sqrt{x - x^2} - \arccos \sqrt{x} + 5, \quad \frac{1}{9} \leq x \leq 1.$$

$$19. y = -\arccos x + \sqrt{1 - x^2} + 1, \quad 0 \leq x \leq \frac{9}{16}.$$

$$20. y = \ln \sin x, \quad \rho/3 \leq x \leq \rho/2.$$

$$21. y = \ln 7 - \ln x, \quad \sqrt{3} \leq x \leq \sqrt{8}.$$

$$22. y = chx + 3, \quad 0 \leq x \leq 1.$$

$$23. y = 1 + \arcsin x - \sqrt{1 - x^2}, \quad 0 \leq x \leq \frac{3}{4}.$$

$$24. y = \ln \cos x + 2, \quad 0 \leq x \leq \rho/6.$$

$$25. y = e^x + 26, \quad \ln \sqrt{8} \leq x \leq \ln \sqrt{24}.$$

$$26. y = \frac{e^x + e^{-x}}{2} + 3, \quad 0 \leq x \leq 2.$$

$$27. y = \arccos \sqrt{x} - \sqrt{x - x^2} + 4, \quad 0 \leq x \leq \frac{1}{2}.$$

$$28. y = \frac{e^{2x} + e^{-2x} + 3}{3}, \quad 0 \leq x \leq 2.$$

$$29. y = e^x + e, \quad \ln \sqrt{3} \leq x \leq \ln \sqrt{15}.$$

$$30. y = \frac{1 - e^x - e^{-x}}{2}, \quad 0 \leq x \leq 3.$$

Javoblar. 11.1 $\frac{1}{2} \ln \frac{9}{5} + 2$; 11.2 $\frac{3}{4} + \frac{1}{2} \ln 2$; 11.3 $\frac{2\sqrt{2}}{3}$; 11.4 $1 + \frac{1}{2} \ln \frac{3}{2}$; 11.5 $\ln \sqrt{3}$;

$$\begin{aligned}
& 11.6 \ 1 + \frac{1}{2} \ln \frac{6}{5}; 11.7 \ 1 + \ln \frac{3}{2}; 11.8 \ 1; 11.9 \ln \frac{5}{3} - \frac{1}{4}; 11.10 \ \text{sh}1; 11.11 \ \ln \sqrt{3}; 11.12 \ 1 + \frac{1}{2} \ln \frac{10}{9}; \\
& 11.13 \ 1; 11.14 \ 1 + \frac{1}{2} \ln \frac{3}{2}; 11.15 \ \frac{3}{\sqrt{2}}; 11.16 \ \frac{\ln 3}{2}; 11.17 \ 1 + \ln \frac{6}{5}; 11.18 \ \frac{4}{3}; 11.19 \ \frac{1}{\sqrt{2}}; 11.20 \ \frac{1}{2} \ln 3; \\
& 11.21 \ 1 + \frac{1}{2} \ln \frac{3}{2}; 11.22 \ \text{sh}1; 11.23 \ \sqrt{2}; 11.24 \ \ln \sqrt{3}; 11.25 \ 2 + \frac{1}{2} \ln \frac{4}{3}; 11.26 \ \frac{1}{2} (e^2 - e^{-2}); \\
& 11.27 \ \sqrt{2}; 11.28 \ \frac{1}{2} (e^4 - e^{-4}); 11.29 \ 2 + \frac{1}{2} \ln \frac{9}{5}; 11.30 \ \frac{1}{2} (e^3 - e^{-3}).
\end{aligned}$$

Parametrik ko'rinishda berilgan egri chiziq yoyining uzunligi

Agar egri chiziq

$$\begin{aligned}
x &= x(t) \\
y &= y(t)
\end{aligned}
\quad a \leq t \leq b$$

ko'rinishida berilgan bo'lib, $x(t)$, $y(t)$ uzluksiz funksiyalar bo'lsa, u holda egri chiziq yoyining uzunligi

$$l = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt$$

Formula bilan hisoblanadi. Bunda a va b lar t parametrning yoy uchlariga mos qiymatlaridir ($a < b$).

12–masala. Parametrik ko'rinishda berilgan egri chiziq yoyining uzunligini hisoblang.

$$\begin{aligned}
x &= 4(\cos t + t \sin t), \\
y &= 4(\sin t - t \cos t), \\
0 &\leq t \leq 2\pi.
\end{aligned}$$

$$\begin{aligned}
x' &= 4(-\sin t + \sin t + t \cos t) = 4t \cos t, \\
y' &= 4(\cos t - \cos t + t \sin t) = 4t \sin t.
\end{aligned}$$

$$l = \int_a^b \sqrt{(x')^2 + (y')^2} dt,$$

$$l = \int_0^{2p} \sqrt{16t^2 \cos^2 t + 16t^2 \sin^2 t} dt = \int_0^{2p} 4t dt = 2t^2 \Big|_0^{2p} = 2 \times (2p)^2 = 8p^2.$$

$$\begin{aligned} 1. \quad & \begin{cases} \dot{x} = 5(t - \sin t), \\ \dot{y} = 5(1 - \cos t), \end{cases} \\ & 0 \leq t \leq p. \end{aligned}$$

$$\begin{aligned} 2. \quad & \begin{cases} \dot{x} = 3(2 \cos t - \cos 2t), \\ \dot{y} = 3(2 \sin t - \sin 2t), \end{cases} \\ & 0 \leq t \leq 2p. \end{aligned}$$

$$\begin{aligned} 3. \quad & \begin{cases} \dot{x} = (t^2 - 2) \sin t + 2t \cos t, \\ \dot{y} = (2 - t^2) \cos t + 2t \sin t, \end{cases} \\ & 0 \leq t \leq p. \end{aligned}$$

$$\begin{aligned} 4. \quad & \begin{cases} \dot{x} = 10 \cos^3 t, \\ \dot{y} = 10 \sin^3 t, \end{cases} \\ & 0 \leq t \leq p/2. \end{aligned}$$

$$\begin{aligned} 5. \quad & \begin{cases} \dot{x} = e^t (\cos t + \sin t), \\ \dot{y} = e^t (\cos t - \sin t), \end{cases} \\ & 0 \leq t \leq p. \end{aligned}$$

$$\begin{aligned} 6. \quad & \begin{cases} \dot{x} = 3(t - \sin t), \\ \dot{y} = 3(t - \cos t), \end{cases} \\ & p \leq t \leq 2p. \end{aligned}$$

$$\begin{aligned} 7. \quad & \begin{cases} \dot{x} = \frac{1}{2} \cos t - \frac{1}{4} \cos 2t, \\ \dot{y} = \frac{1}{2} \sin t - \frac{1}{4} \sin 2t, \end{cases} \\ & p/2 \leq t \leq 2p/3. \end{aligned}$$

$$\begin{aligned} 8. \quad & \begin{cases} \dot{x} = 3(\cos t + t \sin t), \\ \dot{y} = 3(\sin t - t \cos t), \end{cases} \\ & 0 \leq t \leq p/3. \end{aligned}$$

$$\begin{aligned} 9. \quad & \begin{cases} \dot{x} = (t^2 - 2) \sin t + 2t \cos t, \\ \dot{y} = (2 - t^2) \cos t + 2t \sin t, \end{cases} \\ & 0 \leq t \leq p/3. \end{aligned}$$

$$\begin{aligned} 10. \quad & \begin{cases} \dot{x} = 6 \cos^3 t, \\ \dot{y} = 6 \sin^3 t, \end{cases} \\ & 0 \leq t \leq p/3. \end{aligned}$$

$$\begin{aligned} 11. \quad & \begin{cases} \dot{x} = e^t (\cos t + \sin t), \\ \dot{y} = e^t (\cos t - \sin t), \end{cases} \\ & p/2 \leq t \leq p. \end{aligned}$$

$$\begin{aligned} 12. \quad & \begin{cases} \dot{x} = 2,5(t - \sin t), \\ \dot{y} = 2,5(1 - \cos t), \end{cases} \\ & p/2 \leq t \leq p. \end{aligned}$$

$$\begin{aligned} 13. \quad & \begin{cases} \dot{x} = 3,5(2 \cos t - \cos 2t), \\ \dot{y} = 3,5(2 \sin t - \sin 2t), \end{cases} \\ & 0 \leq t \leq p/2. \end{aligned}$$

$$\begin{aligned} 14. \quad & \begin{cases} \dot{x} = 6(\cos t + t \sin t), \\ \dot{y} = 6(\sin t - t \cos t), \end{cases} \\ & 0 \leq t \leq p. \end{aligned}$$

$$\begin{aligned} \dot{x} &= (t^2 - 2) \sin t + 2t \cos t, \\ 15. \dot{y} &= (2 - t^2) \cos t + 2t \sin t, \\ 0 \leq t \leq p/2. \end{aligned}$$

$$\begin{aligned} \dot{x} &= e^t (\cos t + \sin t), \\ 17. \dot{y} &= e^t (\cos t - \sin t), \\ 0 \leq t \leq 2p. \end{aligned}$$

$$\begin{aligned} \dot{x} &= 2(2 \cos t - \cos 2t), \\ 19. \dot{y} &= 2(2 \sin t - \sin 2t), \\ 0 \leq t \leq p/3. \end{aligned}$$

$$\begin{aligned} \dot{x} &= (t^2 - 2) \sin t + 2t \cos t, \\ 21. \dot{y} &= (2 - t^2) \cos t + 2t \sin t, \\ 0 \leq t \leq 2p. \end{aligned}$$

$$\begin{aligned} \dot{x} &= e^t (\cos t + \sin t), \\ 23. \dot{y} &= e^t (\cos t - \sin t), \\ 0 \leq t \leq 3p/2. \end{aligned}$$

$$\begin{aligned} \dot{x} &= 4(2 \cos t - \cos 2t), \\ 25. \dot{y} &= 4(2 \sin t - \sin 2t), \\ 0 \leq t \leq p. \end{aligned}$$

$$\begin{aligned} \dot{x} &= (t^2 - 2) \sin t + 2t \cos t, \\ 27. \dot{y} &= (2 - t^2) \cos t + 2t \sin t, \\ 0 \leq t \leq 3p. \end{aligned}$$

$$\begin{aligned} \dot{x} &= e^t (\cos t + \sin t), \\ 29. \dot{y} &= e^t (\cos t - \sin t), \\ p/6 \leq t \leq p/4. \end{aligned}$$

$$\begin{aligned} \dot{x} &= 8 \cos^3 t, \\ 16. \dot{y} &= 8 \sin^3 t, \\ 0 \leq t \leq p/6. \end{aligned}$$

$$\begin{aligned} \dot{x} &= 4(t - \sin t), \\ 18. \dot{y} &= 4(1 - \cos t), \\ p/2 \leq t \leq 2p/3. \end{aligned}$$

$$\begin{aligned} \dot{x} &= 8(\cos t + t \sin t), \\ 20. \dot{y} &= 8(\sin t - t \cos t), \\ 0 \leq t \leq p/4. \end{aligned}$$

$$\begin{aligned} \dot{x} &= 4 \cos^3 t, \\ 22. \dot{y} &= 4 \sin^3 t, \\ p/6 \leq t \leq p/4. \end{aligned}$$

$$\begin{aligned} \dot{x} &= 2(t - \sin t), \\ 24. \dot{y} &= 2(1 - \cos t), \\ 0 \leq t \leq p/2. \end{aligned}$$

$$\begin{aligned} \dot{x} &= 2(\cos t + t \sin t), \\ 26. \dot{y} &= 2(\sin t - t \cos t), \\ 0 \leq t \leq p/2. \end{aligned}$$

$$\begin{aligned} \dot{x} &= 2 \cos^3 t, \\ 28. \dot{y} &= 2 \sin^3 t, \\ 0 \leq t \leq p/4. \end{aligned}$$

$$\begin{aligned} \dot{x} &= (t^2 - 2) \sin t + 2t \cos t, \\ 30. \dot{y} &= (2 - t^2) \cos t + 2t \sin t, \\ 0 \leq t \leq p. \end{aligned}$$

Javoblar. 12.1 20; 12.2 48; 12.3 $\frac{p^3}{3}$; 12.4 15; 12.5 $2(e^p - 1)$; 12.6 12; 12.7 $\sqrt{2} - 1$; 12.8 $\frac{p^2}{6}$;

$$\begin{aligned}
&12.9 \frac{\rho^3}{81}; 12.10 \frac{27}{4}; 12.11 2(e^p - e^{p/2}); 12.12 5\sqrt{2}; 12.13 14(2 - \sqrt{2}); 12.14 3\rho^2; \\
&12.15 \frac{\rho^3}{24}; 12.16 3; 12.17 2(e^{2p} - 1); 12.18 8(\sqrt{2} - 1); 12.19 8(2 - \sqrt{3}); 12.20 \frac{\rho^2}{4}; \\
&12.21 \frac{8\rho^3}{3}; 12.22 + \infty; 12.23 2(e^{3p/2} - 1); 12.24 4(2 - \sqrt{2}); 12.25 32; 12.26 \frac{\rho^2}{4}; \\
&12.27 9\rho^3; 12.28 \frac{3}{2}; 12.29 2(e^{p/4} - e^{p/6}); 12.30 \frac{\rho^3}{3}.
\end{aligned}$$

Qutb koordinatalari sistemasida berilgan egri chiziq yoyining uzunligi

Qutb koordinalar sistemasida berilgan silliq egri chiziq

$r = f(j)$, $j_0 \leq j \leq j_1$ yoyining uzunligi

$$l = \int_{j_0}^{j_1} \sqrt{r^2 + r'^2} dj$$

orqali hisoblanadi. Bunda j_0 va j_1 —qutb burchagi j ning yoy uchlaridagi qiymatlari ($j_0 < j_1$).

13–masala. Tenglamalari qutb koordinatalari sistemasida berilgan egri chiziq yoyining uzunligini hisoblang.

$$\begin{aligned}
r &= 2e^{4j/3}, \\
-p/2 &\leq j \leq p/2.
\end{aligned}$$

$$\begin{aligned}
L &= \int_a^b \sqrt{r^2 + (r')^2} dj; \\
r' &= \frac{8}{3}e^{4j/3}.
\end{aligned}$$

$$L = \int_{-p/2}^{p/2} \sqrt{4e^{8j/3} + \frac{64}{9}e^{8j/3}} dj = \int_{-p/2}^{p/2} \sqrt{\frac{100}{9}e^{8j/3}} dj = \int_{-p/2}^{p/2} \frac{10}{3}e^{4j/3} dj =$$

$$= \frac{10}{3} \times \frac{3}{4} e^{4j/3} \Big|_{-p/2}^{p/2} = \frac{5}{2} \left(e^{\frac{2p}{3}} - e^{-\frac{2p}{3}} \right) = 5 \sinh \frac{2p}{3}$$

$$\begin{array}{lll} 1. & r = 3e^{3j/4}, & \\ & -p/2 \leq j \leq p/2. & \end{array}$$

$$\begin{array}{lll} 2. & r = \sqrt{2}e^j, & \\ & -p/2 \leq j \leq p/2. & \end{array}$$

$$\begin{array}{lll} 3. & r = 5e^{5j/12}, & \\ & -p/2 \leq j \leq p/2. & \end{array}$$

$$\begin{array}{lll} 4. & r = 6e^{12j/5}, & \\ & -p/2 \leq j \leq p/2. & \end{array}$$

$$\begin{array}{lll} 5. & r = 3e^{3j/4}, & \\ & 0 \leq j \leq p/3. & \end{array}$$

$$\begin{array}{lll} 6. & r = 4e^{4j/3}, & \\ & 0 \leq j \leq p/3. & \end{array}$$

$$\begin{array}{lll} 7. & r = \sqrt{2}e^j, & \\ & 0 \leq j \leq p/3. & \end{array}$$

$$\begin{array}{lll} 8. & r = 5e^{5j/12}, & \\ & 0 \leq j \leq p/3. & \end{array}$$

$$\begin{array}{lll} 9. & r = 12e^{12j/5}, & \\ & 0 \leq j \leq p/3. & \end{array}$$

$$\begin{array}{lll} 10. & r = 1 - \sin j, & \\ & -p/2 \leq j \leq -p/6. & \end{array}$$

$$\begin{array}{lll} 11. & r = 2(1 - \cos j), & \\ & -p \leq j \leq -p/2. & \end{array}$$

$$\begin{array}{lll} 12. & r = 3(1 + \sin j), & \\ & -p/6 \leq j \leq 0. & \end{array}$$

$$\begin{array}{lll} 13. & r = 4(1 - \sin j), & \\ & 0 \leq j \leq p/6. & \end{array}$$

$$\begin{array}{lll} 14. & r = 5(1 - \cos j), & \\ & -p/3 \leq j \leq 0. & \end{array}$$

$$\begin{array}{lll} 15. & r = 6(1 + \sin j), & \\ & -p/2 \leq j \leq 0. & \end{array}$$

$$\begin{array}{lll} 16. & r = 7(1 - \sin j), & \\ & -p/6 \leq j \leq p/6. & \end{array}$$

$$\begin{array}{lll} 17. & r = 8(1 - \cos j), & \\ & -2p/3 \leq j \leq 0. & \end{array}$$

$$\begin{array}{lll} 18. & r = 2j, & \\ & 0 \leq j \leq 3/4. & \end{array}$$

$$\begin{array}{lll} 19. & r = 2j, & \\ & 0 \leq j \leq 4/3. & \end{array}$$

$$\begin{array}{lll} 20. & r = 2j, & \\ & 0 \leq j \leq \frac{5}{12}. & \end{array}$$

$$\begin{array}{lll} 21. & r = 2j, & \\ & 0 \leq j \leq \frac{12}{5}. & \end{array}$$

$$22. \quad r = 4j, \\ 0 \leq j \leq 3/4.$$

$$23. \quad r = 3j, \\ 0 \leq j \leq 4/3.$$

$$24. \quad r = 5j, \\ 0 \leq j \leq \frac{12}{5}.$$

$$25. \quad r = 2 \cos j, \\ 0 \leq j \leq p/6.$$

$$26. \quad r = 8 \cos j, \\ 0 \leq j \leq p/4.$$

$$27. \quad r = 6 \cos j, \\ 0 \leq j \leq p/3.$$

$$28. \quad r = 2 \sin j, \\ 0 \leq j \leq p/6.$$

$$29. \quad r = 8 \sin j, \\ 0 \leq j \leq p/4.$$

$$30. \quad r = 6 \sin j, \\ 0 \leq j \leq p/3.$$

Javoblar. 13.1 $10 \times \operatorname{sh} \frac{3p}{8}$; 13.2 $4 \times \operatorname{sh} \frac{p}{2}$; 13.3 $26 \times \operatorname{sh} \frac{5p}{24}$; 13.4 $13 \times \operatorname{sh} \frac{6p}{5}$; 13.5 $5 \times (e^{p/4} - 1)$;

13.6 $\frac{5}{3} \times (e^{4p/9} - 1)$; 13.7 $2 \times (e^{p/3} - 1)$; 13.8 $13 \times (e^{5p/36} - 1)$; 13.9 $13 \times (e^{4p/5} - 1)$;

13.10 2 ; 13.11 $-4\sqrt{2}$; 13.12 $6(\sqrt{3} - \sqrt{2})$; 13.13 $8(\sqrt{3} - \sqrt{2})$; 13.14 $20\sqrt{\frac{3}{4}}$;

13.15 $12(2 - \sqrt{2})$; 13.16 $10,249$; 13.17 16 ; 13.18 $\frac{j}{2} \sqrt{j^2 + 1} + \frac{1}{2} \ln |j + \sqrt{j^2 + 1}|$;

13.19 $\frac{20}{9} + \ln 3$; 13.20 $\frac{65}{144} + \ln \frac{3}{2}$; 13.21 $\frac{156}{25} + 5$; 13.22 $\frac{15}{8} + \ln 4$; 13.23 $\frac{10}{3} + \frac{3}{2} \ln 3$;

13.24 $\frac{78}{5} + \frac{5}{2} \ln 5$; 13.25 $\frac{p}{3}$; 13.26 $2p$; 13.27 $2p$; 13.28 $\frac{p}{3}$; 13.29 $2p$; 13.30 $2p$.

Aylanma jism sirining yuzi

1. $y = f(x)$ ($a \leq x \leq b$) silliq egri chiziq yoyining Ox o'qi atrofida aylanishidan hosil bo'lgan jism sirining yuzi

$$S = 2\pi \int_a^b y \sqrt{1 + y'^2} dx$$

formula bilan hisoblanadi.

2. Agar silliq egri chiziq

$$\begin{aligned} x &= x(t) \\ y &= y(t) \end{aligned} \quad a \leq t \leq b$$

Parametrik ko'rinishda berilgan bo'lsa, sirt yuzi

$$S = 2\pi \int_a^b y(t) \sqrt{x'^2 + y'^2} dt$$

formula bilan hisoblanadi.

3. Agar silliq egri chiziqli qutb koordinatalar sistemasida

$$r = f(\varphi), \quad \varphi_0 \leq \varphi \leq \varphi_1$$

ko'rinishda berilgan bo'lsa, uning qutb o'qi atrofida aylanishidan hosil bo'lgan jism sirtining yuzi

$$S = 2\pi \int_{\varphi_0}^{\varphi_1} r^2 \sin \varphi \sqrt{r'^2 + r^2} d\varphi$$

formula bilan hisoblanadi.

Jismlarning hajmini ularning ko'ndalang kesimlari bo'yicha hisoblash

Agar ko'ndalang kesim yuzi S ni Ox o'qqa perpendikulyar tekislik orqali x ning funksiyasi sifatida $S = S(x)$ kabi ifodalash mumkin bo'lsa, u holda hajm differensial uchun asosi S , balandligi dx bo'lgan silindrning hajmi olinadi, ya'ni


$$dV = S(x)dx.$$

$[a, b]$ intervalda integral olib, berilgan jismning hajmini hisoblash uchun formula hosil qilamiz:

$$V = \int_a^b S(x)dx.$$

14–masala. Quyidagi sirtlar bilan chegaralangan jismning hajmini toping.

$$\frac{x^2}{9} + \frac{y^2}{16} - \frac{z^2}{64} = -1, \quad z = 16.$$



Jismning ko'ndalang kesimida $z = \text{const}$ ellips hosil bo'ladi:

$$\frac{x^2}{9} + \frac{y^2}{16} = \frac{z^2}{64} - 1.$$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \text{ ko'rinishidagi ellipsning yuzi } p \times a \times b \text{ ga teng.}$$

Ellipsning radiusini topamiz:

$$\frac{x^2}{9 \times \frac{z^2 - 64}{64}} + \frac{y^2}{16 \times \frac{z^2 - 64}{64}} = 1$$

$$a = \frac{3}{8} \sqrt{z^2 - 64}; \quad b = \frac{1}{2} \sqrt{z^2 - 64}$$

$$S = pab = p \times \frac{3}{8} \times \sqrt{z^2 - 64} = \frac{3p}{8} \sqrt{z^2 - 64}.$$

$$\begin{aligned}
 V &= \oint_{\gamma} S(z) dz = \frac{3\rho}{16} \oint_{\gamma} (z^2 - 64) dz = \frac{3\rho}{16} \oint_{\gamma} z^3 - 64z = \\
 &= \frac{3\rho}{16} \oint_{\gamma} z^3 - 64 \oint_{\gamma} 1 = \\
 &= \frac{\rho}{16} \oint_{\gamma} z^3 - 3 \oint_{\gamma} 1 = \\
 &= \frac{\rho}{16} (4096 - 3072 - 512 + 1536) = 128\rho.
 \end{aligned}$$

$$1. \frac{x^2}{9} + y^2 = 1, \quad z = y, \quad z = 0 \quad (y^3 = 0).$$

$$2. z = x^2 + 4y^2, \quad z = 2.$$

$$3. \frac{x^2}{9} + \frac{y^2}{4} - z^2 = 1, \quad z = 0, \quad z = 3.$$

$$4. \frac{x^2}{9} + \frac{y^2}{4} - \frac{z^2}{36} = -1, \quad z = 12.$$

$$5. \frac{x^2}{16} + \frac{y^2}{9} + \frac{z^2}{4} = -1, \quad z = 1, \quad z = 0.$$

$$6. x^2 + y^2 = 9, \quad z = y, \quad z = 0 \quad (y^3 = 0).$$

$$7. z = x^2 + 9y^2, \quad z = 3.$$

$$8. \frac{x^2}{4} + y^2 - z^2 = 1, \quad z = 0, \quad z = 3.$$

$$9. \frac{x^2}{16} + \frac{y^2}{9} + \frac{z^2}{64} = 1, \quad z = 4, \quad z = 0.$$

$$10. \frac{x^2}{16} + \frac{y^2}{9} + \frac{z^2}{16} = 1, \quad z = 2, \quad z = 0.$$

$$11. \frac{x^2}{3} + \frac{y^2}{4} = 1, \quad z = y\sqrt{3}, \quad z = 0 \quad (y \neq 0).$$

$$12. z = 2x^2 + 8y^2, \quad z = 4.$$

$$13. \frac{x^2}{81} + \frac{y^2}{25} - z^2 = 1, \quad z = 0, \quad z = 2.$$

$$14. \frac{x^2}{4} + \frac{y^2}{9} - \frac{z^2}{36} = -1, \quad z = 12.$$

$$15. \frac{x^2}{16} + \frac{y^2}{9} + \frac{z^2}{36} = 1, \quad z = 3, \quad z = 0.$$

$$16. \frac{x^2}{3} + \frac{y^2}{16} = 1, \quad z = y\sqrt{3}, \quad z = 0 \quad (y \neq 0).$$

$$17. z = x^2 + 5y^2, \quad z = 5.$$

$$18. \frac{x^2}{9} + \frac{y^2}{4} - z^2 = 1, \quad z = 0, \quad z = 4.$$

$$19. \frac{x^2}{9} + \frac{y^2}{25} - \frac{z^2}{100} = -1, \quad z = 20.$$

$$20. \frac{x^2}{27} + \frac{y^2}{25} = 1, \quad z = \frac{y}{\sqrt{3}}, \quad z = 0 \quad (y \neq 0).$$

$$21. z = 4x^2 + 9y^2, \quad z = 6.$$

$$22. x^2 + \frac{y^2}{4} - z^2 = 1, \quad z = 0, \quad z = 3.$$

$$23. \frac{x^2}{25} + \frac{y^2}{9} - \frac{z^2}{100} = -1, \quad z = 20.$$

$$24. \frac{x^2}{16} + \frac{y^2}{9} + \frac{z^2}{100} = 1, \quad z = 5, \quad z = 0.$$

$$25. \frac{x^2}{27} + y^2 = 1, \quad z = \frac{y}{\sqrt{3}}, \quad z = 0 \quad (y \neq 0).$$

26. $z = 2x^2 + 18y^2, \quad z = 6.$

27. $\frac{x^2}{25} + \frac{y^2}{9} - z^2 = 1, \quad z = 0, \quad z = 2.$

28. $\frac{x^2}{16} + \frac{y^2}{9} - \frac{z^2}{64} = -1, \quad z = 16.$

29. $\frac{x^2}{16} + \frac{y^2}{9} + \frac{z^2}{144} = 1, \quad z = 6, \quad z = 0.$

30. $\frac{x^2}{16} + \frac{y^2}{9} + \frac{z^2}{196} = 1, \quad z = 7, \quad z = 0.$

Javoblar. 14.1 2; 14.2 p ; 14.3 $72p$; 14.4 $48p$; 14.5 $11p$; 14.6 26; 14.7 $\frac{3p}{2}$; 14.8 $12p$;

14.9 $44p$; 14.10 $22p$; 14.11 32; 14.12 $2p$; 14.13 $210p$; 14.14 $48p$; 14.15 $33p$;

14.16 32; 14.17 $\frac{5p\sqrt{5}}{2}$; 14.18 $152p$; 14.19 $200p$; 14.20 1250; 14.21 $3p$; 14.22 $24p$;

14.23 $200p$; 14.24 $55p$; 14.25 2; 14.26 $3p$; 14.27 $70p$; 14.28 $128p$; 14.29 $66p$;

14.30 $77p$.

To'g'ri burchakli dekart koordinatalar sistemasida aylanma jism hajmi

1. $y = f(x)$ egri chiziqlik Ox o'q va $x = a$, $x = b$ to'g'ri chiziqlar bilan

chegaralangan egri chiziqli trapetsiyaning Ox o'q atrofida aylanishidan hosil bo'lgan jism hajmi

$$V = \rho \int_a^b (f(x))^2 dx = \rho \int_a^b y^2 dx$$

formula bilan hisoblandi.

2. $x = f(y)$ egri chiziq Oy o'q va $y = c$, $y = d$ to'g'ri chiziqlar bilan chegaralangan egri chiziqli trapetsiyaning Oy o'q atrofida aylanishidan hosil bo'lgan jism hajmi

$$V = \rho \int_c^d (f(y))^2 dy = \rho \int_c^d x^2 dy$$

formula bilan hisoblandi.

3. $y = f(x)$ egri chiziq Ox o'q va $x = a$, $y = b$ to'g'ri chiziqlar bilan chegaralangan egri chiziqli trapetsiyaning Oy o'q atrofida aylanishidan hosil bo'lgan jism hajmi

$$V = 2\rho \int_a^b xy dx$$

formula bilan hisoblandi.

4. Umumiy holda, $y = f_1(x)$, $y = f_2(x)$ ($0 \leq f_1(x) \leq f_2(x)$) egri chiziqlar va $x = a$, $y = b$ to'g'ri chiziqlar bilan chegaralangan egri chiziqlar bilan chegaralangan figuraning Ox va Oy o'qlari atrofida aylanishidan hosil bo'lgan jism hajmi mos ravishda quyidagi


$$V_x = \rho \int_a^b (f_2^2(x) - f_1^2(x)) dx$$

$$V_y = 2\pi \int_a^b x(f_2(x) - f_1(x))dx$$

formula bilan hisoblandi.

15–masala. Quyidagi chiziqlar bilan chegaralangan shaklni Ox o'qi atrofida aylantirishdan hosil bo'lgan jismning hajmi toping.

$$y = 2x - x^2, \quad y = -x + 2.$$



Echim:

Ox o'q atrofida aylantirishdan hosil bo'lgan figuraning hajmi:

$$V = \pi \int_a^b y^2 dx.$$

Natijada:

$$y = 2x - x^2 = -x + 2 \Rightarrow x_1 = 1; \quad x_2 = 2.$$

Sirtning hajmi:

$$V = V_1 - V_2,$$

bunda

$$\begin{aligned}
 V_1 &= \rho \int_1^2 (2x - x^2)^2 dx = \rho \int_1^2 (4x^2 - 4x^3 + x^4) dx = \rho \left[\frac{4x^3}{3} - 4\frac{x^4}{4} + \frac{x^5}{5} \right]_1^2 = \\
 &= \rho \left[\frac{4 \cdot 8}{3} - 4\frac{16}{4} + \frac{32}{5} - \left(\frac{4}{3} - 4\frac{1}{4} + \frac{1}{5} \right) \right] = \rho \left[\frac{32}{3} - 16 + \frac{32}{5} - \frac{4}{3} + 1 - \frac{1}{5} \right] = \\
 &= \rho \left[\frac{28}{3} - 15 + \frac{31}{5} \right] = \rho \times \frac{28 \times 5 - 15 \times 15 + 31 \times 3}{15} = \rho \times \frac{8}{15}
 \end{aligned}$$

$$\begin{aligned}
 V_2 &= \rho \int_1^2 (-x + 2)^2 dx = \rho \int_1^2 (4 - 4x + x^2) dx = \rho \left[4x - 4\frac{x^2}{2} + \frac{x^3}{3} \right]_1^2 = \\
 &= \rho \left[8 - 8 + \frac{8}{3} - \left(4 - 2 + \frac{1}{3} \right) \right] = \rho \left[\frac{8}{3} - 2 - \frac{1}{3} \right] = \frac{1}{3} \rho.
 \end{aligned}$$

Demak,

$$V = V_1 - V_2 = \rho \times \frac{8}{15} - \rho \times \frac{1}{3} = \rho \times \frac{8 - 5}{15} = \frac{3}{15} \rho = \frac{1}{5} \rho.$$

15–masala. Quyidagi chiziqlar bilan chegaralangan shaklni (1-16 variantlarda) Ox o'qi atrofida, (17-30 variantlarda) Oy o'qi atrofida aylantirishdan hosil bo'lgan jismning hajmi toping.

1. $y = -x^2 + 5x - 6$, $y = 0$.
2. $2x - x^2 - y = 0$, $2x^2 - 4x + y = 0$.
3. $y = 3\sin x$, $y = \sin x$, $0 \leq x \leq \rho$.
4. $y = 5\cos x$, $y = \cos x$, $x = 0$, $x^3 = 0$.
5. $y = \sin^2 x$, $x = \rho/2$, $y = 0$.
6. $x = \sqrt[3]{y - 2}$, $x = 1$, $y = 1$.

$$7. y = xe^x, \quad y = 0, \quad x = 1.$$

$$8. y = 2x - x^2, \quad y = -x + 2, \quad x = 0.$$

$$9. y = e^{1-x}, \quad y = 0, \quad x = 0, \quad x = 1.$$

$$10. y = x^2, \quad y^2 - x = 0.$$

$$11. x^2 + (y - 2)^2 = 1.$$

$$12. y = 1 - x^2, \quad x = 0, \quad x = \sqrt{y - 2}, \quad x = 1$$

$$13. y = x^2, \quad y = 1, \quad x = 2.$$

$$14. y = x^3, \quad y = \sqrt{x}.$$

$$15. y = \sin \frac{\rho x}{2}, \quad y = x^2.$$

$$16. y = \arccos \frac{x}{3}, \quad y = \arccos x, \quad y = 0.$$

$$17. y = \arcsin \frac{x}{5}, \quad y = \arcsin x, \quad y = \frac{\rho}{2}.$$

$$18. y = x^2, \quad x = 2, \quad y = 0.$$

$$19. y = x^2 + 1, \quad y = x, \quad x = 0, \quad x = 1.$$

$$20. y = \sqrt{x - 1}, \quad y = 0, \quad y = 1, \quad x = 0,5.$$

$$21. y = \ln x, \quad x = 2, \quad y = 0.$$

$$22. y = (x - 1)^2, \quad y = 1.$$

$$23. y^2 = x - 2, \quad y = 0, \quad y = x^3, \quad y = 1.$$

$$24. y = x^3, \quad y = x^2.$$

$$25. y = \arccos \frac{x}{5}, \quad y = \arccos \frac{x}{3}, \quad y = 0.$$

$$26. y = \arcsin x, \quad y = \arccos x, \quad y = 0.$$

$$27. y = x^2 - 2x + 1, \quad x = 2, \quad y = 0.$$

28. $y = x^3$, $y = x$.

29. $y = \arccos x$, $y = \arcsin x$, $x = 0$.

30. $y = (x - 1)^2$, $x = 0$, $x = 2$, $y = 0$.

Javoblar. 15.1 $\frac{p}{30}$; 15.2 $\frac{16p}{5}$; 15.3 $4p^2$; 15.4 $4p$; 15.5 $\frac{3p^2}{16}$; 15.6 $\frac{44p}{7}$; 15.7 $p \times \frac{e^2 - 1}{4}$;

15.8 $\frac{9p}{5}$; 15.9 $2p \times (e^2 - 1)$; 15.10 $\frac{3p}{10}$; 15.11 $4p^2$; 15.12 $5p$; 15.13 $\frac{26p}{5}$; 15.14 $\frac{5p}{14}$;

15.15 $0,3p$; 15.16 $19,739$; 15.17 $6p^2$; 15.18 $8p$; 15.19 $\frac{5p}{6}$; 15.20 $\frac{97p}{60}$;

15.21 $p \times \frac{4}{e} \ln 2 - \frac{3}{2}$; 15.22 $\frac{8p}{3}$; 15.23 $\frac{24p}{5}$; 15.24 $\frac{p}{10}$; 15.25 $4p^2$; 15.26 $\frac{p}{2}$;

15.27 $\frac{7p}{6}$; 15.28 $\frac{4p}{15}$; 15.29 $\frac{p}{2} + \frac{p^2}{4}$; 15.30 $\frac{4p}{3}$.

Elementar funksiyalarning hosilalari jadvali

$u = u(x)$, $v = v(x)$ funksiyalar differensiallanuvchi funksiyalar bo'lsin.

$$1. y = C, y' = 0.$$

$$2. y = u + v + w, y' = u' + v' + w'$$

$$3. y = Cu, y' = Cu'$$

$$4. y = uv, y' = uv'$$

$$5. y = u^n, y' = nu^{n-1}u'$$

$$6. y = \frac{u}{v}, y' = \frac{u'v - uv'}{v^2}.$$

$$7. y = a^u, y' = a^u \ln a \cdot u'$$

$$8. y = e^u, y' = e^u u'$$

$$9. y = \ln u, y' = \frac{u'}{u}, u > 0$$

$$10. y = \log_a u, y' = \frac{u'}{u} \log_a e, u > 0$$

$$11. y = \sin u, y' = u' \cos u.$$

$$12. y = \cos u, y' = -u' \sin u.$$

$$13. y = \operatorname{tg} u, y' = \frac{u'}{\cos^2 u}.$$

$$14. y = \operatorname{ctg} u, y' = -\frac{u'}{\sin^2 u}$$

$$15. y = \arcsin u, y' = \frac{u'}{\sqrt{1-u^2}}$$

$$16. y = \arcsin u, y' = -\frac{u'}{\sqrt{1-u^2}}.$$

$$17. y = \operatorname{arctg} u, y' = \frac{u'}{1+u^2}.$$

$$18. y = \operatorname{arcctg} u, y' = -\frac{u'}{1+u^2}.$$

$$19. y = f(u), u = u(x), y' = f'_u(u) \cdot u'.$$

$$20. x = x(t), y = y(t), y' = \frac{y'_t}{x'_t}.$$

Asosiy integrallar jadvali

$$\text{I. } \int x^m dx = \frac{x^{m+1}}{m+1} + C, \quad (m \neq -1).$$

$$\text{II. } \int \frac{dx}{x} = \ln |x| + C.$$

$$\text{III. } \int \sin x dx = -\cos x + C.$$

$$\text{IV. } \int \cos x dx = \sin x + C.$$

$$\text{V. } \int a^x dx = \frac{a^x}{\ln a} + C.$$

$$\text{VI. } \int e^x dx = e^x + C.$$

$$\text{VII. } \int \frac{dx}{\cos^2 x} = \operatorname{tg} x + C.$$

$$\text{VIII. } \int \frac{dx}{\sin^2 x} = -\operatorname{ctg} x + C.$$

$$\text{IX. } \int \frac{dx}{1+x^2} = \operatorname{arctg} x + C.$$

$$\text{X. } \int \frac{dx}{\sqrt{1-x^2}} = \arcsin x + C.$$

$$\text{XI. } \int \operatorname{tg} x dx = -\ln |\cos x| + C.$$

$$\text{XII. } \int \operatorname{ctg} x dx = \ln |\sin x| + C.$$

$$\text{XIII. } \int \frac{dx}{a^2 + x^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C.$$

$$\text{XIV. } \int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin \frac{x}{a} + C.$$

$$\text{XV. } \int \frac{dx}{\sqrt{x^2 \pm a^2}} = \ln \left| x + \sqrt{x^2 \pm a^2} \right| + C.$$

$$\text{XVI. } \int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \ln \left| \frac{a+x}{a-x} \right| + C.$$

FOYDANILGAN ADABIYOTLAR RO`YXATI.

1. Azlarov.T.A., Mansurov H. Matematik analiz. 1 va 2-qismlar-T., "O`qituvchi" 1986, 1989.

2. Sa'dullayev A., Mansurov H., Xudoyberganov G., Vorisov A., G'ulomov R. *Matematik analiz kursidan misol va masalalar to'plami, 1 va 2-qismlar*.-T., "O'zbekiston", 1993, 1995.
3. B.A.Shoimqulov, T.T.To'ychiyev, D.X.Djumaboyev «*Matematik analizdan mustaqil ishlar*», T., «O'zbekiston faylasuflari milliy jamiyati» nashriyoti, 2008
4. Зорич. А. Математический анализ, Т. 1-2, М. «Наука», 1981.
5. Ильин В.А., Садовничий В.А., Сендов Б.Х. Математический анализ. Т. 1-2, М., «Наука», 1979.
6. Фихтенгольц Г.М. Курс дифференциального и интегрального исчисления, Т. 1-3, М. «Физ-матгиз», 1962.
7. Tuychiev T.T., Djumaboyev D.X. *Matematik analiz fanidan 1 va 2-kurs talabalari uchun laboratoriya ishlari*. T., «Universitet», 2003.
8. Кузнецов Л.А. Сборник заданий по высшей математике-М., "Высшая школа", 1982.
9. Демидович Б.П. Сборник задач и упражнений по математическому анализу.-М., "Наука", 1972.
10. Кудрявцев Л.Д., Кутасов А.Д., Шеглов В.И., Шабунин М.И.. Сборник задач по математическому анализу, Т. 1, 2-М., "Наука", 1984, 1986.
11. Виноградова И.А., Олехник С.Н., Садовничий В.А. Задачи и упражнения по математическому анализу, Т. 1.-Издательство Московского Университета, 1988.
12. Бутузов В.Ф., Крутицкая Н.Ш., Медведев Г.Н., Шишкин А.А. Математический анализ в вопросах и задачах, Т. 1, 2-М., "Высшая школа", 1984, 1988.
13. Бруй И.Н., Гаврилюк А.В. и др. Лабораторный практикум по математическому анализу -Минск, "Высшая школа", 1991.
14. К.Н.Лунгу, Д.Т.Письменный, С.Н.Федин, Ю.А.Шевченко. Сборник задач по высшей математике. М. «Айрис пресс», 2001.
15. Н.Ш.Кремер, Б.А.Путко, И.М.Тришин, М.Н.Фридман. «Высшая математика для экономистов», М., ЮНИТИ, 2004.

So'z boshi	3
I BOB. Analitik geometriya	4
Vektorni bazis bo'yicha yoyilmasi	4
Vektorlarning kollinearligi	9
Vektorlar orasidagi burchak	11
Parallelogramning yuzi	13
Vektorlar komplanarligi	15
Tetraedrning balandligi va hajmi	17
Nuqtadan tekislikkacha bo'lgan masofa	22
Normal vektori berilgan tekislik tenglamasi	25
Tekisliklar orasidagi burchak	27
Bir xil uzoqlikda yotgan nuqtaning koordinatalari	30
To'g'ri chiziqning kononik tenglamasi	35
To'g'ri chiziq va tekislikning kesishish nuqtasi	37
To'g'ri chiziqqa nisbatan nuqtalarning simmetrikligi	39
II BOB. Limitlar nazariyasi	42
Sonli ketma-ketlikning limiti	46
Funktsiyaning limiti	60
III BOB. Funktsiyaning hosilasi	84
Funktsiyaning differentsiali	92
Differentsiallashning asosiy qoidalari	98
Murakkab funktsiyaning hosilasi	101
Funktsiyaning hosilasi	104
Parametrik berilgan funktsiyalarning hosilasi	128
Egri chiziqqa nuqtadan o'tkazilgan urinma va normal tenglamasi	131
Yuqori tartibli hosilalar	135
IV BOB. Grafiklar	147
Funktsiyaning grafigi	147
Funktsiyaning eng katta va eng kichik qiymatlari	149
Funktsiyaning asimptotalari	153
Funktsiyalarni tekshirish va grafiklarini chizish	156
V BOB. Aniqmas integral	167
Aniqmas integralning ta'rifi va xossalari	167
Integrallashning asosiy usullari	169
Yoyish (integral ostidagi ifodani yoyib integrallash) usuli	169

Bevosita integrallash usuli	170
Bo'laklab integrallash usuli	174
Kvadrat uchhadni o'z ichiga olgan funktsiyalarni integrallash	176
Ratsional kasrlarni integrallash	180
Noma'lum koeffitsientlar usuli	185
Maxrajning ildizlari kompleks va karrali	189
$\int x^m (a + bx^n)^p dx$ differentsial binomial integrali	194
VI BOB. Aniq integral	197
Aniq integralda o'zgaruvchini almashtirish	197
Bo'laklab integrallash	200
$R(\sin x, \cos x)dx$ ko'rinishdagi integrallar	203
$\int \sin^m x \cos^n x dx$ ko'rinishdagi integrallar	209
$\int R\left(x, \frac{ax+b}{cx+d}, \frac{ax+b}{cx+d}, \dots, \frac{ax+b}{cx+d}\right) dx$ ko'rinishdagi integrallar	212
$R(x, \sqrt{a^2 \pm x^2})$ va $R(x, \sqrt{x^2 - a^2})$ ko'rinishdagi integrallar.	215
Yassi figuralar yuzlarini hisoblash	217
Figuraning yuzini qutb koordinalar sistemasida hisoblash	224
Yoy uzunligini hisoblash	227
Parametrik ko'rinishda berilgan egri chiziqli yoyining uzunligi	230
Qutb koordinalari sistemasida berilgan egri chiziqli yoyining uzunligi	233
Aylanma jism sirining yuzi	235
Jismlarning hajmini ularning ko'ndalang kesimlari bo'yicha hisoblash	236
To'g'ri burchakli dekart koordinalar sistemasida aylanma jism hajmi	241
Ilovalar	246
Foydalanilgan adabiyotlar ro'yxati	249

